

Correlation of Self-Efficacy, Action Planning, and Coping Planning to ARV Adherence in HIV Patients in Ternate, Indonesia

Nani Supriyatni^{1*}, Lutfi Agus Salim², Arief hargono³, Tati Sumiati⁴, Tutik Lestari⁵, Andani⁶

^{1,4,5,6}Faculty of Public Health, Universitas Muhammadiyah Maluku Utara

^{2,3} Department of Epidemiology and Biostatistics Faculty of Public Health, Universitas Airlangga

How to Cite: Supriyatni, N., Agus Salim, L., Hargono, A., Sumiati, T., Lestari, T., & Andiani. Correlation of Self-Efficacy, Action Planning, and Coping Planning to ARV Adherence in HIV Patients in Ternate, Indonesia. *Jurnal Kesehatan*, 18(3), 375-386. <https://doi.org/10.23917/jk.v18i3.12648>

Article Information

Article History:

Submission: 03 September 2025

Revision: 26 September 2025

Reception: 20 October 2025

Keywords: Adherence, ARV, action planning, coping planning, self-efficacy

ABSTRACT

Introduction: HIV-positive patients require adherence to Antiretroviral therapy (ART) to control their disease. However, in Ternate City, ARV adherence remains low due to side effect fears, stigma, depression, and limited health service access. Therefore, this study analyzes the correlation between self-efficacy, action planning, and coping planning with ARV adherence in HIV-positive patients in Ternate City, North Maluku. **Method:** An observational, cross-sectional study used stratified random sampling of HIV-positive patients at five service centers, with data analyzed using the Spearman test. **Results:** Most respondents were aged 19-49 (94.9%), male (69.1%), held a high school education (57.3%), were self-employed (24.2%), and earned under 1 million rupiah per month (33.1%). Furthermore, self-efficacy ($p=0.000$; $r=0.404$), action planning ($p=0.000$; $r=0.399$), and coping planning ($p=0.000$; $r=0.265$) were significantly associated with ARV adherence. **Conclusion:** These findings suggest that self-efficacy, action planning, and coping planning benefit ARV adherence in HIV-positive patients.

Corresponding Authors: (*)

Prodi Kesehatan Masyarakat, Fakultas Ilmu Kesehatan, Universitas Muhammadiyah Universitas Muhammadiyah Maluku Utara, Jl. KH ahmad Dahlan no 100 kelurahan Sasa, Kec. Ternate Selatan, Maluku Utara 57169, Indonesia

Email: nanisupriyatni123@gmail.com

INTRODUCTION

HIV/AIDS is an infectious disease that has caused 40.4 million deaths since it was declared an epidemic. By the end of 2022, 39 million people worldwide were HIV-positive. Approximately 630,000 people worldwide had died from the disease (World Health Organization, 2023a). According to the WHO, HIV/AIDS is the second-leading cause of death among adolescents globally. It is also among the top ten causes of death in low-income countries. HIV/AIDS is projected to become a leading cause of disease burden by 2030 (Mathers and Loncar, 2006; World Health Organization, 2020)

Indonesia has the third-highest number of people living with HIV and new infections in the Asia-Pacific region, after India and China. Data collected by the World Health Organization in 2019 showed that 81% of people living with HIV knew their status, 67% were receiving antiretroviral therapy, and 59% had achieved HIV viral suppression without the risk of infecting others. In Indonesia, data shows 143,590 people living with HIV, with only 77,748 people, or 33.5%, receiving ARV therapy. This falls short of the ARV therapy coverage target of over 80%. 31,415 people have not received ARVs, 31,487 have lost follow-up, and 2,940 have discontinued ARV therapy. A preliminary study of Ternate City Health Office data, based on cumulative HIV/AIDS data, revealed that as of December 2022, 2,363 people were living with HIV, of whom 2,125 were eligible for ARV treatment, 764 had previously received ARV therapy, and 286 (37.4%) were still on ARV treatment. Additionally, 262 people living with HIV died, and 129 lost follow-up (Kementerian Kesehatan RI, 2022).

The high mortality rate among HIV/AIDS sufferers is influenced by factors such as late HIV diagnosis, low awareness of serostatus, low CD4 count, high viral load, difficulty maintaining treatment, advanced disease stage, poor adherence to diet, low body weight, anemia, low socioeconomic status, late initiation of ARV, and poor adherence to antiretroviral therapy (ARV) (Colecraft, 2008; Bhattacharya, Dubey and Sharma, 2011; Arrivé *et al.*, 2012; Hickson and Mayers, 2020; Ramaiya *et al.*, 2020; Christopher *et al.*, 2021). Currently, there is no fully effective cure for HIV infection. However, with access to effective HIV prevention, diagnosis, treatment, and care—including treatment of opportunistic infections—HIV has become a manageable chronic condition, allowing people living with HIV to lead long, healthy lives (World Health Organization, 2023b).

Effective treatment with antiretroviral drugs (ARVs) can control the infection and disease progression, enabling HIV-positive individuals to enjoy healthy and productive lives. ARVs work by reducing HIV replication and preventing infection of new cells, which also improves immune system function. Consequently, ARV therapy positively impacts the quality of life and survival of those with HIV (Seyoum *et al.*, 2017; Nuraidah *et al.*, 2022). Several studies on HIV-positive patients indicate that adherence to ARV therapy in Indonesia remains low (Safira, Lubis and Fahdhy, 2018; Martiana, Waluyo and Yona, 2019; Nurfalah, Yona and Waluyo, 2019). Non-adherence to ARV therapy is dangerous because it can lead to increased viral load, disease progression, drug resistance, and a higher risk of transmitting resistant HIV to others, especially among sexually active adolescents (Attia *et al.*, 2009; Adejumo *et al.*, 2015).

Various barriers impede HIV-positive patients from adhering to ARV therapy, including feeling healthy (not yet experiencing opportunistic infections), fear of side effects, reluctance to take medication, boredom with treatment, stigma, discrimination, depression, limited access to healthcare facilities, transportation costs, and therapy expenses (Horne *et al.*, 2013; Bezabhe *et al.*, 2014; Wardhani and Yona, 2021; Purwaningsih, Asmoro and Prastiwi, 2022; Hutahaeon, Stutterheim and Jonas, 2023). These barriers are largely driven by low self-efficacy and a lack of planning skills in HIV-positive individuals. Self-efficacy is the belief in one's ability to adhere to a treatment plan despite challenges such as side effects, disruptions to daily life, barriers to treatment, depression, and lack of social support. Multiple studies have demonstrated that self-efficacy influences health behavior changes (Isa *et al.*, 2017; Tharek *et al.*, 2018; Dzerounian *et al.*, 2022). Conversely, having good planning skills can enhance behavioral control. Planning serves as a self-regulation strategy to anticipate and manage challenges related to treatment.

There is an urgent need to address these barriers, as improving adherence to antiretroviral therapy is critical for the health outcomes of HIV-positive individuals and for

controlling the spread of HIV. The limitations of previous studies are that the last research from the research location did not cover all health services in the city of Ternate, limited to referral hospitals only, the geographical location is not accessible to all PLHIV, and the previous research method of the variables studied was limited to the threat of disease on adherence to taking ARV medication. For this reason, this study expands the research location to all HIV Support and Treatment (PDP) service locations in Ternate City. This study expands the research variables that show the influence of planning on ARV therapy adherence behavior. The purpose of this study is to study the correlation of self-efficacy, action planning, and coping planning with ARV therapy adherence in HIV-positive patients in Ternate, North Maluku Province.

LITERATURE REVIEW

Human Immunodeficiency Virus (HIV) is a virus that causes a decline in the human immune system, while Acquired Immune Deficiency Syndrome (AIDS) is a further development in people infected with HIV. "Acquired" refers to an infectious disease that is not hereditary but develops after direct contact with an HIV-infected person (Kristiono and Astuti, 2019).

HIV (Human Immunodeficiency Virus) is a virus that attacks the immune system, leaving the body vulnerable to various diseases. Acquired Immune Deficiency Syndrome (AIDS) can be defined as a collection of symptoms or diseases caused by a decline in the immune system due to HIV infection, specifically affecting T lymphocytes and reducing the number of CD4 cells, which are responsible for fighting infection. AIDS is the final stage of HIV infection (Kemenkes, 2020). The syndrome arises from a reduction in the body's immune system (CD4) that occurs approximately 5–10 years after HIV infection, progressing to AIDS, characterized by a CD4 count of less than 200 cells per μL of blood as the threshold criterion (Keputusan Menteri Kesehatan RI, 2019; Kristiono and Astuti, 2019).

Antiretrovirals (ARVs) are drugs that inhibit the replication of the Human Immunodeficiency Virus (HIV). ARV therapy is the most clinically successful strategy to date. Antiretroviral therapy (ART) means treating HIV infection with drugs. These drugs (called ARVs) do not kill the virus but can slow its growth, slowing the viral growth rate and thus the course of HIV disease. Because HIV is a retrovirus, these drugs are commonly referred to as antiretroviral therapy (Nurihwani, 2017).

"Adherence" is an English term that refers to a client's compliance with treatment (on time, at the correct dose, and taken in the right way). Treatment adherence can be defined as the patient's behavior in following all the advice and instructions recommended by medical professionals, such as doctors and pharmacists, regarding everything that needs to be done to achieve treatment goals, one of which is adherence to taking medication. This is one of the successes of the treatment carried out (Kementrian Kesehatan, 2019).

Non-adherence to ARV therapy is very dangerous because it can lead to increased viral load, disease progression, drug resistance, and an increased risk of transmitting resistant HIV to others in sexually active adolescents (Adejumo *et al.*, 2015)

Self-efficacy has been shown to have a stronger influence on ARV medication adherence in HIV-positive patients than other factors such as sociodemographics, duration of therapy, disease condition, interpersonal correlations such as social support, and other potentially related factors such as drug use (Colbert, Sereika and Erlen, 2013; Mi *et al.*, 2020).

Action planning links goal-directed behavior to specific environmental cues by clarifying when, where, and how to act. This process can unintentionally trigger action and encourages individuals to follow their plans to reach their goals (Sniehotta *et al.*, 2005; Kwasnicka *et al.*, 2013). Research by Wee and Dilon (2022) found that action planning relates to increased physical activity. It serves as a bridge, helping turn intentions into actions

(Lippke, Ziegelmann and Schwarzer, 2005; Norman and Conner, 2005; Scholz *et al.*, 2008; Schwarzer and Luszczynska, 2008)

Coping planning is a self-regulation strategy that focuses on obstacles. This represents the mental connection between the anticipated risk situation and the appropriate coping response. Coping planning is a detailed plan to anticipate potential difficulties or obstacles that may arise when implementing an intention, consisting of a series of activities to identify unwanted obstacles or disruptions and how to overcome them (Wee and Dillon, 2022).

METHOD

The type of research is observational with a cross-sectional design. The research was conducted in five HIV service centers in Ternate City, North Maluku Province, namely Chasan Boesoirie Regional General Hospital, Siko Care Health Center, City Health Center, BLUD Kalumpang Health Center, and Kalumata Health Center. The population in this study was all HIV patients who had participated in the ARV treatment program in five HIV service centers in Ternate City, totaling 318 respondents. The minimum sample in this study was 178 PLHIV using a stratified random sampling technique. The sampling steps were (1) recording the number of PLHIV currently undergoing ARV treatment in five HIV service centers in Ternate City, (2) from the number of PLHIV in each service center, a representative sample was taken randomly, with 132 samples of PLHIV in Chasan Boesoirie Regional General Hospital, 10 samples in Siko Health Center, 10 samples in City Health Center, 11 samples in Kalumpang Health Center, and 15 samples in Kalumata Health Center. The inclusion criteria for this study were: HIV-positive patients registered in the medical records of Ternate City HIV services, willing to be interviewed, and undergoing ARV treatment. Respondents from outside the island and who were difficult to contact, as well as those who had moved, were excluded.

Data collection for this study was conducted over four months, from August to December 2023. Data collection was conducted using structured interviews guided by a questionnaire. The questionnaire's validity and reliability were tested on a sample of 20 people living with HIV. A calculated *r* value (correlation coefficient) of more than 0.444 indicated the instrument's validity. A Cronbach's alpha value of more than 0.6 indicated its reliability. The questionnaire consisted of eight questions on adherence, 12 questions on self-efficacy, three questions on action planning, and three questions on coping planning. Statistical analysis using Spearman's rho test was conducted to determine the relationship between self-efficacy, action planning, and coping planning and ARV adherence.

All participants in the study agreed to participate freely, as evidenced by the completion of an informed consent form. This research has passed the ethical review of Airlangga University under reference number 954/HRECC.FODM/VIII/2023.

RESULTS AND DISCUSSION

Statistical analysis revealed a significant relationship between self-efficacy, action planning, and coping planning and ARV therapy adherence in HIV-positive patients. The relationship between these variables was positive, indicating that the higher the self-efficacy, action planning, and coping planning, the higher the adherence of HIV-positive patients to ARV therapy. The results of the analysis can be seen in Table 1.

Socio-demographic factors consist of age, gender, education, occupation, income, and ARV. Based on the research results, most of the respondents were HIV patients with the age of 19-49 years (94.9%), male 123 (69.1%), high school education 102 (57.3%), the employment status of the majority of respondents was self-employed, including

housewives, traders, motorcycle taxi drivers, fishermen, massage parlors or beauty salons and the average income of respondents was low <1 million rupiah per month (33.1%) (Table 1).

Table 1. Frequency Distribution of Socio-Demographic Factors

Variable	Categories	Frequency (n)	Percentage (%)
Age (years)	10 - 18	1	6
	19 - 49	169	94.6
	>50	8	4.5
Gender	Males	123	69.1
	Female	55	30.9
Education	Non-School	2	1.1
	Elementary School	2	1.1
	Junior High School	18	10.1
	High School	102	57.1
	College	54	30.1
Occupation	Not working	37	20.8
	Civil Servant	29	16.3
	Private Employees	31	17.4
	Wireless	43	24.2
	Other	38	21.3
Revenue (Rupiahs*)	<1 million	59	33.1
	1 million - 2.5 million	52	29.2
	2 million - 5 million	53	29.8
	>5 million	14	7.9

Table 2. Frequency Distribution of Self-efficacy, Action planning, Coping planning, ARV compliance

Variable	Categories	Frequency (n)	Percentage (%)
Self Efficacy	Low	23	12.9
	Medium	120	67.4
	High	35	19.7
Action Planning	Low	17	9.6
	Medium	98	55.1
	High	63	35.4
Coping Planning	Low	38	21.3
	Medium	98	55.1
	High	42	23.6
ARV Adherence	Low	23	15.9
	Medium	80	44.9
	High	75	42.1

The results in Table 2 above show that the majority of respondents had moderate self-efficacy (n = 120, 67.4%). Only 35 patients (19.7%) had high self-efficacy. This indicates that only a few respondents had good self-efficacy. The average self-efficacy score of respondents was 39.52 out of a maximum score of 48, with a standard deviation of 6.26. The majority of respondents reported moderate action planning (n = 98, 55.1%). 63 patients (35.4%) had high action planning. Although not as low as the self-efficacy variable, respondents' action planning was still low, indicating that more than half of the respondents did not have good action planning skills. The average action planning score of respondents was 10.15 out of a maximum score of 12 and had a standard deviation of 1,777. The majority of respondents had moderate coping planning (n=98, 55.1%). Only 42 (23.6%)

respondents had high coping planning. This indicates that only a few respondents had good coping planning. The average score for coping planning was 9.35 out of a maximum score of 12 and had a standard deviation of 1,817. The majority of respondents had moderate adherence to ARV therapy (n=80, 44.9%). Only 75 (42.1%) respondents had high adherence to ARV therapy. This indicates that more than half of the respondents still did not have good adherence. The average score for respondent adherence was 28.61 out of a maximum score of 32 and had a standard deviation of 4,717.

Table 3. Correlation between self-efficacy, action planning, and coping planning with ARV adherence in HIV-positive patients

Variabel	Mean	SD*s	Minimum	Maximum	p	r
Self-efficacy	39.52	6.263	12	48	0.000	0.404
Action planning	10.15	1.777	3	12	0.000	0.399
Coping planning	9.35	1.817	3	12	0.000	0.265
ARV Adherence	28.61	4.717	8	32		

*SD: Standard Deviation

Table 3 above shows the results of the Spearman test, which found that self-efficacy (p=0.000), action planning (p=0.000), and coping planning (p=0.000) were significantly correlated with ARV therapy adherence in HIV-positive patients. Furthermore, the correlation coefficient was positive, indicating that the greater the self-efficacy (p=0.404), action planning (p=0.399), and coping planning (p=0.265), the greater the ARV therapy adherence in HIV-positive patients in Ternate.

Self-efficacy is often associated with positive health behaviors, such as engaging in recommended health behaviors, improving communication with healthcare providers, adapting to the disease, and treatment adherence (Oshotse *et al.*, 2018). This study found that self-efficacy was significantly associated with ARV therapy adherence in HIV-positive patients. These findings align with other studies (Colbert, Sereika and Erlen, 2013; Brown *et al.*, 2015; Machtinger *et al.*, 2015; Archiopoli *et al.*, 2016; Peng *et al.*, 2020). Self-efficacy in ARV treatment refers to a person's belief that they will be able to take ARVs as recommended. Self-efficacy has been shown to have a stronger influence on ARV therapy adherence in HIV-positive patients compared to other factors such as sociodemographics, duration of therapy, disease condition, interpersonal correlations such as social support, and other potentially related factors such as drug use (Colbert, Sereika and Erlen, 2013). Rather than seeing the difficulties and side effects of ARV therapy, someone with high treatment self-efficacy is more likely to see the benefits of treatment and try to prevent potential negative impacts (Mi *et al.*, 2020). Furthermore, self-efficacy can act as a protective barrier to reduce the negative impact of stigma on quality of life and is known to be a facilitator of ARV adherence (Zhang *et al.*, 2016; Zhou *et al.*, 2018). Therefore, self-efficacy-related interventions in HIV-positive patients, such as measuring self-efficacy levels, can be an early detection method for poor ARV therapy adherence. Kim *et al.* stated that measuring self-efficacy during routine visits to the clinic by HIV-positive patients can be an early detection method for future non-adherence to ARV therapy. Furthermore, interventions aimed at improving self-efficacy can improve adherence (Kim *et al.*, 2017).

Action planning is a detailed specification of when, where, and how to act in accordance with one's intentions. Action planning encourages a person to act according to a predetermined plan to achieve a goal (Sniehotta *et al.*, 2005). The results of this study indicate that action planning is significantly related to ARV therapy adherence in HIV-positive patients. Individuals with an action plan will find it easier to adhere to ARV therapy as recommended. This includes creating a specific schedule for taking ARVs,

organizing treatment-related needs such as finding a health facility or obtaining medication, planning questions to ask health professionals during consultations, and planning the best strategy for visiting health facilities, such as scheduling clinic visits and transportation. With careful planning, barriers to ARV therapy can be avoided.

To our knowledge, there has been no research examining the effect of action planning on ARV therapy adherence in HIV-positive patients. However, research on the effects of action planning has been conducted for other health behavior changes, such as regularly taking supplements, exercising, quitting smoking, and adopting a healthy (Godinho *et al.*, 2014; Hsieh, Kanda and Fujii, 2019; Wee and Dillon, 2022). In their study, Wee and Dillon (2022) found that action planning is associated with increased physical activity. By conducting action planning, a person can design the "how," "when," and "where" to achieve the desired behavior. This can remind them of the important steps needed to improve positive behavior (Wee and Dillon, 2022). Action planning serves as a bridge to translate a plan into action. Therefore, action planning can only be implemented after a person has the intention to do something (Ziegelmann, Lippke and Schwarzer, 2006). Coping planning is a detailed plan to anticipate difficulties or obstacles that may arise when implementing an intention. It consists of a series of activities that identify unwanted obstacles or disruptions and how to overcome them (Scholz *et al.*, 2008; Wee and Dillon, 2022). In our research, we found a significant relationship between coping planning and ARV therapy adherence in HIV-positive patients. Many obstacles contribute to non-adherence to ARV therapy (e.g., forgetting to take medication), and individuals with good coping planning will find potential solutions to address these obstacles (e.g., setting a medication alarm). Having a coping plan does not ensure a person's adherence to ARV therapy, but it does prevent someone who is already on ARV therapy from stopping due to obstacles such as reluctance to take medication or fear of side effects. This coping plan is more useful for maintaining behavior (Sniehotta *et al.*, 2005; Ziegelmann, Lippke and Schwarzer, 2006; Kwasnicka *et al.*, 2013). No research has examined the relationship between coping planning and ARV therapy adherence, but the influence of coping planning on other health behaviors has been extensively studied (Kwasnicka *et al.*, 2013). Research by Lin *et al.* found a strong relationship between coping planning and epilepsy treatment adherence. This indicates that patient anticipation of adherence barriers is a crucial factor, necessitating interventions to improve adherence by encouraging patients to overcome these barriers (Lin, Updegraff and Pakpour, 2016).

Self-efficacy, action planning, and coping planning all play a crucial role in helping individuals adhere to ARV therapy. Sometimes, individuals develop an action plan to adhere to therapy but fail due to a lack of ability to resist temptation, be assertive in the face of peer pressure, or overcome a temporary lack of motivation (Araújo-Soares, McIntyre and Sniehotta, 2009). Scholz *et al.* demonstrated how failure or mastery in implementing an action plan can influence self-efficacy for carrying it out. Self-efficacy in action planning is a key determinant of successful change. Those who adhere to action planning with coping planning are more likely to resist temptation, implement their plans, and experience mastery (Scholz *et al.*, 2007).

Physical activity-related interventions involving action planning and coping planning are more effective for behavior change and barrier management than other interventions. The combination of the two is considered more effective than interventions solely focused on action planning (Gollwitzer and Sheeran, 2006; Dombrowski *et al.*, 2012). On the other hand, Kwasnicka *et al.* argue that planning can serve as an effective behavior change technique, similar to physicians encouraging patients to adhere to treatment, but such efforts do not appear to be as effective in interventions that do not provide any support in formulating plans (Kwasnicka *et al.*, 2013).

Research on the influence of action planning and coping planning on ARV adherence in HIV-positive patients is limited, so this study was compared with research on other health behaviors. Therefore, the results of this study cannot be compared with findings in other areas related to ARV adherence. Further research on the relationship between action planning, coping planning, and ARV adherence, as well as related interventions, is needed.

CONCLUSION

Self-efficacy, action planning, and coping planning are significantly associated with ARV therapy adherence in HIV-positive patients in Ternate, Indonesia. Self-efficacy boosts patients' confidence in ARV adherence, action planning helps patients realize their intention to adhere to ARV, while coping planning helps patients maintain ARV adherence by finding solutions to overcome emerging obstacles. The recommendation from this study is for interventions to provide training on self-efficacy, action planning, and coping planning, which can be beneficial for maintaining ARV adherence in HIV patients in Ternate.

ACKNOWLEDGEMENT

Thank you to the Muhammadiyah University of North Maluku for providing research funding assistance, and thank you to the research institutions that have provided research permits, namely Chasan Boesoirie Regional Hospital, Siko Care Health Center, City Health Center, BLUD Kalumpang Health Center, and Kalumata Health Center.

REFERENCES

- Adejumo, O.A. *et al.* (2015) 'Contemporary issues on the epidemiology and antiretroviral adherence of HIV-infected adolescents in sub-Saharan Africa: A narrative review', *Journal of the International AIDS Society*, 18(1). doi:10.7448/IAS.18.1.20049.
- Araújo-Soares, V., McIntyre, T. and Sniehotta, F.F. (2009) 'Predicting changes in physical activity among adolescents: The role of self-efficacy, intention, action planning and coping planning', *Health Education Research*, 24(1), pp. 128–139. doi:10.1093/her/cyn005.
- Archiopoli, A. *et al.* (2016) 'Factors of interpersonal communication and behavioral health on medication self-efficacy and medication adherence', *AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV*, 28(12), pp. 1607–1614. doi:10.1080/09540121.2016.1192577.
- Arrivé, E. *et al.* (2012) 'HIV status disclosure and retention in care in HIV-infected adolescents on antiretroviral therapy (ART) in West Africa', *PLoS ONE*, 7(3), pp. 1–8. doi:10.1371/journal.pone.0033690.
- Attia, S. *et al.* (2009) 'Sexual transmission of HIV according to viral load and antiretroviral therapy: Systematic review and meta-analysis', *Aids*, 23(11), pp. 1397–1404. doi:10.1097/QAD.0b013e32832b7dca.
- Bezabhe, W.M. *et al.* (2014) 'Barriers and facilitators of adherence to antiretroviral drug therapy and retention in care among adult HIV-positive patients: A qualitative study from Ethiopia', *PLoS ONE*, 9(5). doi:10.1371/journal.pone.0097353.
- Bhattacharya, M., Dubey, A.P. and Sharma, M. (2011) 'Patterns of diagnosis disclosure and its correlates in HIV-infected north Indian children', *Journal of Tropical Pediatrics*, 57(6), pp. 405–411. doi:10.1093/tropej/fmq115.
- Brown, M.J. *et al.* (2015) 'Disclosure and Self-Efficacy among HIV-Positive Men Who Have Sex with Men: A Comparison between Older and Younger Adults', *AIDS Patient Care and STDs*, 29(11), pp. 625–633. doi:10.1089/apc.2015.0133.
- Christopher, S. *et al.* (2021) 'Children and Youth Services Review Disclosure of HIV / AIDS

- status to infected children in Ghana – A north-south comparison of barriers and enablers', *Children and Youth Services Review*, 122, p. 105753. doi:10.1016/j.chidyouth.2020.105753.
- Colbert, A.M., Sereika, S.M. and Erlen, J.A. (2013) 'Functional health literacy, medication-taking self-efficacy and adherence to antiretroviral therapy', *Journal of Advanced Nursing*, 69(2), pp. 295–304. doi:10.1111/j.1365-2648.2012.06007.x.
- Colecraft, E. (2008) 'HIV/AIDS: Nutritional implications and impact on human development', *Proceedings of the Nutrition Society*, 67(1), pp. 109–113. doi:10.1017/S0029665108006095.
- Dombrowski, S.U. *et al.* (2012) 'Identifying active ingredients in complex behavioural interventions for obese adults with obesity-related co-morbidities or additional risk factors for co-morbidities: a systematic review', *Health Psychology Review*, 6(1), pp. 7–32. doi:10.1080/17437199.2010.513298.
- Dzerounian, J. *et al.* (2022) 'Health knowledge and self-efficacy to make health behaviour changes: a survey of older adults living in Ontario social housing', *BMC Geriatrics*, 22(1), pp. 1–17. doi:10.1186/s12877-022-03116-1.
- Godinho, C.A. *et al.* (2014) 'Will is not enough: Coping planning and action control as mediators in the prediction of fruit and vegetable intake', *British Journal of Health Psychology*, 19(4), pp. 856–870. doi:10.1111/bjhp.12084.
- Gollwitzer, P.M. and Sheeran, P. (2006) 'Implementation Intentions and Goal Achievement: A Meta-analysis of Effects and Processes', *Advances in Experimental Social Psychology*, 38(06), pp. 69–119. doi:10.1016/S0065-2601(06)38002-1.
- Hickson, W. and Mayers, P.M. (2020) 'Understanding how young people become motivated to take their human immunodeficiency virus medication (Antiretroviral therapy) and how the need for adherence is communicated', *Health SA Gesondheid*, 25, pp. 1–9. doi:10.4102/hsag.v25i0.1458.
- Horne, R. *et al.* (2013) 'Understanding patients' adherence-related Beliefs about Medicines prescribed for long-term conditions: A meta-analytic review of the Necessity-Concerns Framework', *PLoS ONE*, 8(12). doi:10.1371/journal.pone.0080633.
- Hsieh, H.S., Kanda, Y. and Fujii, S. (2019) 'Incorporation of coping planning into the behavior change model that accounts for implementation intention', *Transportation Research Part F: Traffic Psychology and Behaviour*, 60(101), pp. 228–244. doi:10.1016/j.trf.2018.10.025.
- Hutahaean, B.S.H., Stutterheim, S.E. and Jonas, K.J. (2023) 'Barriers and Facilitators to HIV Treatment Adherence in Indonesia: Perspectives of People Living with HIV and HIV Service Providers', *Tropical Medicine and Infectious Disease*, 8(3). doi:10.3390/tropicalmed8030138.
- Isa, A. *et al.* (2017) 'Association between Self-Efficacy and Health Behaviour in Disease Control: A Systematic Review', *Global Journal of Health Science*, 10(1), p. 18. doi:10.5539/gjhs.v10n1p18.
- Kemenkes, P. (2020) 'Pusat Data dan Informasi Kementerian Kesehatan', *INFODATIN* [Preprint], (2442–7659). Available at: <http://www.depkes.go.id/resources/download/pusdatin/infodatin/InfodatinAIDS.pdf>.
- Kementerian Kesehatan RI (2022) 'Distribusi ODHIV yang di tes per Provinsi dapat dilihat pada grafik berikut ini.', *Laporan Eksekutif Perkembangan Hiv Aids Dan Penyakit Infeksi Menular Seksual (Pims) Triwulan I Tahun 2022* [Preprint].
- Kementerian Kesehatan, R. (2019) *Petunjuk Teknis Standar Pelayanan Kefarmasian di Rumah Sakit*.

- Keputusan Menteri Kesehatan RI (2019) 'Pedoman Nasional Pelayanan Kedokteran Tata Laksana HIV'. Available at: http://www.ghbook.ir/index.php?option=com_dbook&task=readonline&book_id=13650&page=73&chckhashk=ED9C9491B4&Itemid=218&lang=fa&tmpl=component%0Ahttp://www.albayan.ae%0Ahttps://scholar.google.co.id/scholar?hl=en&q=APLIKASI+PENGENA.
- Kim, M.H. *et al.* (2017) 'High self-reported non-adherence to antiretroviral therapy amongst adolescents living with HIV in Malawi: Barriers and associated factors', *Journal of the International AIDS Society*, 20(1), pp. 1–12. doi:10.7448/IAS.20.1.21437.
- Kristiono, N. and Astuti, I. (2019) *Mengenal Hiv Dan Aids*, <https://www.researchgate.net/publication/337049851>.
- Kwasnicka, D. *et al.* (2013) 'Does planning how to cope with anticipated barriers facilitate health-related behaviour change? A systematic review', *Health Psychology Review*, 7(2), pp. 129–145. doi:10.1080/17437199.2013.766832.
- Lin, C.Y., Updegraff, J.A. and Pakpour, A.H. (2016) 'The relationship between the theory of planned behavior and medication adherence in patients with epilepsy', *Epilepsy and Behavior*, 61, pp. 231–236. doi:10.1016/j.yebeh.2016.05.030.
- Lippke, S., Ziegelmann, J.P. and Schwarzer, R. (2005) 'Stage-specific adoption and maintenance of physical activity: Testing a three-stage model', *Psychology of Sport and Exercise*, 6(5), pp. 585–603. doi:10.1016/j.psychsport.2004.11.002.
- Machtiger, E.L. *et al.* (2015) 'An Expressive Therapy Group Disclosure Intervention for Women Living With HIV Improves Social Support, Self-efficacy, and the Safety and Quality of Relationships: A Qualitative Analysis', *Journal of the Association of Nurses in AIDS Care*, 26(2), pp. 187–198. doi:10.1016/j.jana.2014.05.001.
- Martiana, I., Waluyo, A. and Yona, S. (2019) 'Assessing the relationship between knowledge of antiretroviral therapy and stigma regarding adherence to ART among men who have sex with men', *Enfermería Clínica*, 29, pp. 321–325. doi:<https://doi.org/10.1016/j.enfcli.2019.06.004>.
- Mathers, C.D. and Loncar, D. (2006) 'Projections of global mortality and burden of disease from 2002 to 2030', *PLoS Medicine*, 3(11), pp. 2011–2030. doi:10.1371/journal.pmed.0030442.
- Mi, T. *et al.* (2020) 'HIV Disclosure to Family Members and Medication Adherence: Role of Social Support and Self-efficacy', *AIDS and Behavior*, 24(1), pp. 45–54. doi:10.1007/s10461-019-02456-1.
- Norman, P. and Conner, M. (2005) 'The theory of planned behavior and exercise: Evidence for the mediating and moderating roles of planning on intention-behavior relationships.', *Journal of Sport & Exercise Psychology*, 27(4), pp. 488–504.
- Nuraidah *et al.* (2022) "'I can live a normal life": Exploring adherence to antiretroviral therapy in Indonesian adolescents living with HIV', *Belitung Nursing Journal*, 8(2), pp. 108–114. doi:10.33546/bnj.2024.
- Nurfalah, F., Yona, S. and Waluyo, A. (2019) 'The relationship between HIV stigma and adherence to antiretroviral (ARV) drug therapy among women with HIV in Lampung, Indonesia', *Enfermería Clínica*, 29, pp. 234–237. doi:<https://doi.org/10.1016/j.enfcli.2019.04.138>.
- Nurihwani (2017) 'Hubungan Dukungan Keluarga Terhadap Kepatuhan Pengobatan Antiretroviral (ARV) Pada Orang Dengan HIV Dan Aids (ODHA) di Puskesmas Jumpandang Baru'.
- Oshotse, C. *et al.* (2018) 'Self-efficacy and adherence behaviors in rheumatoid arthritis patients', *Preventing Chronic Disease*, 15(10), pp. 1–10. doi:10.5888/pcd15.180218.

- Peng, Z. *et al.* (2020) 'Reliability and validity of the lifewindows information-motivation-behavioral skills antiretroviral therapy adherence questionnaire among HIV+ patients in Shanghai', *Patient Preference and Adherence*, 14, pp. 507-515. doi:10.2147/PPA.S234041.
- Purwaningsih, P., Asmoro, C.P. and Prastiwi, Y.A. (2022) 'Self-esteem and motivation with adherence of people living with HIV/AIDS (PLWHA) in Indonesia with antiretroviral therapy: A cross-sectional study', *International Journal of Adolescent Medicine and Health*, 34(1), pp. 1-7. doi:10.1515/ijamh-2019-0051.
- Ramaiya, M.K. *et al.* (2020) 'Patient-Provider Communication and Information, Motivation, and Behavioral Skills in HIV-Positive Adults Initiating Antiretroviral Therapy in Haiti', 19, pp. 1-10. doi:10.1177/2325958220952631.
- Safira, N., Lubis, R. and Fahdhy, M. (2018) 'Factors Affecting Adherence to Antiretroviral Therapy', *KnE Life Sciences*, 4(4), p. 60. doi:10.18502/cls.v4i4.2264.
- Scholz, U. *et al.* (2007) 'Dynamics in self-regulation: Plan execution self-efficacy and mastery of action plans', *Journal of Applied Social Psychology*, 37(11), pp. 2706-2725. doi:10.1111/j.1559-1816.2007.00277.x.
- Scholz, U. *et al.* (2008) 'Beyond behavioural intentions: Planning mediates between intentions and physical activity', *British Journal of Health Psychology*, 13(3), pp. 479-494. doi:10.1348/135910707X216062.
- Schwarzer, R. and Luszczynska, A. (2008) 'How to overcome health-compromising behaviors: The health action process approach', *European Psychologist*, 13(2), pp. 141-151. doi:10.1027/1016-9040.13.2.141.
- Seyoum, D. *et al.* (2017) 'Risk factors for mortality among adult HIV/AIDS patients following antiretroviral therapy in southwestern ethiopia: An assessment through survival models', *International Journal of Environmental Research and Public Health*, 14(3), pp. 1-12. doi:10.3390/ijerph14030296.
- Sniehotta, F.F. *et al.* (2005) 'Action planning and coping planning for long-term lifestyle change, Theory assessment', *European Journal of Social Psychology*, 35(December 2004), pp. 565-576.
- Tharek, Z. *et al.* (2018) 'Relationship between self-efficacy, self-care behaviour and glycaemic control among patients with type 2 diabetes mellitus in the Malaysian primary care setting', *BMC Family Practice*, 19(1), pp. 1-10. doi:10.1186/s12875-018-0725-6.
- Wardhani, S.F. and Yona, S. (2021) 'Spousal intimacy, type of antiretroviral drug and antiretroviral therapy adherence among hiv patients in bandung, indonesia', *Journal of Public Health Research*, 10(s1), pp. 17-21. doi:10.4081/jphr.2021.2336.
- Wee, Z.Q.C. and Dillon, D. (2022) 'Increasing Physical Exercise through Action and Coping Planning', *International Journal of Environmental Research and Public Health*, 19(7). doi:10.3390/ijerph19073883.
- World Health Organization (2020) *The top 10 causes of death*, World Health Organization.
- World Health Organization (2023a) *HIV*, World Health Organization.
- World Health Organization (2023b) *HIV and AIDS*, World Health Organization.
- Zhang, L. *et al.* (2016) 'Side effects, adherence self-efficacy, and adherence to antiretroviral treatment: a mediation analysis in a Chinese sample', *AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV*, 28(7), pp. 919-926. doi:10.1080/09540121.2015.1124984.
- Zhou, G. *et al.* (2018) 'Influence of Side Effects on ART Adherence Among PLWH in China: The Moderator Role of ART-Related Knowledge', *AIDS and Behavior*, 22(3), pp. 961-970. doi:10.1007/s10461-017-1791-9.
- Ziegelmann, J.P., Lippke, S. and Schwarzer, R. (2006) 'Adoption and maintenance of

physical activity: Planning interventions in young, middle-aged, and older adults', *Psychology and Health*, 21(2), pp. 145–163. doi:10.1080/1476832050018891.