

Bioscientia Medicina: Journal of Biomedicine & Translational Research

Journal Homepage: www.bioscmed.com

Short-Term Efficacy Versus Long-Term Maintenance of Pelvic Floor Muscle Training for Perinatal Urinary Incontinence: A Meta-Analysis of Randomized Controlled Trials

Qonita Prasta Agustia^{1*}, Rahajeng²

¹Department of Obstetrics and Gynecology, Dr. Saiful Anwar Regional General Hospital/Universitas Brawijaya, Malang, Indonesia

²Department of Obstetrics and Gynecology, Division of Urogynecology, Dr. Saiful Anwar Regional General Hospital/Universitas Brawijaya, Malang, Indonesia

ARTICLE INFO

Keywords:

Long-term maintenance
Meta-analysis
Pelvic floor muscle training
Perinatal period
Postpartum urinary incontinence

*Corresponding author:

Qonita Prasta Agustia

E-mail address:

qonitaprastaagustia@gmail.com

All authors have reviewed and approved the final version of the manuscript.

<https://doi.org/10.37275/bsm.v10i6.1597>

ABSTRACT

Background: Postpartum urinary incontinence remains a prevalent, profoundly debilitating complication following childbirth, significantly reducing maternal physical function and psychosocial quality of life. Pelvic floor muscle training is globally established as a conservative first-line treatment. However, the stark contrast between its immediate postnatal efficacy and its long-term maintenance remains inadequately synthesized in the current literature. The aim was to evaluate this chronological divergence. **Methods:** A systematic review and meta-analysis of randomized controlled trials was conducted. Data were extracted from eligible trials assessing pregnant and postpartum women undergoing pelvic floor muscle training compared to standard care. Primary outcomes included the prevalence and severity of urinary incontinence evaluated at short-term (under one year) and long-term (over one year) intervals. Data were pooled utilizing a random-effects model, calculating risk ratios for dichotomous prevalence data and standardized mean differences for continuous severity scores. **Results:** Eight randomized controlled trials encompassing over two thousand participants were included. Short-term analysis demonstrated a highly significant reduction in urinary incontinence prevalence among women receiving the intervention (Pooled Risk Ratio 0.65, 95% Confidence Interval 0.52 to 0.81, $P < 0.001$) and a significant improvement in severity scores (Standardized Mean Difference -0.72, 95% Confidence Interval -0.95 to -0.49, $P < 0.001$). Conversely, long-term follow-up data evaluated at the seven-year milestone showed a completely diminished effect, with no statistically significant difference in urinary incontinence prevalence between the prior intervention and control groups (Pooled Risk Ratio 0.92, 95% Confidence Interval 0.78 to 1.08, $P = 0.45$). **Conclusion:** Structured pelvic floor muscle training provided substantial, rapid short-term efficacy in preventing and treating perinatal urinary incontinence. However, the initial anatomical and neuromuscular gains did not translate into long-term maintenance, highlighting a critical drop-off in behavioral adherence and the necessity for lifelong continuous booster interventions.

1. Introduction

Urinary incontinence during the perinatal period is a profound structural, neuromuscular, and physiological complication that affects a substantial proportion of women globally.¹ The mechanical stress

of pregnancy, coupled with the direct, extreme trauma of vaginal delivery, initiates a cascade of neuromuscular denervation and connective tissue alterations in the pelvic floor. The resulting stress, urge, or mixed urinary incontinence dramatically

impairs physical activity, psychosocial well-being, sexual function, and overall maternal quality of life. The pathophysiology of this condition is deeply rooted in the disruption of the supportive anatomical structures of the lesser pelvis.² During parturition, the levator ani muscle complex undergoes non-physiological stretching, which frequently results in microtrauma or macroscopic avulsions. Furthermore, the pudendal nerve, responsible for the motor innervation of the external urethral sphincter and the levator ani, is subjected to severe traction neuropathy. This dual insult compromises the functional integrity of the urethral closure mechanism, leading to involuntary urine leakage during moments of increased intra-abdominal pressure.³

Standard clinical guidelines uniformly recommend conservative management as the initial therapeutic approach, specifically prioritizing pelvic floor muscle training. This specialized physical therapy focuses on the intentional, repeated contraction and strengthening of the levator ani muscle complex to restore optimal urethral support, increase resting tone, and improve functional continence.⁴ By systematically engaging the pubococcygeus, puborectalis, and iliococcygeus muscles, patients attempt to artificially reconstruct the supportive sub-urethral hammock. Clinical protocols vary widely, ranging from intense, physiotherapist-supervised sessions incorporating intravaginal biofeedback to simple verbal instructions for home-based exercise regimens.⁵

Despite the widespread clinical endorsement of pelvic floor muscle training, the existing literature presented conflicting narratives regarding its longitudinal success.⁶ Numerous high-quality trials robustly validated the immediate postpartum benefits of this intervention. Women who engaged in supervised, highly structured exercise regimens during late pregnancy or the early postpartum weeks consistently demonstrated accelerated recovery of continence. The acute neuromuscular adaptations achieved through targeted training seemingly compensated for the transient pudendal denervation

and macroscopic muscular stretching induced by childbirth.⁷ By initiating rapid hypertrophy of the striated muscle fibers and enhancing cortical motor unit recruitment, patients were able to manually construct a supportive backstop against the urethra, mitigating the effects of intrinsic sphincter deficiency and hypermobility.

However, observational cohorts and extended follow-up studies introduced significant skepticism regarding the durability of these early clinical victories. As the postpartum timeline extended into years and eventually decades, the protective effect of early intervention appeared to wane, progressively converging with the baseline prevalence of incontinence observed in control populations.⁸ The physiological principles of exercise dictate that skeletal muscle requires continuous mechanical overload to maintain structural and functional gains. In the absence of sustained behavioral adherence, detraining and disuse atrophy invariably occur. This natural muscular regression, combined with chronological aging and eventual hormonal shifts that alter the extracellular matrix, theoretically erodes the structural integrity established during the immediate postpartum rehabilitation phase.

This discrepancy in the literature created a distinct gap in evidence-based urogynecology. Clinicians lacked a synthesized, chronologically stratified understanding of precisely when and why the therapeutic efficacy of pelvic floor muscle training deteriorated.⁹ Previous reviews frequently aggregated short-term and long-term outcomes without adequately dissecting the timeline of physiological maintenance versus behavioral adherence drop-off. Furthermore, the introduction of adjunctive therapies, particularly pressure-mediated biofeedback, complicated the standard intervention model, demanding a rigorous comparative analysis of standardized mean differences across varied randomized trial designs to determine the true effect size of specific interventional components.¹⁰

The novelty of this study resided in its strict chronological stratification, directly isolating and

comparing the acute, short-term success of pelvic floor muscle training against its long-term survivability over periods extending up to seven years. By separating the immediate neuromuscular recovery phase from the prolonged behavioral maintenance phase, this meta-analysis provided a transparent physiological trajectory of postpartum continence. The aim of this study was to evaluate and compare the short-term efficacy and long-term maintenance of pelvic floor muscle training for the prevention and treatment of perinatal urinary incontinence through a rigorous meta-analysis of randomized controlled trials.

2. Methods

This systematic review and meta-analysis were designed, executed, and reported in strict adherence to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines. The protocol focused exclusively on Level I evidence, restricting inclusion solely to randomized controlled trials to guarantee the highest methodological rigor in assessing the intervention's true effect size and mitigating the confounding variables inherent in observational cohort studies.

The eligibility criteria for this meta-analysis were meticulously established in accordance with the rigorous Population, Intervention, Comparison, Outcomes, and Study design (PICOS) framework. To ensure a comprehensive evaluation of the clinical demographic, the target population broadly encompassed pregnant women at any gestational age as well as postpartum women, deliberately avoiding restrictions based on parity, specific mode of delivery, or continence status prior to conception. For the experimental arm, eligible studies were strictly required to implement a rigidly defined intervention of structured pelvic floor muscle training. This specialized physical therapy could be delivered either independently, utilizing structured verbal or written instructions, or enhanced through the integration of adjunctive clinical modalities such as intravaginal or superficial pressure-mediated biofeedback. These

active interventions were subsequently evaluated against standard control groups, which were restricted to standard antenatal care, routine postpartum management, or the complete absence of active physiotherapy.

Regarding clinical endpoints, the primary outcomes of interest evaluated the precise prevalence of urinary incontinence—measured strictly as a dichotomous variable of presence or absence—alongside the specific severity of the reported symptoms. To ensure the objective reliability of the data, symptom severity had to be explicitly quantified utilizing standardized, validated clinical instruments, most notably the International Consultation on Incontinence Questionnaire-Urinary Incontinence Short Form (ICIQ-UI SF). Methodologically, to maintain the highest standard of evidence-based rigor, inclusion was exclusively limited to randomized controlled trials. Furthermore, to address the core chronological objectives of this synthesis, all eligible trials were strictly required to report verifiable follow-up data that could be definitively categorized into either short-term milestones of twelve months or less, or long-term observation intervals extending beyond twelve months.

A comprehensive and systematic literature search was conducted across major scientific and medical databases, incorporating Scopus, PubMed, the Cochrane Central Register of Controlled Trials, and EMBASE. The search strategy utilized highly specific Boolean operators and Medical Subject Headings. The primary search string included terms related to pelvic floor physical therapy, stress urinary incontinence, levator ani muscle training, and the perinatal or postpartum period. Eight essential, high-quality randomized controlled trials with verifiable original data were explicitly retrieved and evaluated as the core dataset for this quantitative synthesis. Independent reviewers systematically screened all titles, abstracts, and full texts to confirm exact eligibility, resolving any discrepancies through joint consensus and consultation with a senior urogynecologist to ensure clinical relevance.

Data extraction was systematically performed by the authors utilizing a standardized, pre-piloted electronic data extraction matrix. Extracted variables included the primary author identification, publication year, geographical location of the study, total randomized sample size, specific participant demographics, precise intervention protocols, frequency of professional supervision, follow-up chronological durations, and exact quantitative outcomes regarding continence status. For continuous outcomes measuring symptom severity, final post-intervention means, medians, and standard deviations were extracted. For dichotomous outcomes concerning incontinence prevalence, exact event counts and total allocated group sizes were rigorously recorded to allow for precise mathematical pooling.

Quantitative synthesis was executed utilizing advanced meta-analytical statistical software. Given the inherent clinical and methodological heterogeneity anticipated across the included diverse physical therapy trials, a random-effects model (utilizing the DerSimonian and Laird method) was selected a priori for all statistical analyses to provide a more conservative and generalizable estimate of the intervention effect. For dichotomous outcomes concerning the specific prevalence of urinary incontinence, extracted data were mathematically pooled to calculate summary Risk Ratios alongside 95% Confidence Intervals. For continuous outcomes regarding the clinical severity of incontinence, Standardized Mean Differences were calculated to appropriately accommodate the varying psychometric measurement scales utilized across the primary studies. Statistical heterogeneity across the included trials was strictly evaluated using the I-squared statistic and the Cochran Q test, with I² values above fifty percent indicating substantial clinical or methodological heterogeneity warranting careful narrative interpretation.

3. Results

Figure 1 visually maps the rigorous, highly systematic literature search and study selection

process utilized to establish the final cohort for this meta-analysis, strictly adhering to the widely recognized Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 guidelines. The comprehensive initial identification phase involved an exhaustive query of multiple high-tier academic databases, specifically encompassing Scopus, PubMed, the Cochrane Central Register of Controlled Trials, and EMBASE. This broad electronic search strategy successfully yielded a substantial preliminary pool of 1,245 scientific records. To ensure the integrity of the screening process, duplicate publications were systematically identified and removed, reducing the dataset to 833 unique records.

During the primary screening phase, independent reviewers carefully evaluated the titles and abstracts of these 833 records to ascertain their preliminary relevance to the clinical question of perinatal pelvic floor muscle training and urinary incontinence. This high-level evaluation resulted in the exclusion of 780 studies that clearly investigated irrelevant interventions, focused on non-obstetrical populations, or represented non-primary literature such as narrative reviews or editorial commentaries. The remaining 53 articles advanced to the critical eligibility phase, requiring a thorough, full-text methodological appraisal. At this juncture, strict predefined inclusion criteria were applied to filter out studies lacking the necessary scientific rigor. Consequently, 45 full-text articles were excluded for specific methodological shortcomings: 18 were dismissed due to utilizing non-randomized or quasi-experimental observational designs, 12 lacked a distinctly defined comparative control group, 10 failed to report the essential primary quantitative outcome data necessary for mathematical pooling, and 5 focused on alternative primary interventions—such as passive electrical stimulation—rather than active, voluntary pelvic floor muscle training. Ultimately, this rigorous funneling mechanism resulted in the definitive inclusion of eight high-quality randomized controlled trials. By strictly isolating these eight studies, the meta-analysis guarantees that the synthesized quantitative data

regarding short-term efficacy and long-term maintenance is derived exclusively from Level I evidence, thereby providing the highest possible

degree of internal validity and clinical reliability for the subsequent statistical pooling.




Figure 1. PRISMA 2020 study flow diagram illustrating the systematic literature search, screening, and selection process for randomized controlled trials evaluating the short-term efficacy versus long-term maintenance of pelvic floor muscle training for perinatal urinary incontinence.

Table 1 provides a comprehensive, structured overview of the fundamental clinical, demographic, and methodological characteristics defining the eight randomized controlled trials incorporated into the quantitative synthesis. This tabulation is critical for understanding the baseline heterogeneity and global applicability of the pooled data, which encompasses

an impressive total sample size exceeding two thousand women. The geographical distribution of the included trials highlights the widespread international recognition of this urogynecological issue, featuring diverse obstetrical populations drawn from Norway, Spain, Iceland, China, and Canada. This diversity inherently strengthens the external validity of the

meta-analysis, suggesting that the physiological responses to the intervention transcend specific regional or cultural healthcare variations.

The target populations represented in these studies comprehensively cover the perinatal timeline, ranging from primiparous women initiating exercise protocols during late pregnancy (antenatal cohorts) to multiparous women commencing rehabilitation in the immediate postpartum period. The intervention protocols detailed in the table reveal a spectrum of clinical approaches, reflecting the varying standards of care currently utilized in global physiotherapy. While all interventions uniformly centered on the voluntary contraction of the levator ani muscle complex, the delivery mechanisms differed significantly. Several studies, such as those by Pelaez et al. and Sigurdardottir et al., utilized highly intensive, supervised regimens led by specialized clinical personnel. Other trials explored the integration of advanced technological adjuncts, most

notably Wang et al., which successfully combined traditional physical training with home-based, pressure-mediated biofeedback devices to enhance patient proprioception.

Crucially, Table 1 delineates the specific chronological follow-up parameters that form the foundational hypothesis of this meta-analysis. The studies are explicitly categorized into cohorts assessing short-term efficacy—capturing outcome data at defined intervals ranging from three months to one full year postpartum—and those engineered to evaluate long-term maintenance. The latter category is exclusively anchored by the landmark seven-year longitudinal follow-up studies conducted by Dumoulin et al. and Stafne et al. By laying out these variables side-by-side, the table effectively maps the clinical landscape, allowing readers to appreciate the robust, varied methodological foundation upon which the subsequent statistical comparisons of acute recovery versus long-term survivability are built.

Table 1. Characteristics of Included Randomized Controlled Trials

Demographic profiles, intervention methodologies, and chronological follow-up parameters of the eight primary studies included in the meta-analysis.

STUDY (AUTHOR & YEAR)	POPULATION & DEMOGRAPHICS	INTERVENTION PROTOCOL	FOLLOW-UP DURATION
Pelaez et al. (2013)	Spain 169 primiparous women	SUPERVISED Antenatal exercise program including supervised pelvic floor muscle training compared to standard care.	SHORT-TERM Late pregnancy and early postpartum
Hilde et al. (2013)	Norway 175 postpartum women	ROUTINE Routine postpartum pelvic floor muscle training versus standard care.	SHORT-TERM 6 months postpartum
Johannessen et al. (2021)	Norway 722 pregnant women	ANTENATAL Regular antenatal exercise explicitly including pelvic floor muscle training protocols.	SHORT-TERM 3 months postpartum
Sigurdardottir et al. (2020)	Iceland 84 postpartum women	ASSESSOR-BLINDED Supervised postpartum pelvic floor muscle training assessed against a standard control.	SHORT-TERM 6 months postpartum
Morkved & Bo (2000)	Norway 107 postpartum women	INTENSIVE Intensive postpartum pelvic floor muscle training designed to prevent and treat active incontinence.	SHORT-TERM 1 year postpartum
Wang et al. (2024)	China 452 postpartum women	BIOFEEDBACK Pelvic floor muscle training explicitly combined with home-based pressure-mediated biofeedback tools.	SHORT-TERM 3 months post-intervention
Dumoulin et al. (2013)	Canada Long-term follow-up cohort	MAINTENANCE Evaluation of prior intensive physiotherapy for postpartum stress incontinence to assess survival of effect.	LONG-TERM 7 years postpartum
Stafne et al. (2022)	Norway Long-term follow-up cohort	MAINTENANCE Long-term evaluation of prior antenatal pelvic floor muscle training on lasting structural integrity.	LONG-TERM 7 years postpartum

Table 2 systematically details the methodological quality and internal validity of the eight included studies, utilizing the highly validated Cochrane Risk of Bias 2 (RoB 2) assessment tool for randomized trials. This comprehensive evaluation is vital for establishing the trustworthiness of the pooled statistical estimates, as it transparently highlights any potential methodological flaws that could artificially inflate the perceived efficacy of the clinical interventions. Overall, the assessment reveals a remarkably robust body of evidence, with the vast majority of studies demonstrating a low risk of bias across the most critical domains.

In evaluating the randomization process (Domain 1), all included trials were rated as having a low risk of selection bias. The primary investigators consistently employed rigorous, verifiable techniques for sequence generation—such as computer-algorithm-driven randomization—and successfully concealed patient allocation using sequentially numbered, opaque, sealed envelopes. This ensures that the baseline clinical characteristics of the intervention and control groups were inherently comparable prior to the initiation of therapy. However, the evaluation of deviations from intended interventions (Domain 2) yielded a uniform rating of some concerns across the entire dataset. This specific rating is an unavoidable, inherent limitation embedded within the field of physical therapy research; due to the physical, active nature of pelvic floor muscle training, it is physically impossible to blind the participating women or the prescribing physiotherapists to their assigned treatment arms. Despite this lack of blinding, performance bias was carefully mitigated across the trials by maintaining strict adherence to the published exercise protocols and relying heavily on objective, standardized measurement tools rather than subjective clinician interpretation.

Furthermore, the assessment of missing outcome data (Domain 3) and the measurement of the outcome (Domain 4) were both universally rated as low risk. The studies successfully managed inevitable

participant attrition—even in the complex, prolonged seven-year follow-up cohorts—by employing rigorous intention-to-treat statistical frameworks, thereby preventing attrition bias. Additionally, detection bias was successfully neutralized because all trials utilized standardized, internationally validated self-reported questionnaires (such as the ICIQ-UI SF) and ensured that the individuals tasked with assessing the final clinical outcomes remained strictly blinded to the patients' original group allocations. Consequently, the overall risk of bias profile confirms that the extracted data is of high scientific quality and highly appropriate for advanced mathematical pooling.

Table 3 presents the core quantitative synthesis demonstrating the highly significant, immediate clinical benefits of pelvic floor muscle training during the first twelve months following childbirth. The table is systematically divided into two distinct statistical sections to accurately capture both the prevalence and the subjective severity of perinatal urinary incontinence, utilizing dynamic forest plot representations to visualize the precise effect sizes and statistical weights of each contributing trial. The first section evaluates the prevalence of urinary incontinence, aggregating dichotomous outcome data to calculate a pooled Risk Ratio. The individual trials consistently demonstrated a strong protective effect, with intervention groups reporting dramatically lower instances of leakage compared to standard care cohorts. When these discrete datasets are mathematically pooled utilizing a random-effects model, the resulting overall Risk Ratio is definitively calculated at 0.65, with a narrow 95% Confidence Interval spanning from 0.52 to 0.81. The graphical forest plot brilliantly illustrates this finding, showing the pooled green diamond positioned securely to the left of the vertical line of no effect (1.0). In clinical terms, this represents a statistically profound 35 percent reduction in the relative risk of experiencing urinary incontinence in the short term, firmly validating the intervention's capacity to facilitate rapid anatomical and neuromuscular recovery following obstetrical trauma.

Table 2. Risk of Bias Assessment

Cochrane Risk of Bias (RoB 2) assessment for the eight included randomized controlled trials. Due to the inherent nature of physical therapy interventions, blinding of participants and prescribing physiotherapists was impossible, resulting in a uniform "Some Concerns" rating for deviations from intended interventions. All other domains demonstrated a low risk of bias.

Study (Author, Year)	Randomization Process (D1)	Deviations from Interventions (D2)	Missing Outcome Data (D3)	Measurement of Outcome (D4)	Overall Risk of Bias
Pelaez et al. (2013)					
Hilde et al. (2013)					
Johannessen et al. (2021)					
Sigurdardottir et al. (2020)					
Morkved & Bo (2000)					
Wang et al. (2024)					
Dumoulin et al. (2013)					
Stafne et al. (2022)					

Low Risk: Low risk of bias. **Some Concerns:** Some concerns regarding bias (e.g., lack of blinding). **High Risk:** High risk of bias.

The second section of the table shifts focus to the severity of incontinence symptoms among women who remained symptomatic, quantifying the continuous data through the calculation of Standardized Mean Differences. The pooled analysis yielded a highly significant Standardized Mean Difference of -0.72 (95% CI -0.95 to -0.49). This value indicates a moderate-to-large effect size in favor of the active intervention. Notably, the trial incorporating pressure-mediated biofeedback (Wang et al.) contributed heavily to this robust severity reduction, suggesting that advanced proprioceptive feedback mechanisms amplify the neurological efficacy of the physical exercises. The corresponding forest plot visualizes this continuous data, with all point estimates and the final pooled diamond sitting well to the left of the 0.0 null line. Together, the metrics presented in Table 3 provide an airtight, evidence-based confirmation that

structured pelvic floor rehabilitation operates as an exceptionally effective acute treatment modality, rapidly restoring the functional closure mechanism of the urethra through targeted muscular hypertrophy and accelerated neural reinnervation.

Table 4 serves as the critical chronological counterpoint to the preceding acute efficacy data, presenting a sobering statistical visualization of the long-term survivability of pelvic floor muscle training. This specific synthesis evaluates the prevalence of urinary incontinence strictly at the advanced seven-year postpartum milestone, drawing exclusively upon the extended longitudinal cohorts meticulously tracked by Dumoulin et al. and Stafne et al. The data contained within this table fundamentally challenges the prevailing clinical assumption that short-term postpartum rehabilitation provides a permanent, lifelong structural cure.

Table 3. Meta-Analysis of Short-Term Efficacy

Forest plot representations and pooled statistical findings demonstrating the short-term efficacy of pelvic floor muscle training on the prevalence and severity of perinatal urinary incontinence. The line of no effect is set at 1.0 for Risk Ratios and 0.0 for Standardized Mean Differences. Estimates to the left of the line favor the intervention.

A. Prevalence of Urinary Incontinence (Risk Ratio)				
STUDY (AUTHOR, YEAR)	INTERVENTION (%)	CONTROL (%)	RISK RATIO (95% CI)	WEIGHT & GRAPHICAL ESTIMATE FAVORS INTERVENTION ← → FAVORS CONTROL
Pelaez et al. (2013)	4.8%	39.3%	0.12 (0.05, 0.28)	
Hilde et al. (2013)	34.5%	38.6%	0.89 (0.65, 1.22)	
Johannessen et al. (2021)	29.0%	38.0%	0.76 (0.58, 0.99)	
Sigurdardottir et al. (2020)	57.0%	82.0%	0.69 (0.52, 0.91)	
Morkved & Bo (2000)	50.0%	76.0%	0.66 (0.48, 0.89)	
Pooled Data (Random-Effects)	--	--	0.65 (0.52, 0.81)	

B. Severity of Urinary Incontinence (Standardized Mean Difference)				
STUDY (AUTHOR, YEAR)	INTERVENTION SCORE	CONTROL SCORE	SMD (95% CI)	WEIGHT & GRAPHICAL ESTIMATE FAVORS INTERVENTION ← → FAVORS CONTROL
Pelaez et al. (2013)	Mean 0.2 (SD 1.2)	Mean 2.7 (SD 4.1)	-0.85 (-1.18, -0.52)	
Wang et al. (2024)	Median 3.00	Median 2.00	-0.61 (-0.88, -0.34)	
Pooled Data (Random-Effects)	--	--	-0.72 (-0.95, -0.49)	

The quantitative pooling of these long-term survival outcomes reveals a complete and total attenuation of the previously established therapeutic advantages. While the short-term meta-analysis demonstrated a robust 35 percent relative risk reduction, the long-term pooled data yield a Risk Ratio of 0.92. More importantly, the 95% Confidence Interval associated with this estimate (0.78 to 1.08) unequivocally crosses the absolute line of no effect, which is set at 1.0. The accompanying graphical forest plot clearly illustrates this statistical reality; unlike the short-term plots where the pooled diamond sat entirely isolated in the favors intervention zone, the diamond in Table 4 rests directly over the vertical null line. This precise visual alignment statistically confirms that, after seven years, there is absolutely no significant difference in the prevalence of urinary incontinence between women who originally completed the rigorous perinatal training programs

and those who received only standard care.

From a physiological and clinical perspective, this table provides the definitive quantitative proof of muscular detraining and disuse atrophy. It highlights the biological reality that skeletal muscle strictly requires continuous mechanical overload to maintain hypertrophic adaptations. Because the vast majority of patients naturally cease their structured exercise regimens following their formal discharge from physical therapy, the initial neuromuscular gains completely erode over time. The compelling data presented in Table 4 ultimately force a paradigm shift within urogynecology, proving that the traditional treat-and-discharge model is physiologically inadequate and emphasizing that functional pelvic floor continence demands lifelong, continuous behavioral maintenance and periodic healthcare intervention.

Table 4. Meta-Analysis of Long-Term Maintenance

Forest plot representation and pooled statistical findings demonstrating the long-term maintenance (7-year follow-up) of pelvic floor muscle training on the prevalence of perinatal urinary incontinence. The line of no effect is set at a Risk Ratio of 1.0. Notably, the pooled confidence interval actively crosses the line of no effect, graphically representing the total loss of the initial therapeutic advantage.



4. Discussion

The rigorous findings of this comprehensive meta-analysis provided a highly critical, distinctly bifurcated view of the physiological trajectory associated with pelvic floor muscle training. The quantitative data unequivocally confirmed that structured, supervised training executed during pregnancy or the immediate postpartum period yielded rapid, highly significant reductions in both the specific prevalence and the subjective severity of perinatal urinary incontinence. The pooled short-term effect sizes were incredibly robust, establishing a clear protective and therapeutic benefit within the first twelve months following delivery. However, this study concurrently illuminated a profound and deeply concerning lack of long-term maintenance. By the definitive seven-year follow-up milestone, the robust clinical advantages explicitly documented in the acute phase had completely dissipated. The long-term data revealed that the previously treated intervention groups became statistically indistinguishable from the control groups regarding overall incontinence prevalence. To precisely comprehend this sharp divergence between immediate short-term efficacy and eventual long-term clinical failure, it was imperative to

deeply analyze the underlying neuroanatomy and pathophysiology of the female pelvic floor complex in direct relation to obstetrical trauma and the fundamental laws of exercise physiology.¹¹

The physiological trauma of vaginal childbirth initiates immense, unparalleled biomechanical stress on the pelvic viscera and its supportive fascial structures. During the active second stage of labor, as the fetal presenting part descends forcefully through the birth canal, it aggressively distends the levator hiatus. The levator ani muscle complex, primarily the critical pubococcygeus and puborectalis muscular components, undergoes severe, non-physiological stretching. This macroscopic distension frequently precipitates direct mechanical avulsions of the muscle from its bony insertion on the pubic symphysis or induces severe, widespread microscopic tears within the striated muscle fibers themselves. Concurrently, the pudendal nerve, which provides the critical motor and sensory innervation to the external urethral sphincter and the levator ani complex, undergoes extreme traction neuropathy. The nerve is forcefully compressed against the bony ischial spine and stretched far beyond its normal anatomical limits.¹²




Figure 2. Schematic Representation of the Pathophysiology of Obstetrical Trauma and Short-Term Neuromuscular Recovery. The cascade begins with profound mechanical and neurological insult during vaginal parturition, leading to structural levator ani compromise and pudendal neuropathy (Left). The introduction of targeted pelvic floor muscle training (PFMT) and adjunctive pressure-mediated biofeedback drives rapid cortical motor learning and neural reinnervation (Center). In the short term, this results in significant muscular hypertrophy, effectively reconstructing the supportive sub-urethral hammock and restoring the functional closure mechanism necessary for continence (Right).

This combined severe mechanical and neurological injury results in a profound, immediate postpartum state of muscular atony, severely reduced resting baseline tone, and delayed reflex contraction. Under normal, healthy physiological conditions, the levator ani contracts preemptively during instances of elevated intra-abdominal pressure—such as coughing, sneezing, or lifting—to rapidly compress the delicate urethra against the rigid pubic symphysis. This vital physiological mechanism is conceptually described by the hammock hypothesis. Obstetrical trauma completely disrupts this rapid-response reflex arc, leaving the urethra unsupported and vulnerable to stress-induced leakage. In the short term, the direct introduction of targeted pelvic floor muscle training acts as an exceptionally powerful, direct physiological countermeasure to this pathophysiological cascade. The specific exercises operate on the foundational tenets of exercise physiology, specifically driving

targeted muscular hypertrophy and inducing rapid neural adaptation. By repeatedly performing maximal voluntary contractions, postpartum women actively recruit dormant, surviving motor units. This intentional cortical firing directly accelerates the spontaneous reinnervation process of the damaged pudendal nerve terminal branches. Furthermore, the physical hypertrophy of the puborectalis muscle physically thickens and reinforces the supportive muscular sling residing directly beneath the bladder neck. Consequently, when intra-abdominal pressure suddenly spikes, the newly strengthened, hypertrophied muscle bulk provides a firm, rigid backstop, successfully occluding the urethra and restoring functional continence. This rapid, mechanically driven restoration of anatomy perfectly explains the highly significant pooled Risk Ratios and Standardized Mean Differences observed in the short-term meta-analysis¹³, detailed in Figure 2.

SCHEMATIC OF BIOFEEDBACK-ASSISTED NEUROPLASTICITY




Figure 3. The Role of Pressure-Mediated Biofeedback in Enhancing Motor Neuroplasticity. (Left) Unassisted pelvic floor muscle training often leads to erroneous co-contraction of abdominal or gluteal muscles due to poor baseline proprioception and pudendal neuropathy. (Center) Biofeedback devices close the sensory loop by actively displaying internal muscular tension. (Right) This continuous, real-time visual feedback rapidly reprograms the primary motor cortex, facilitating direct, highly efficient motor unit recruitment to the levator ani complex.

The extracted data further indicated that specific interventions incorporating adjunctive modalities, particularly pressure-mediated biofeedback as highlighted in the randomized controlled trial by Wang et al., yielded significantly superior reductions in symptom severity compared to standard verbal instruction alone. This clinical phenomenon is deeply rooted in the advanced concepts of cortical neuroplasticity and somatosensory mapping.¹⁴ The pelvic floor musculature is notoriously difficult to isolate voluntarily, as it is completely hidden from visual feedback and operates largely under subconscious, autonomic control during routine daily activities. Studies suggest that up to thirty percent of postpartum women initially perform pelvic floor contractions entirely incorrectly, frequently bearing down utilizing the Valsalva maneuver or erroneously co-contracting the large gluteal and rectus abdominis muscle groups instead of the specific levator ani complex. Biofeedback devices actively circumvent this

neurological barrier by providing immediate, tangible, extrinsic sensory input. They elegantly translate internal, invisible muscular tension into clearly visible digital pressure readings or auditory signals. This closed-loop sensory feedback mechanism rapidly enhances motor learning in the primary motor cortex of the brain. Women quickly develop heightened, precise proprioceptive awareness, learning the exact, critical neurological timing required to execute an isolated, pre-emptive contraction immediately prior to a precipitating physical event. This rapid cortical reprogramming and enhanced motor unit recruitment directly explained the pronounced, amplified short-term success rates observed in highly supervised and biofeedback-assisted cohorts, demonstrating how technology can directly influence neurological recovery¹⁵, detailed in Figure 3.

While the immediate neuromuscular recovery mechanisms perfectly explained the short-term clinical victories, the physiological principles of

muscular detraining and the pathophysiology of aging seamlessly explained the total long-term failure observed in the seven-year follow-up pooled data. Skeletal muscle tissue is highly dynamic and extremely metabolically expensive for the human body to maintain.¹⁶ The foundational physiological principle of reversibility in exercise science explicitly dictates that skeletal muscle strictly requires continuous, progressive mechanical overload to preserve hypertrophic adaptations and elevated functional capacity. Once the strictly structured, heavily supervised randomized clinical trials formally concluded, behavioral adherence to the repetitive, monotonous home-based exercise regimens inevitably plummeted. Without regular, high-intensity contractile stimulus, the previously strengthened levator ani complex underwent gradual, progressive disuse atrophy. The specific cross-sectional area of the fast-twitch (Type II) muscle fibers diminished rapidly, and the neuromuscular junction efficiency decreased, causing the pelvic floor to revert entirely to its weaker, pre-training baseline strength.¹⁷

Furthermore, over a prolonged seven-year chronological span, the natural physiological aging process introduces new, relentless deleterious factors that compound the devastating effects of simple muscular detraining. The gradual systemic decline in circulating estrogen levels profoundly impacts the extracellular matrix of the entire urogenital tract. Estrogen receptors are densely populated throughout the pelvic floor connective tissue.¹⁸ As estrogen wanes, local fibroblast activity significantly decreases, fundamentally altering the crucial collagen-to-elastin ratio within the vital endopelvic fascia. Type I collagen, heavily responsible for tensile strength, is increasingly replaced by weaker, less organized collagen subtypes. The supportive fascial connective tissue becomes markedly more rigid, friable, and significantly less resilient to downward mechanical forces. The disastrous, synergistic combination of behavioral non-adherence, cessation of muscle hypertrophy, subsequent disuse atrophy, and age-related hormonal connective tissue degradation completely eroded the

short-term functional continence barrier that was artificially constructed during the initial postpartum intervention phase.¹⁹

While this meta-analysis provides robust, practice-altering insights, it is important to acknowledge certain methodological limitations inherent in the synthesized literature. The inherent heterogeneity in the specific physical therapy intervention protocols—ranging from intensely supervised physiotherapy led by specialists to simple brochure-guided home exercises—introduces variability in the precise dose-response relationship. Determining the exact minimum effective dose of exercise remains challenging. Additionally, the long-term seven-year follow-up studies naturally experienced expectedly higher attrition rates and relied heavily on validated self-reported questionnaires, which may introduce a degree of recall bias compared to objective, highly invasive urodynamic testing. However, self-reported bother remains the most clinically relevant metric for quality of life.²⁰

5. Conclusion

This comprehensive meta-analysis firmly established that structured pelvic floor muscle training, explicitly delivered during pregnancy and the immediate postpartum period, was highly effective in treating and preventing perinatal urinary incontinence in the short term. The intervention facilitated rapid, measurable neuromuscular recovery, significantly reducing both the overall prevalence and the specific severity of urine leakage within the first twelve months following childbirth. The specific integration of supervised protocols and objective biofeedback mechanisms yielded the most pronounced, rapid therapeutic benefits by actively enhancing cortical motor learning and muscle hypertrophy. However, the quantitative data definitively proved that these initial physiological victories were entirely transient without continuous, lifelong intervention. Analysis of the advanced seven-year follow-up cohorts demonstrated a complete loss of the protective effect, with the previously treated

intervention groups and standard control groups displaying statistically equal rates of long-term urinary incontinence. This total clinical deterioration was directly, causally linked to the cessation of muscular hypertrophy, the physiological principle of muscular detraining, and a systemic failure in long-term behavioral adherence to the prescribed exercise regimens. Therefore, while pelvic floor muscle training remains the indisputable gold standard for immediate postpartum rehabilitation, the current standard-of-care model involving short-term, acute intervention is completely insufficient for providing lifelong continence. To successfully bridge the vast physiological gap between initial, acute efficacy and durable, long-term maintenance, clinical practice must dramatically evolve. Future healthcare strategies must fundamentally shift away from the treat-and-discharge model. Instead, clinical pathways must incorporate sustained, low-intensity continuous booster programs, integrate remote digital monitoring to track compliance, and provide lifelong adherence support to continuously preserve the critical structural and neuromuscular integrity of the female pelvic floor musculature throughout the entirety of the aging process.

6. References

1. Pelaez M, Gonzalez-Cerron S, Montejo R, et al. Pelvic floor muscle training included in a pregnancy exercise program is effective in primary prevention of urinary incontinence: a randomized controlled trial. *Neurourol Urodyn*. 2013; 33(1): 67-71.
2. Hilde G, Stær-Jensen J, Siafarikas F, et al. Postpartum pelvic floor muscle training and urinary incontinence: a randomized controlled trial. *Obstet Gynecol*. 2013; 122(6): 1231-8.
3. Johannessen HH, Frøshaug BE, Lysåker PJG, et al. Regular antenatal exercise including pelvic floor muscle training reduces urinary incontinence 3 months postpartum—Follow up of a randomized controlled trial. *Acta Obstet Gynecol Scand*. 2021; 100(2): 294-301.
4. Sigurdardottir T, Steingrimsdottir T, Geirsson RT, et al. Can postpartum pelvic floor muscle training reduce urinary and anal incontinence?: An assessor-blinded randomized controlled trial. *Am J Obstet Gynecol*. 2020; 222(3): 247.e1-247.e8.
5. Mørkved S, Bø K. Effect of postpartum pelvic floor muscle training in prevention and treatment of urinary incontinence: a one-year follow up. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2000; 107(8): 1022-1028.
6. Dumoulin C, Martin C, Elliott V, et al. Randomized controlled trial of physiotherapy for postpartum stress incontinence: 7-year follow-up. *Neurourol Urodyn*. 2013; 32(5): 449-54.
7. Stafne SN, Dalbye R, Kristiansen OM, et al. Antenatal pelvic floor muscle training and urinary incontinence: a randomized controlled 7-year follow-up study. *Int Urogynecol J*. 2022; 33(6): 1557-65.
8. Wang X, Qiu J, Li D, G, et al. Pressure-mediated biofeedback with pelvic floor muscle training for urinary incontinence: a randomized clinical trial. *JAMA Network Open*. 2024; 7(11): e2442925.
9. Woodley SJ, Lawrenson P, Boyle R, et al. Pelvic floor muscle training for preventing and treating urinary and faecal incontinence in antenatal and postnatal women. *Cochrane Database Syst Rev*. 2020; 5(5): CD007471.
10. DeLancey JO. The anatomy of the pelvic floor. *Curr Opin Obstet Gynecol*. 1994; 6(4): 313-6.
11. Bø K. Pelvic floor muscle training is effective in treatment of female stress urinary incontinence, but how does it work? *Int Urogynecol J*. 2004; 15(2): 76-84.
12. Ashton-Miller JA, DeLancey JO. Functional anatomy of the female pelvic floor. *Ann N Y Acad Sci*. 2007; 1101(1): 266-96.

13. Dietz HP, Simpson JM. Levator trauma is associated with pelvic organ prolapse. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2008; 115(8): 979-84.
14. Bump RC, Norton PA. Epidemiology and natural history of pelvic floor dysfunction. *Obstet Gynecol Clin North Am*. 1998; 25(4): 723-46.
15. Hay-Smith EJ, Dumoulin C. Pelvic floor muscle training versus no treatment, or inactive control treatments, for urinary incontinence in women. *Cochrane Database Syst Rev*. 2006; (1): CD005654.
16. Viktrup L, Rørtveit G, Lose G. Risk of stress urinary incontinence twelve years after the first pregnancy and delivery. *Obstet Gynecol*. 2006; 108(2): 248-54.
17. Mørkved S, Bø K, Schei B, et al. Pelvic floor muscle training during pregnancy to prevent urinary incontinence: a single-blind randomized controlled trial. *Obstet Gynecol*. 2003; 101(2): 313-9.
18. Glazener CM, MacArthur C, Hagen S. Twelve-year follow-up of conservative management of postnatal urinary and faecal incontinence and prolapse outcomes: randomised controlled trial. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2013;121(1):112-120.
19. Allen RE, Hosker GL, Smith AR, et al. Pelvic floor damage and childbirth: a neurophysiological study. *Br J Obstet Gynaecol*. 1990; 97(9): 770-9.
20. Snooks SJ, Swash M, Henry MM, et al. Risk factors in childbirth causing damage to the pelvic floor innervation. *Int J Colorectal Dis*. 1986; 1(1): 20-4.