

Prospect Study of Anti-inflammatory Activity by Identification of *Muntingia calabura* Leaf Infusion

Shenny S.S Permana,¹ Muchamad R.A Yusuf,¹ Aufa A.D Putri,¹ Jasmine Y. Sukmayani,¹
Ratu Z.A.P Sudrajat,¹ Miski A. Khairinisa,² Indah S. Wahyuni,³ Enny Rohmawaty,⁴
Muhammad H. Bashari,⁴ Ronny Lesmana,⁴ Aziiz Mardanarian Rosdianto^{1,4}

¹Veterinary Study Program, Faculty of Medicine, Universitas Padjadjaran, Jatinangor, Indonesia.

²Department of Pharmacology and Clinical Pharmacy, Faculty of Pharmacy,
Universitas Padjadjaran, Jatinangor, Indonesia.

³Department of Oral Medicine, Faculty of Dentistry, Universitas Padjadjaran, Jatinangor, Indonesia.

⁴Department of Biomedical Sciences, Faculty of Medicine, Universitas Padjadjaran,
Jatinangor, Indonesia.

Abstract

Many diseases occur due to inflammation that is not handled properly. One of the feature of inflammation is swelling or edema. Inflammation can be handled with traditional medicine, such as *Muntingia calabura* L. (*M. calabura* L.) Pharmacologically, this plant extract is reported to have antipyretic, analgesic, anti-inflammatory, antioxidant, and antibacterial. Metabolite compound contained in *M. calabura* L. has the potential as an anti-inflammatory agent. The objective of this study is to ascertain the secondary metabolites contained in *M. calabura* L. leaves infusion. This experimental research method includes the simplicia making and infusion of *M. calabura* L. leaves by cold infusion and phytochemical screening. This study also showed that the infusion of *M. calabura* L. leaves contained flavonoids, alkaloids, and triterpenoids. Overall, the findings indicate potential as an anti-inflammatory agent that requires further investigation, specifically in preclinical testing.

Keywords: Anti-inflammatory, infuse, *M. calabura* L, phytochemical screening

Introduction

Several diseases arise when inflammation symptoms are not promptly and accurately addressed. Osteoarthritis becomes a problem that arises due to inadequately managed inflammation. This condition is prevalent among racehorses and other equine athletes.

It occurs due to repetitive joint trauma, leading to cartilage damage and erosion of bone¹. A study found that 33% of thoroughbred racehorses aged 2-3 years suffered from

osteoarthritis and lesions in their articular cartilage.² Other diseases that can emerge from inflammation in horses is muscle and orthopedic pain, perioperative pain, corneal ulcers, uveitis, laminitis, and gastrointestinal pain (colic).³ Efforts to prevent the worsening of trauma can be reduced using anti-inflammatory drugs. Indonesia has a wealth of natural medicines and traditional remedies that many people have been using for generations.⁴

Corresponding author: Aziiz M. Rosdianto. Department of Biomedical Sciences, Faculty of Medicine, Universitas Padjadjaran, Jatinangor, Indonesia. Email: a.m.rosdianto@unpad.ac.id

Received: 22 Mei 2024. Revised: 3 Juli 2024. Published: 5 August 2024

M. calabura L., also known as kersen, belongs to the family Muntingiaceae and is widely found along the way. This plant typically grows in the wild. Nevertheless, due to its dense foliage, it is deliberately cultivated as a roadside shade tree and an air pollution absorber.⁵ This plant is one of the potential candidates for further development as a traditional medicine for anti-inflammatory purposes. Some research journals state that flavonoids are the main constituents in kersen leaves. Flavonoids themselves are polyphenolic compounds with various activities, one of which is anti-inflammatory.⁵

One direct benefit of traditional medicine formulations for the community is the easy availability of these remedies. The demand for plant-based ingredients in traditional medicine is increasing because they are proven healthier and have fewer side effects than chemical-based alternatives. However, a challenge with traditional medicine is the limited knowledge and information about the different types of plants used as ingredients and how to use them effectively.⁶

Based on this information, the primary goal of this study is to identify the secondary metabolites present in the infusion of *M. calabura L.* leaf so that further tests can be carried out in vivo and in vitro as anti-alternative inflammatory drugs.

Methods

This study is a laboratory experiment to discover the secondary metabolites found in *M. calabura L.* leaf infusion (MCLI). This study was conducted at the Chemical Application Laboratory and Services, Universitas Padjadjaran. This research has received approval and authorization from the Research Ethics Committee of Universitas Padjadjaran (Document No. 1066/UN6.KEP/EC/2023).

Research Procedures

Collection and preparation of sample

Samples of *M. calabura L.* were collected from the Sukapura Village in Kiaracandong District, Bandung City, West Java Province. Before further investigation, the sample was recognize at the Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Padjadjaran (Document No. No.212/LBM/IT/VIII/2023). The sample was cleaned of attached dirt, cut into small pieces, and dried in the dryer.

Preparation of M. calabura L. Infusion

The simplicia of *M. calabura L.* leaves (50 g) was extracted with 200mL distilled water (1:4). At room temperature (25-30°C), let it stays for 1 hour before being placed in the refrigerator for 19 hours. Next, the infusion substance was filtered and put into a falcon tube. Adaptive method for cold infusion was followed Farmakope Herbal Indonesia year 2017.

Phytochemical Screening

Phenolic Test

Two grams of sample was mixed with 5 mL methanol. The solution was refined then 5% ferric chloride solution was added. When phenolic compounds are present, a positive outcome is indicated by the formation of a bluish-black color.⁷

Tannins Test

Roughly 2 grams of the plant extract were boiling with 5 mL of distilled water. After that, added 0.1% Ferric chloride to the mixture and watched for the appearance of a greenish-black color, which would suggest that tannins were present.⁷

Alkaloid Test (Wagner's Test)

Two grams of sample was mixed with 5 mL chloroform, then add 5 mL sulphuric acid. Next, put a few drops of Wagner's reagent

into the solution. If a reddish-brown or brown clumpy substance forms, it means the test is positive.⁷

Flavonoid Test

1. Shinoda test: combined 2 grams of the sample with 5 mL of methanol solution. Next, put in a tiny piece of magnesium chunk. Then, added 2 drops of concentrated HCl. When the color turned orange, it meant that there were flavonoids present.
2. Two grams of sample was mixed with 5 mL methanol solution; then plant solution was treated with 2-3 drops of 10% sodium hydroxide solution. The development of a strong yellow color suggests that flavonoids are present.
3. Mixed two grams of the sample with a 5 mL methanol solution and then reacted it with two drops of 2N hydrochloric acid. The sudden appearance of a acute yellow color shows that there are flavonoids present.⁷

Saponins (Foam Test)

Two grams of sample was mixed with 5 mL aquadest, filtered the solution, shaken vigorously for 2 minutes, then reacted with two drops hydrochloride acid 2N. The formation of a persistent froth for 10 minutes indicated the presence of saponins.⁷

Triterpenoids and Steroids Tests

Mixed two grams of each sample with ethanol. 2 mL chloroform was extracted with 1 mL chloroform and water (1:1). Then added with 1 mL concentrated sulphuric acid. Reddish-brown coloration indicated the presence of triterpenoids. Green, blue, and violet pigment indicated the presence of steroids.⁷

Results and Discussion

Extraction Yield

The cold infusion technique was employed in this research to extract compounds from the leaves of *M. calabura L.*, resulting in an extract yield of 0.1%. The cold infusion technique differs from the hot infusion technique. This method involves using refrigerator temperatures where the sample is left to steep for a minimum of 12-24 hours.⁸ A non-thermal extraction method was selected to prevent any potential damage to heat-resistant active metabolites that may be present in the sample. Biological factors (plant part, plant species, location of growth, and harvesting time) and chemical factors (size, hardness, dryness of the material, levels and types of active compounds contained in plant material, type of solvent use, extraction methods) Can affect the quality of the extract.⁹

Table 1. Phytochemical Screening Results

No	Test	Testing Method	Result
1.	Phenolic	FeCl ₃ reagent 5%	-
2.	Tannins	FeCl ₃ reagent 1%	-
3.	Flavonoids	Concentrated HCL reagent + Mg	-
		Reagent H ₂ SO ₄ 2N	-
		NaOH reagent 10%	+
4.	Saponins	Reagent HCL 2N	-
5.	Triterpenoids		+
	Steroids	Concentrated H ₂ SO ₄ reagent	-
6.	Alkaloids	Wagner's reagent	+

Note: (-) : Not detected | (+) : Detected




Figure 1. Phytochemical screening results of phenolics compounds with 5% ferric chloride solution: (a) before and (b) after adding the reagent.




Figure 2. Phytochemical screening results of tannin compounds with 1% ferric chloride solution: (a) before and (b) after adding the reagent.




Figure 3. Basic Flavonoids Structure¹⁴




Figure 4. Pinocembrin Structure²¹




Figure 5. Phytochemical screening results of flavonoids compounds found in 10% sodium hydroxide solution reagent: (a) before and (b) after adding the reagent.




Figure 6. Phytochemical screening results of saponins compounds with hydrochloric acid 2N: (a) and (b) after adding the reagent.




Figure 7. Phytochemical screening results of triterpenoids and steroids compounds with concentrated sulphuric reagent: (a) before and (b) after adding the reagent.




Figure 8. Phytochemical screening results of alkaloids compounds using Wagner's reagent: (a) before and (b) after adding the reagent

Meanwhile, the extraction process employs the infusion method due to its relatively shorter and faster preparation time and the easy availability of materials and tools.¹⁰ Water is a commonly used solvent for extracting active constituents soluble in water from plant materials, despite the drawback of the infusion being susceptible to mold growth.

Phytochemical Screening

The content of secondary metabolite compounds in the leaves through phytochemical screening for the test was phenolics, tannins, flavonoids, saponin, triterpenoids, steroids, and alkaloids showed the following results as in Table 1.

Phenolics

Test for phenolic compounds obtained negative results (Fig. 1). The level of phenolics in plant sources rests on factors such as maturation processes, cultivation techniques, cultivars, growing conditions, and processing and storage conditions, among others.¹¹

Phenolic compounds of MCLI that were not detected may have been influenced by extraction method factors. According to a study, Many traditional methods used to extract phenolic compounds from plants, like soxhlet extraction, percolation, and maceration, have their own problems. These include getting only small amounts of the compounds, using a lot of extraction solvents, taking a long time, and creating a lot of waste.

Because of these issues, new methods like Microwave Assisted Extraction. (MAE), (Ultrasonic Assisted Extracction) Ultrasonic Assisted Extracction (UAE) , Supercritical Carbon Dioxide (SC-CO₂), Enzyme Assisted Extraction (EAE), and Pressure Liquid Extraction (PLE) have come up. These techniques are unconventional but help fix the

problems of the methods before¹². Therefore, further testing is necessary to identify the phenolic compounds in *M. calabura L.* leaf.

Tannins

Based on the results of a phytochemical test of tannin compounds, MCLI showed negative results. The results showed an acute yellow solution was formed after adding 1% ferric chloride solution, while positive results were obtained if a greenish-black coloration solution was formed. The results of the study can be viewed in Figure 2.

The tannin content in *M. calabura L.* leaves might be low in amounts detectable through cold infusion methods. The analytical methods employed should be highly sensitive to detect tannins at low concentrations. Several advanced technologies, such as microwaves and ultrasonication, have demonstrated the potential to extract tannins efficiently. Additionally, controlling factors like temperature, solid-to-solvent ratio, particle size, source of the material, and extraction time contribute to obtaining higher-quality tannins.¹³

Flavonoid

Previous study reported that *M. calabura L.* leaves extract rich in flavonoid active metabolites, which show a potential anti-inflammatory activity. Flavonoids direct to inhibiting the biosynthesis of prostaglandins (PGE) and lipooxygenase (LOX), which are enzymes involved in the inflammatory process.

Flavonoid compounds are categorized into various subgroups, such as flavones, flavanones, flavans, and biflavans.¹⁵ One of the flavanone compounds found in kersen leaves is pinocembrin.¹⁶ The study states that pinocembrin possesses anti-inflammatory activity and has been demonstrated in various

disease models.¹⁷ Pinocembrin works by decrease of c-Jun N-terminal Kinase levels, p38/MAPK, and NF- κ B, thereby inhibiting cytokines products.

In a rheumatoid arthritis rat model, pinocembrin decrease joint erosion and the percentage of inflammatory cells.¹⁸ In the ulcerative colitis animal study, administering high doses of pinocembrin demonstrated therapeutic and anti-inflammatory effects, likely due to inhibiting the NF- κ B pathway.¹⁹ Additionally, pinocembrin has shown promising results in respiratory allergic inflammation by reducing Th2-type cytokines (IL-4, IL-5, and IL-13) in bronchoalveolar lavage fluid from sensitized rats through the inhibition of NF- κ B activation blocking).²⁰

Based on the phytochemical screening test results, positive results were found in a 10% sodium hydroxide solution reagent characterized by acute yellow formation. In the other reagent (Table 1.), negative results were obtained because there was no formation of acute yellow color in the sample. The results of the flavanoid compound test using a 10% sodium hydroxide reagent be seen in Figure 5.

Saponin

Saponins are potent surface-active substances that lead to the formation of foam upon being heated in water. The saponins compounds were not detected in MCLI. The saponins test results can be seen in Figure 6.

These results might occur due to insufficient extraction time and temperature. According to a study, higher temperatures and longer extraction durations increase saponin levels.²² Additionally, active saponin compounds are more effectively generated when extracted using methanol as a solvent. This results because methanol's universal nature allows it to capture saponins with polar and non-polar

groups.²³

Triterpenoids and steroids

The test results of triterpenoid compounds in *M. calabura L.* leaves using the 1 mL concentrated sulphuric reagent showed that the leaves contained triterpenoids but not steroid compounds in MCLI.

In a study, it is mentioned that the compound group belongs to the triterpenoid class treatment effects on inflammation complex range, which persists for an onset and duration treatment to treated of chronic diseases, including periodontitis, cerebral edema, sepsis, liver injury, gastric ulcer, acute lung injury, allergic reaction.²⁴ Another group (Escin) showed a significant reduction in inflammation induced in mice in vivo. Moreover, its effects better compared to the positive treatment by NSAID. The Escin mechanism is linked to the glucocorticoid receptor (GR). Through GR activation, escin inhibits the activation of NF- κ B, subsequently reducing the release of inflammatory cytokines (TNF- α and IL-1 β) along with nitric oxide (NO).

Alkaloids

Based on phytochemical screening results, the MCLI tested showed the presence of alkaloid compounds. This was seen from the formation of brown sediment at the bottom of the test tube from the MCLI. The alkaloid compound test resulting from using Wagner's reagents can be seen in Figure 7.

Several studies have indicated that alkaloids can inhibit various pro-inflammatory factors expression, such as histamine, lipid mediators, cytokines, and enzymes involved in the inflammatory response.²⁵ Some alkaloids direct mechanism to treated dermatomyositis rheumatoid arthritis, ankylosing spondylitis, myasthenia gravis, ankylosing spondylitis, systemic lupus erythematosus, Behcet's

disease, and other rheumatic immune diseases.²⁶

Conclusion

The investigation of this study showed that local *M. calabura* L. contains flavonoids, triterpenoids, and alkaloids and each has the potential to developing for alternative inflammation traetment by natural products.

Prospect Future Studies

The study revealed that the anti-inflammatory potential of the studied medicinal plants. The cold maceration extraction method can be use to obtain more secondary metabolites without damaging heat-sensitive compounds. *Muntingia calabura* L. leaves extract contains a wide range of non-polar, semi-polar, and polar secondary metabolite chemicals. Using the fractional method to separate some of these compounds based on their polarity is better. As anti-inflammatory drugs, further tests can be carried out in vivo and in vitro. By purifying and separating the beneficial components from these plant extracts, we could gain a better understanding of how they function and discover potential ingredients for developing new drugs.

Acknowledgement

Big appreciation for technical support from Prof. Dikdik Kurnia and Siti from Chemical Application Laboratory of Universitas Padjadjaran.

Funding

This work was supported by Research Funding from RKI and RPLK for Aziiz Mardanarian Rosdianto (Document No. 2213/UN6.3.1/TU.00/2023 and No. 1549/UN6.3.1/PT.00/2023).

Conflict of Interest

None declared.

References

1. Brokken MT, Head MJ, Boswell JC. Joint Disorders in Horses.pdf. MSD Veterinary Manual. Pet Owner Version. Published online 2019. <https://www.merckvetmanual.com/horse-owners/bone,-joint,-and-muscle-disorders-in-horses/joint-disorders-in-horses#>
2. Neundorf RH, Lowerison MB, Antonio M Cruz, Thomason JJ, McEwen BJ, Hurtig MB. Determination of the prevalence and severity of metacarpophalangeal joint osteoarthritis in Thoroughbred racehorses via quantitative macroscopic evaluation. *American Journal of Veterinary Research*. Published online 2010.
3. Flood J, Stewart AJ. Non-Steroidal Anti-Inflammatory Drugs and Associated Toxicities in Horses. *Animals*. 2022;12(21). doi:10.3390/ani12212939
4. Ansori ANM, Kharisma VD, Solikhah TI. Medicinal properties of muntingia calabura l.: A review. *Research Journal of Pharmacy and Technology*. 2021;14(8):4509-4512. doi:10.52711/0974-360X.2021.00784
5. Puspitasari, A. D. and Prayogo, L. S. (2016) 'Pengaruh waktu perebusan terhadap kadar flavonoid total daun kersen (Muntingia calabura). *Jurnal Inovasi Teknik Kimia*, 1(2), pp. 104–108.
6. Sumarni S, Sadino A, Sumiwi SA. Literature Review: Chemical Content and Pharmacological Activity of Kersen Leaf (Muntingia calabura L.). *Jurnal Farmasi Sains dan Praktis*. 2022;8(1):13-20. doi:10.31603/pharmacy.v8i1.3802
7. Howlader M, Ahmed S, Kubra K, Bhuiyan MK. Biochemical and phytochemical evaluation of Stevia rebaudiana. *Asian Journal of Medical and Biological Research*. 2016;2:121. doi:10.3329/ajmbr.v2i1.27577
8. Hoon, J. H. and Chung, B. M. (2020) 'Effects of extraction conditions on acrylamide / furan content , antioxidant

- activity , and sensory properties of cold brew coffee', *Food Science and Biotechnology*. 2020 Mar 30;29(8):1071-1080. doi: 10.1007/s10068-020-00747-1.
9. Prastiwi R, Siska, Nila M. *Parameter Fisikokimia dan Analisis Kadar Allyl Disulfide dalam Ekstrak Etanol 70% Bawang Putih (Allium sativum L.) dengan Perbandingan Daerah Tempat Tumbuh Parameter Physico-Chemical Parameters and Quantitative Analysis of Allyl Disulfide in Allium sativu*. Fakultas Farmasi dan Sains, Univ Muhammadiyah. Published online 2017:33.
10. Wahyuningsih R, Wiryosoendjoyo K. Uji Aktivitas Anti Jamur Ekstrak Infusa Daun Sirsak (*Annona muricata* L.) Terhadap *Candida albicans* Anti Fungus Activity Test for Soursop (*Annona muricata* L.) Leaf Infusion Extract Against *Candida albicans*. *Media Informasi Kesehatan*. 2019;6(2):167-176.
11. Pertiwi RD, Suwaldi, Martien R, Setyowati EP. Radical Scavenging Activity and Quercetin Content of *Muntingia calabura* L. Leaves Extracted by Various Ethanol Concentration. *Journal of Food and Pharmaceutical Sciences*. 2020;(September):1. doi:10.22146/jfps.581
12. Alara OR, Abdurahman NH, Ukaegbu CI. Extraction of phenolic compounds: A review. *Current Research in Food Science*. 2021;4(March):200-214. doi:10.1016/j.crfs.2021.03.011
13. Das AK, Islam MN, Faruk MO, Ashaduzzaman M, Dungani R. Review on tannins: Extraction processes, applications and possibilities. *South African Journal of Botany*. 2020;135:58-70. doi:10.1016/j.sajb.2020.08.008
14. Rodríguez De Luna SL, Ramírez-Garza RE, Serna Saldívar SO. Environmentally Friendly Methods for Flavonoid Extraction from Plant Material: Impact of Their Operating Conditions on Yield and Antioxidant Properties. *The Scientific World Journal*. 2020;2020. doi:10.1155/2020/6792069
15. Widyaningrum NR, Wahyuningsih SS, Priyono C. Antiinflammation activity of *Muntingia calabura* L. leaves ethanol, ethylacetate and chloroform extracts. *Natural Science: Journal of Science and Technology*. 2022;11(01). doi:10.22487/25411969.2022.v11.i01.15699
16. Zakaria ZA, Mahmood ND, Omar MH, Taher M, Basir R. Methanol extract of *Muntingia calabura* leaves attenuates CCl₄-induced liver injury: possible synergistic action of flavonoids and volatile bioactive compounds on endogenous defence system. *Pharmaceutical Biology*. 2019;57(1):335-344. doi:10.1080/13880209.2019.1606836
17. Elbatreek MH, Mahdi I, Ouchari W, Mahmoud MF, Sobeh M. Biomedicine & Pharmacotherapy Current advances on the therapeutic potential of pinocembrin : An updated review. *Biomedicine & Pharmacotherapy*. 2023;157:114032. doi:10.1016/j.biopha.2022.114032
18. Ahmed EA, Ibrahim H-IM, Khalil HE. Pinocembrin Reduces Arthritic Symptoms in Mouse Model via Targeting Sox4 Signaling Molecules. *Journal of Medicinal Food*. 2021 Mar;24(3):282-291. doi: 10.1089/jmf.2020.4862.
19. Yue B, Ren J, Yu Z, et al. Pinocembrin alleviates ulcerative colitis in mice via regulating gut microbiota, suppressing TLR4 / MD2 / NF- κ B pathway and promoting intestinal barrier. *Bioscience Report*. 2020 Jul 31; 40(7): BSR20200986
20. Gu X, Zhang Q, Du Q, Shen H, Zhu Z. Pinocembrin attenuates allergic airway inflammation via inhibition of NF- κ B pathway in mice. *International Immunopharmacology*. 2017 Dec;53:90-

95. doi: 10.1016/j.intimp.2017.10.005. Epub 2017 Oct 18.
21. National Center for Biotechnology Information. *PubChem Compound Summary for CID 68071, Pinocembrin*. Published 2023. Accessed August 28, 2023. <https://pubchem.ncbi.nlm.nih.gov/compound/68071>
22. Chairunnisa S, Wartini NM, Suhendra L. Pengaruh Suhu dan Waktu Maserasi terhadap Karakteristik Ekstrak Daun Bidara (*Ziziphus mauritiana* L.) sebagai Sumber Saponin. *Jurnal Rekayasa Dan Manajemen Agroindustri*. 2019;7(4):551. doi:10.24843/jrma.2019.v07.i04.p07
23. Labagu R, Asri ;, Naiu S, et al. Kadar Saponin Ekstrak Buah Mangrove (*Sonneratia alba*) dan Daya Hambatnya Terhadap Radikal Bebas DPPH Levels of Saponin in Magrove Fruit (*Sonneratia alba*) Extract and Its Inhibition Against DPPH Free Radical. *Jambura Fish Process Journal*. 2022;4(1):1-11. <http://ejurnal.ung.ac.id/index.php/jfpj/issue/archive>
24. Zhang X, Zhang S, Yang Y, Wang D, Gao H. Natural barrigenol-like triterpenoids: A comprehensive review of their contributions to medicinal chemistry. *Phytochemistry*. 2019;161:41-74. doi:10.1016/j.phytochem.2019.01.017
25. Souza CRM, Bezerra WP, Souto JT. Marine alkaloids with anti-inflammatory activity: Current knowledge and future perspectives. *Marine Drugs*. 2020;18(3). doi:10.3390/md18030147
26. Wei T, Xiaojun X, Peilong C. Biomedicine & Pharmacotherapy Magno fl orine improves sensitivity to doxorubicin (DOX) of breast cancer cells via inducing apoptosis and autophagy through AKT / mTOR and p38 signaling pathways. *Biomedicine & Pharmacotherapy*. 2020;121(June 2019):109139. doi:10.1016/j.biopha.2019.109139.