



Assessment of Iron Contamination in Groundwater of Catchment Area Water

Aïchetou Brahim Boutebib¹, Abdoulaye Demba N'diaye^{1,*}, Sid'Ahmed Baba Elhoumed², Bocar Kalidou M'Baye³, Youssef Aoulad El Hadj Ali⁴, Belkheir Hammouti^{5,*}, Bakari Mohamed Semega¹

¹ Unité de Recherche, Eau Pollution et Environnement, Département de Chimie, Faculté des Sciences et Techniques, Université de Nouakchott, Nouakchott, Mauritanie.

² Cellule Management Qualité, Service Laboratoire, Société Nationale d'Eau (SNDE), Nouakchott, Mauritanie.

³ Laboratoire de Chimie, Institut National de Recherches en Santé Publique, Nouakchott, Mauritanie

⁴ Département de Chimie, Faculté des Sciences, Université Abdelmalek Essadi, Mhannech II, 93002 Tétouan, Maroc

⁵ Euro-Mediterranean University of Fez, fez, Morocco

Correspondence: E-mail: abdouldemba@yahoo.fr; hammoutib@gmail.com

ABSTRACT

This study investigates the occurrence of iron (Fe) in the catchment area water in Derwiche from Hodh El Chargui in Mauritania. The study area was monitored during January and August. The occurrence of Fe in the groundwater showed significant spatial and temporal fluctuations. The relationship between Fe and some physicochemical parameters was also analyzed statistically using Pearson's correlation matrix. Firstly, the results suggested that the concentration of Fe was influenced by the dissolution of iron minerals. Secondly, the results show that the pH value was an important factor that influenced the Fe concentrations in the groundwater. The Water Quality Index (WQI) method was used to evaluate the suitability for human consumption. The results show that WQI is strongly influenced by Fe indicating that Fe removal would contribute to excellent well water. The Treatability Index (TI) is used to assess water quality. TI confirms the results obtained by the WQI. However, TI also shows that the pH is a parameter to be optimized for possible potabilization. These results show that there would be a close relationship between the recorded Fe levels and the pH. TI confirms the results obtained by Pearson's correlation matrix. Therefore, the presence of high Fe concentrations in groundwater in the study area could have a natural origin. The results recommended that different treatment techniques should be employed to purify groundwater before consumption. The primary focus should be cascaded toward Fe contamination.

ARTICLE INFO

Article History:

Submitted/Received 29 Mar 2023

First Revised 01 May 2023

Accepted 14 Jul 2023

First Available online 17 Jul 2023

Publication Date 01 Dec 2023

Keyword:

Groundwater,

Hodh El Chargui,

Iron,

Mauritania,

Quality.

1. INTRODUCTION

Groundwater is a vital natural resource essential for drinking, irrigation, industry, and other economic sectors (Bodrud-Doza et al., 2019). Because of being pathogen-free and available, groundwater is considered a prime source of potable water in most countries of the world (Rahman et al., 2021).

About 80% of the diseases in developing countries are related to contaminated water and the resulting death toll is as much as 10 million per year. Among many contaminants, iron (Fe) is present in chemicals derived from both natural sources and human activities (Tekerlekopoulou et al., 2013).

Fe is a fairly abundant element in rocks and is found in the form of silicates, oxides as well as hydroxides, carbonates, and sulfides. Fe exists in dissolved Fe^{2+} (ferrous ion) and suspended Fe^{3+} (ferric ion) forms (Zhang et al., 2020). They can occur naturally as a result of water–sediment interaction which induces their dissolution. Dissolution processes are controlled by several environmental factors, primarily redox conditions (Weng et al., 2007; McMahon & Chapelle, 2008; Palmucci et al., 2016). Fe is one of the important elements, used by many technologies, including hydrogen energy generation (Nandiyanto et al., 2023).

Although trace of Fe is indispensable to human survival, excessive exposure to them can damage human health. Excessive intake of Fe can lead to chronic intoxication (Zoni et al., 2007). Drinking water containing Fe at concentrations 3 mg/L or higher for a long time causes hemochromatosis and other effects (Weng et al., 2007). At the same time, the oxidation of Fe makes water turbid and odorous, and excessive Fe content in

groundwater also causes pipeline rust and blockage of wells.

Much attention has been paid to this problem (Benrabah et al., 2016; Ye et al., 2020). The World Health Organization (WHO)'s maximum threshold for Fe^{2+} in drinking water is 0.3 mg/L. The objectives of this study are to (1) investigate the contamination levels of Fe in the groundwater in catchment area water in Derwiche from Hodh El Chargui in Mauritania and (2) examine the factors that influence the Fe concentrations in the groundwater. Through this work, we offer relevant recommendations that the government could consider in establishing a set of cost-effective measures for groundwater remedy and management.

2. METHODS

2.1. Presentation of the Study Area

The Hodh El Chargui, a region located in the southeast of Mauritania at 1000 km from the capital city of Nouakchott (Figure 1), is characterized by a hard and rural environment. The temperature rises during the summer to 50 °C and drops in winter to about 20°C. Farmers and agro-pastoralists are the main economic actors. The boreholes are located in the Nema watershed at the level of the pumping station (Dhar-service). These boreholes are located about 150 km from Nema, this aquifer has an area of about 30 thousand km². The alimentation of this aquifer is mainly done by the direct infiltration of rainwater. It is important to note that there is no industrial production or other pollution source in the study area. The selected boreholes were DL2 (N16°50'17.5" and W 06°11'24.2"), DL6 (N16°50'37.4" and W 06°11'07.0"), and DL9 (N 16°51'03.3" and W 06°11'09.3").




Figure 1. Hodh El Chargui localisation.

2.2. Sampling Procedures

Water samples were collected from three sampling sites (Boreholes) scattered along the catchment area water in Derwiche from Hodh El Chargui. The samples were collected two times: the first sampling was done in January and the last in August corresponding to the rainy period. The sampling for the analysis of physico-chemical parameters was carried out at the selected site, using a 1.5 L polyethylene bottle. The collected samples were stored at 4°C in insulated boxes and transported to the laboratory.

2.3. Determination of Physicochemical Parameters

Methods for measurement of physicochemical parameters such as

Temperature (T), pH, Electrical Conductivity (EC), Total Dissolved Solid (TDS), nitrates (NO_3^-), chlorides (Cl^-), oxidizability (Organic Matter: (OM), and iron (Fe) are listed in **Table 1**. It is important to note that each sample was conducted in triplicate and the mean values are reported.

2.4. Calculation of Water Quality Index (WQI)

A WQI is considered a parameter that defined the composite effect attributed to diverse water quality parameters (Sahu & Sikdar, 2008) and was calculated using the Weighted Arithmetic Index method. The quality rating scale for each parameter q_i was calculated by using this expression:

$$q_i = \frac{C_i}{S_i} \times 100 \quad (1)$$

Table 1. Water quality parameters and measurement methods.

| Parameters | Unit | Measuring equipment and method analysis |
|------------------------------|-------|--|
| T | ° C | pH meter HI 991001 |
| pH | | pH meter HI 991001 |
| EC | µS/cm | Conductimeter HI98192 |
| TDS | mg/L | Conductimeter HI98192 |
| Cl ⁻ | mg/L | Volumetric dosage with AgNO ₃ |
| NO ₃ ⁻ | mg/L | Photometer Wagtech 7100 |
| Fe | mg/L | Photometer Wagtech 7100 |
| OM | mg/L | Hot oxidation by KMnO ₄ |

A quality rating scale (q_i) for each parameter is assigned by dividing its concentration (C_i) in each water sample by its respective standard (S_i) and the result is multiplied by 100. Relative weight (W_i) was calculated by a value inversely proportional to the recommended standard (S_i) of the corresponding parameter:

$$W_i = \frac{1}{S_i} \quad (2)$$

The overall WQI was calculated by aggregating the quality rating (q_i) with unit weight (W_i) linearly:

$$WQI = \sum_{i=1}^n (W_i \times q_i) \quad (3)$$

Generally, WQI was discussed for a specific and intended use of water. In this study, the WQI for drinking purposes is considered and the permissible WQI for drinking water is taken as 100:

$$\text{Overall WQI} = \frac{\sum_{i=1}^n q_i w_i}{\sum_{i=1}^n w_i} \quad (4)$$

The determined WQI values are classified into five classes as mentioned in **Table 2** (Yidana & Yidana, 2010).

2.5. Treatability Index (TI) and Borehole Classification Model

The TI is an indicator that is often used to assess water quality (Enitan-Folami *et al.*, 2019; Masindi & Foteinis, 2021). Besides, the TI is defined as the ratio of the Measured Value (MV) of the contaminant under study to its Maximum Allowed Limit

(MAL) for drinking water, as shown in Equation (5):

$$TI = \frac{MV}{MAL} \quad (5)$$

When the TI is > 1 , the water will require treatment to comply with the prescribed limits, since it is not suitable for human consumption. When the TI is < 1 , the water will require zero treatment to comply with the required limit, and it is suitable for human consumption.

Equation (5) also implies that a wide range of physicochemical parameters such as pH, TDS, NO_3^- , Cl^- , OM, and Fe should be examined to assess water quality and its suitability for drinking purposes. The quality of the water will then be determined by the parameter that exhibits the highest TI value, i.e. if one parameter has a TI value higher than unity then water will be unsuitable for human consumption, regardless of the values of the other parameters. Therefore, treatment will be required to lower all TI values below unity before water consumption. When the TI value is higher than unity, the water boards should either treat the water or not use it at all.

Specifically, to classify water quality in Derwiche from Hodh El Chargui, a classification index was developed. In this study, the WHO (2011) in http://www.who.int/water_sanitation_health. Accessed 21 February 2014 for drinking water, was used to assess groundwater's quality. Finally, the water quality was divided into four different classes, as proposed elsewhere (Enitan-Folami *et al.*, 2019).

Table 2. WQI Categories.

| Range | Quality |
|---------|-------------------------|
| <50 | Excellent water |
| 50-100 | Good water |
| 100-200 | Poor water |
| 200-300 | Very poor water |
| >300 | Unsuitable for drinking |

3. RESULTS AND DISCUSSION

3.1. Spatio-Temporal Variation of Fe

First, the physicochemical parameters such as Temperature, pH, EC, TDS, NO_3^- , Cl^- and OM studied in the catchment area water in Derwiche during January and August were represented in **Table 3**. The Temperature, pH, EC, TDS, NO_3^- , Cl^- and OM contents of water samples are within WHO for drinking water.

Concerning the spatio-temporal variation of Fe, Fe values range from 0.75 to 7 mg/L in

January and 0.4 to 2.8 mg/L during August. The highest Fe values are recorded at sites DL2 and DL6, where they reached 4.8 and 7 mg/L in January, respectively. However, the lower Fe values are recorded at DL9 in January (0.4 mg/L).

The decrease in Fe in August is probably due to the dilution effect of rainwater. The concentrations of Fe obtained in this study are very lower than those obtained by [Hossain et al., \(2013\)](#) (24.5 mg/L) and [Rusydi et al., \(2021\)](#) (252 mg/L). The Fe concentration limit is set at 0.3 mg/L.

Table 3. Statistical summary of the physicochemical parameters of the catchment area water in Derwiche.

| | DL2 | DL6 | DL9 |
|-----------------|------------------|-------------------|-------------------|
| T | 25.80 \pm 1.48 | 25.55 \pm 0.80 | 25.45 \pm 0.38 |
| pH | 6.00 \pm 0.84 | 6.00 \pm 0.50 | 5.95 \pm 0.63 |
| EC | 68.15 \pm 3.51 | 59.10 \pm 30.77 | 63.65 \pm 11.68 |
| TDS | 25.5 \pm 10.97 | 29.76 \pm 15.07 | 31.4 \pm 6.05 |
| Cl^- | 21.30 \pm 3.89 | 21.40 \pm 3.94 | 21.42 \pm 3.94 |
| NO_3^- | 1.02 \pm 0.73 | 1.08 \pm 0.31 | 1.00 \pm 0.13 |
| OM | 1.09 \pm 0.68 | 1.21 \pm 0.75 | 1.25 \pm 0.31 |




Figure 2. Spatio-temporal variation in Fe of Borehole waters from the catchment area water in Derwiche.

3.2. Relationship Between Fe and Other Physicochemical Parameters

The degree estimated by Pearson's connection coefficients for the January and August catchment area water in Derwiche is exhibited in **Tables 4** and **5**, respectively. A strong and positive correlation was found between Fe and OM ($r=0.986$) while a strong and negative correlation was found between Fe and pH ($r=-0.952$) (**Table 5**) during January. This means that the higher Fe concentrations might have been the result of the interaction of underground oxidized iron minerals with OM present and can be due to the dissolution of Fe_2CO_3 present in rocks at a low pH. According to [Harvey and Fuller \(1998\)](#), a decrease in pH will accelerate the dissolution of carbonates and hydroxides. Fe bound to carbonates will be released into the water. The Fe–O bonds in the crystal lattices of Fe-bearing minerals will readily be destroyed in an acidic environment to release structural Fe or interlayer Fe and promote Fe dissolution. Similar observations are reported by [\(Applin & Zhao 1989; Mondal et al., 2010\)](#).

A strong and positive correlation was found between Fe–TDS (0.664) and Fe – EC

(0.644) (**Tables 4-5**) during January, suggesting that the concentration of Fe in catchment area water in Derwiche well was influenced by dissolution minerals in groundwater. Similar observations are reported by [Akbar et al., \(2021\)](#).

A negative correlation was found between Fe^{2+} and NO_3^- during January ($r=-0.267$), probably because nitrate is the second preferential electron acceptor after oxygen. Nevertheless, the negative correlations noted between Fe^{2+} and NO_3^- imply a reduction process that encourages Fe dissolution ([Robertson & Thamdrup 2017; Benkaddour et al., 2021](#)).

3.3. Assessment of the Water Quality Using WQI

The parameters pH, TDS, Cl^- , NO_3^- , OM, and Fe, were taken into justification for the calculation of the WQI value for each sampling site in January and August. Furthermore, the WHO (limits were used to compute the WQI). Since the high values of borehole waters, the WQI values were categorized into two categories:

- (i) Including Fe,
- (ii) Without Fe

Table 4. Pearson correlation matrix of January.

| Variables | Fe | pH | EC | TDS | Cl | NO_3^- | MO |
|-----------------|----------|----------|----------|----------|----------|-----------------|----------|
| Fe | 1 | | | | | | |
| pH | -0,952 | 1 | | | | | |
| EC | 0,644 | -0,848 | 1 | | | | |
| TDS | 0,664 | -0,862 | 1,000 | 1 | | | |
| Cl | 0,986 | -0,990 | 0,763 | 0,781 | 1 | | |
| NO_3^- | -0,267 | -0,042 | 0,565 | 0,543 | -0,101 | 1 | |
| MO | 0,986 | -0,990 | 0,763 | 0,781 | 1,000 | -0,101 | 1 |

Table 5. Pearson correlation matrix of August.

| Variables | Fe | pH | EC | TDS | Cl | NO_3^- | MO |
|-----------------|----------|----------|----------|----------|----------|-----------------|----------|
| Fe | 1 | | | | | | |
| pH | 1,000 | 1 | | | | | |
| EC | -0,498 | -0,481 | 1 | | | | |
| TDS | -0,572 | -0,556 | 0,996 | 1 | | | |
| Cl | -0,994 | -0,996 | 0,399 | 0,478 | 1 | | |
| NO_3^- | 0,884 | 0,893 | -0,036 | -0,123 | -0,931 | 1 | |
| MO | -0,963 | -0,969 | 0,248 | 0,331 | 0,987 | -0,977 | 1 |

In the case of the Fe surveyed is included (**Table 6**) in the calculation of the WQI of the catchment area water in Derwiche, the results showed that the lowest WQI values were recorded at DL9 for all sampling months, indicating poor and very poor water quality. However, the highest WQI values were recorded at DL6 and DL2 indicating unsuitable for drinking water for all sampling months.

Without Fe (**Table 7**), it was observed that all WQI values are below 50, which means that the water is excellent for drinking in all boreholes from catchment area water in Derwiche for all sites and for all sampling months. These results obtained by WQI indicate that Fe removal would contribute to excellent quality water.

3.4. Assessment of the Water Quality Using TI

The parameters pH, TDS, Cl⁻, NO₃⁻, OM, and Fe, were taken into justification for the calculation of the TI value for each sampling site in January and August. Since the high values of borehole waters Fe, the TI values were categorized into three categories:

- (i) Including Fe,
- (ii) Without Fe
- (iii) Without Fe and pH

In the case of the Fe surveyed is included (**Table 8**) in the calculation of the TI of the catchment area water in Derwiche, the results showed that if the Fe has a TI value higher than unity then water will be unsuitable for human consumption for all sampling sites and months. However once without Fe (**Table 9**), it was observed that pH has a TI value between 0.75 and 1, indicating water will be acceptable quality for human consumption for all sampling sites and months.

Without Fe and pH (**Table 10**), the results showed that all studied parameters have a TI value higher than unity then water will be of excellent quality for human consumption for all sampling sites and months.

The TI confirms the results obtained by the WQI. However, TI also shows that the pH is an important factor that influenced the Fe concentrations in the groundwater. So, the TI confirms the results obtained by the correlation matrix.

Table 6. WQI values of different sampling sites from catchment area water in Derwiche including Fe.

| Boreholes | January | | August | |
|-----------|---------|-------------------------|--------|-------------------------|
| | WQI | Water type | WQI | Water type |
| DL2 | 1449.3 | Unsuitable for drinking | 846.4 | Unsuitable for drinking |
| DL6 | 2111.8 | Unsuitable for drinking | 1237.7 | Unsuitable for drinking |
| DL9 | 229.9 | Very poor water | 124.7 | Poor water |

Table 7. WQI values of different sampling sites from catchment area water in Derwiche without Fe.

| Boreholes | January | | August | |
|-----------|---------|-----------------|--------|-----------------|
| | WQI | Water type | WQI | Water type |
| DL2 | 4.53 | Excellent water | 37.49 | Excellent water |
| DL6 | 4.86 | Excellent water | 37.10 | Excellent water |
| DL9 | 4.20 | Excellent water | 44.02 | Excellent water |

Table 8. TI values of different sampling sites from catchment area water in Derwiche including Fe.

| | Sites | pH | TDS | Cl ⁻ | NO ₃ ⁻ | OM | Fe | Class |
|---------|-------|------|-------|-----------------|------------------------------|-------|-------|-------|
| January | DL2 | 0.87 | 0.071 | 0.051 | 0.014 | 0.26 | 16 | 4 |
| | DL6 | 0.86 | 0.087 | 0.060 | 0.017 | 0.32 | 23.33 | 4 |
| | DL9 | 0.87 | 0.074 | 0.057 | 0.018 | 0.19 | 2.5 | 4 |
| August | DL2 | 0.73 | 0.031 | 0.071 | 0.027 | 0.178 | 9.33 | 4 |
| | DL6 | 0.74 | 0.032 | 0.071 | 0.026 | 0.164 | 23.67 | 4 |
| | DL9 | 0.72 | 0.052 | 0.071 | 0.018 | 0.32 | 1.33 | 4 |

Table 9. TI values of different sampling sites from catchment area water in Derwiche without Fe.

| | Sites | pH | TDS | Cl ⁻ | NO ₃ ⁻ | OM | Class |
|---------|-------|------|-------|-----------------|------------------------------|-------|-------|
| January | DL2 | 0.87 | 0.071 | 0.051 | 0.014 | 0.26 | 3 |
| | DL6 | 0.86 | 0.087 | 0.060 | 0.017 | 0.32 | 3 |
| | DL9 | 0.87 | 0.074 | 0.057 | 0.018 | 0.19 | 3 |
| August | DL2 | 0.73 | 0.031 | 0.071 | 0.027 | 0.178 | 3 |
| | DL6 | 0.74 | 0.032 | 0.071 | 0.026 | 0.164 | 3 |
| | DL9 | 0.72 | 0.052 | 0.071 | 0.018 | 0.32 | 3 |

Table 10. TI values of different sampling sites from catchment area water in Derwiche without Fe and pH.

| | Sites | TDS | Cl ⁻ | NO ₃ ⁻ | OM | Class |
|---------|-------|-------|-----------------|------------------------------|-------|-------|
| January | DL2 | 0.071 | 0.051 | 0.014 | 0.26 | 1 |
| | DL6 | 0.087 | 0.060 | 0.017 | 0.32 | 1 |
| | DL9 | 0.074 | 0.057 | 0.018 | 0.19 | 1 |
| August | DL2 | 0.031 | 0.071 | 0.027 | 0.178 | 1 |
| | DL6 | 0.032 | 0.071 | 0.026 | 0.164 | 1 |
| | DL9 | 0.052 | 0.071 | 0.018 | 0.32 | 1 |

4. CONCLUSION

This study investigates the occurrence of Fe in the catchment area water in Derwiche from Hodh El Chargui in Mauritania. The occurrence of Fe in the groundwater showed significant spatial and temporal fluctuations and was possibly affected by the change in environmental conditions within the aquifer. The relationship between Fe and some physicochemical parameters was also analyzed statistically using Pearson's correlation matrix. Firstly, the results suggested that the concentration of Fe was influenced by the dissolution of iron minerals. Secondly, the results show that the pH value was an important factor that influenced the Fe concentrations in the groundwater. The WQI is used to identify

the influence of Fe contamination. These results indicate that Fe removal would contribute to excellent well water. The TI confirms the results obtained by the WQI. However, TI also shows that the pH is a parameter to be optimized for possible potabilization.

These results show that there would be a close relationship between the recorded Fe levels and the pH. Therefor, the TI confirms the results obtained by Pearson's correlation matrix. Therefore, the presence of high Fe concentrations in groundwater in the study area could have a natural origin. From the result of the present study, it is highly recommended that different treatment techniques should be employed to purify groundwater before consumption. Adsorption is one of the most widely applied

techniques for Fe and other pollutants removal.

5. AUTHORS' NOTE

The authors declare that there is no conflict of interest regarding the publication of this article. The authors confirmed that the paper was free of plagiarism.

6. REFERENCES

Akbar, N. A., Aziz, H. A., and Adlan, M. N. (2021). Characteristic of groundwater well quality using bivariate analysis: A case study at USM Engineering Campus, Penang. *IOP Conference Series: Earth and Environmental Science*, 646(1), 012060.

Benkaddour, R., Merimi, I., Szumiata, T., and Hammouti, B. (2020). Nitrates in the groundwater of the Triffa plain Eastern Morocco. *Materials Today: Proceedings*, 27, 3171-3174.

Benrabah, S., Attoui, B., and Hannouche, M. (2016). Characterization of groundwater quality destined for drinking water supply of Khenchela City (eastern Algeria). *Journal of Water and Land Development*, 30(1), 13-20.

Bodrud-Doza, M., Bhuiyan, M. A. H., Islam, S. D. U., Quraishi, S. B., Muhib, M. I., Rakib, M. A., and Rahman, M. S. (2019). Delineation of trace metals contamination in groundwater using geostatistical techniques: A study on Dhaka City of Bangladesh. *Groundwater for Sustainable Development*, 9, 100212.

Enitan-Folami, A. M., Mutileni, N., Odiyo, J. O., Swalaha, F. M., and Edokpayi, J. N. (2020). Hydrochemical, bacteriological assessment, and classification of groundwater quality in Thulamela Municipality, South Africa: Potential health risk. *Human and Ecological Risk Assessment: An International Journal*, 26(8), 2044-2058.

Harvey, J. W., and Fuller, C. C. (1998). Effect of enhanced manganese oxidation in the hyporheic zone on basin-scale geochemical mass balance. *Water Resources Research*, 34(4), 623-636.

Hossain, D., Islam, M. S., Sultana, N., and Tusher, T. R. (2013). Assessment of iron contamination in groundwater at Tangail Municipality, Bangladesh. *Journal of Environmental Science and Natural Resources*, 6(1), 117-121.

Masindi, V., and Foteinis, S. (2021). Groundwater contamination in sub-Saharan Africa: Implications for groundwater protection in developing countries. *Cleaner Engineering and Technology*, 2, 100038.

McMahon, P. B., and Chapelle, F. H. (2008). Redox processes and water quality of selected principal aquifer systems. *Groundwater*, 46(2), 259-271.

Mondal, N. C., Singh, V. S., Puranik, S. C., and Singh, V. P. (2010). Trace element concentration in groundwater of Pesarlanka Island, Krishna Delta, India. *Environmental Monitoring and Assessment*, 163, 215-227.

Nandiyanto, A.B.D., Ragadhita, R., Fiandini, M., Al Husaeni, D.N., and Aziz, M. (2023). The role of iron oxide in hydrogen production: Theory and bibliometric analyses. *Moroccan Journal of Chemistry*, 11(4), 897.

Palmucci, W., Rusi, S., and Di Curzio, D. (2016). Mobilisation processes responsible for iron and manganese contamination of groundwater in Central Adriatic Italy. *Environmental Science and Pollution Research*, 23, 11790-11805.

Rahman, M., Tushar, M. A. N., Zahid, A., Mustafa, M. G., Siddique, M. A. M., and Ahmed, K. M. (2021). Spatial distribution of manganese in groundwater and associated human health risk in the southern part of the Bengal Basin. *Environmental Science and Pollution Research*, 28, 41061-41070.

Robertson, E. K., and Thamdrup, B. (2017). The fate of nitrogen is linked to iron (II) availability in a freshwater lake sediment. *Geochimica et Cosmochimica Acta*, 205, 84-99.

Rusydi, A. F., Onodera, S. I., Saito, M., Ioka, S., Maria, R., Ridwansyah, I., and Delinom, R. M. (2021). Vulnerability of groundwater to iron and manganese contamination in the coastal alluvial plain of a developing Indonesian City. *SN Applied Sciences*, 3, 1-12.

Sahu, P., and Sikdar, P. K. (2008). Hydrochemical framework of the aquifer in and around East Kolkata Wetlands, West Bengal, India. *Environmental Geology*, 55, 823-835.

Tekerlekopoulou, A. G., Pavlou, S., and Vayenas, D. V. (2013). Removal of ammonium, iron and manganese from potable water in biofiltration units: A review. *Journal of Chemical Technology and Biotechnology*, 88(5), 751-773.

Weng, H. X., Qin, Y. C., and Chen, X. H. (2007). Elevated iron and manganese concentrations in groundwater derived from the Holocene transgression in the Hang-Jia-Hu Plain, China. *Hydrogeology Journal*, 15, 715-726.

Ye, X., Cui, R., Wang, L., and Du, X. (2020). The influence of riverbank filtration on regional water resources: A case study in the second Songhua River Catchment, China. *Water Supply*, 20(4), 1425-1438.

Yidana, S. M., and Yidana, A. (2010). Assessing water quality using water quality index and multivariate analysis. *Environmental Earth Sciences*, 59, 1461-1473.

Zhang, Z., Xiao, C., Adeyeye, O., Yang, W., and Liang, X. (2020). Source and mobilization mechanism of iron, manganese and arsenic in groundwater of Shuangliao City, Northeast China. *Water*, 12(2), 534.

Zoni, S., Albini, E., and Lucchini, R. (2007). Neuropsychological testing for the assessment of manganese neurotoxicity: A review and a proposal. *American Journal of Industrial Medicine*, 50(11), 812-830.