



Original Research

Design of primer *Ipomoea batatas* chloroplast gene matK

Syamsurizal^{1*}, Ardi¹, Des M¹, Resti Fevria¹, Yusni Atifah¹, Elsa Badriyya², Afifatul Achyar¹

¹Department of Biology, Faculty of Mathematics and Natural Sciences, Universitas Negeri Padang, Padang, Indonesia, 25131

²Department of Pharmacology and Clinical Pharmacy, Faculty of Pharmacy, Universitas Andalas, Padang, Indonesia, 25166

*corresponding author

E-mail address: syam_unp@fmipa.unp.ac.id

Article Info

Article history:

Received 15 January 2021

Received in revised form

20 February 2021

Accepted 25 March 2021

Available online 30 May 2021

Keywords:

Ipomoea batatas

Plant DNA barcoding

matK

How to cite:

Syamsurizal, S., Ardi, A., M, D., Fevria R.,

Atifah, Y., Badriyya, E., Achyar, A. 2021.

Design of primer *Ipomoea batatas*

chloroplast gene matK. *Tropical Genetics*

1(1): 12-16.

Abstract

Sweet potato varieties (*Ipomoea batatas*) Pucuk Hitam Panyalaian and Madu Pucuk Hitam Panyalaian have the potential to be a superior commodity in West Sumatra, but the information of genetic diversity is very limited. Data on the diversity of sweet potato germplasm is beneficial in the selection of plants to obtain superior cultivars in plant breeding. The aim of the study was to find specific primers for the study of sweet potato genetic markers using *I. batatas* chloroplast genes matK. DNA extraction from sweet potato young leaves, designing forward and reverse primers *I. batatas* chloroplast genes matK, PCR, sequencing, bioinformatics analysis and species identification by comparing the NCBI database. Plant DNA barcoding PCR using designed primers matK were successfully resulting single DNA band in different amplicon size in some samples. This indicating that the designed primers used were able to distinguish variation in one species.

Copyright © 2021. The Authors. This is an open access article under the CC BY-NC-SA license (<http://creativecommons.org/licenses/by-nc-sa/4.0/>).

Introduction

Sweet potatoes have good prospects and opportunities to ensure food security, especially if rice and maize production does not meet the food needs of the community. Sweet potatoes are a food source that is not foreign to most Indonesian people. The ease of getting it makes people like to eat foods made from sweet potatoes. In the market you can find sweet potatoes of various colors, ranging from white, pale yellow, orange, to purple.

Many benefits can be obtained from this sweet potato. Sweet potatoes are rich in antioxidants (Pochapski et al., 2011), the content is higher when the color of sweet potatoes gets thicker (Cai et al., 2016). Sweet potatoes are also useful for helping the

body's immunity, because it is rich in beta carotene, sweet potatoes also contain other nutrients such as vitamin C, B complex, iron, and phosphorus (Bartke et al., 2006). This makes sweet potatoes a great immune booster. Besides sweet potatoes are good for overcoming inflammation, overcome asthma, reduce inflammation of the joints, help digestion and a source of carbohydrates for people with diabetes (Jones and de Brauw, 2015). Sweet Potato also find to have antidiabetic and anticancer activity (Mohanraj and Sivasankar, 2014).

Sweet potatoes are widely used and cultivated in West Sumatra, but are still traditional and have not been certified as superior and healthy. The West Sumatra sweet potato production centers include Bukittinggi, Batusangkar and Agam.

Therefore, it is necessary to develop superior standardized sweet potatoes and certified organic sweet potatoes. Strategic efforts to obtain such products require analysis of molecular biology to obtain phylogeny data and tissue culture propagation techniques (Badriyya and Achyar, 2020).

Analysis of molecular biology of sweet potato genetic diversity in West Sumatra was carried out using the DNA barcoding plant method (Fazekas et al., 2012). DNA barcoding is a standardized approach to identify plants and animals by using short gene sequences taken from a standardized portion of the genome, called DNA barcodes. The DNA sequence for barcoding must be able to distinguish each species, but it has a primer annealed region that is conserved or universal for all species. In plants, finding efficient DNA barcode candidates is quite challenging because of the high diversity of plant genomes (Wattoo et al., 2016).

Based on the recommendation of consortium for the Barcode of Life (BOLD) in plant working group, the unique DNA barcode for plant identification is the chloroplast matK (maturase K). These genes meet the criteria needed by a DNA barcode, which is short ± 1500 bp and ± 1400 bp respectively, has a conserved area for universal primer but still capable to distinguish each species. The genes found in the chloroplast are inherited maternally and recoverable even from herbarium samples (Hollingsworth et al., 2011).

In this study, we designed primer for *I. batatas* identification based on matK chloroplast genes as DNA barcode to discover more about genetic diversity of sweet potatoes in West Sumatra.

Materials and Methods

This research is a descriptive study that describes the design results primers used to detect the presence of matK chloroplast genes as DNA barcode. Identify the matK chloroplast genes was performed using the PCR method. Primers are designed based on the *Ipomoea batatas* cultivar Xushu 18

chloroplast genome sequence using the Geneious program. In this study, a pair of primer were used and the PCR product PCR was analyzed by 1.5% agarose gel electrophoresis (Vivantis).

Plant materials

Five varieties of sweet potatoes were studied, four of which were varieties cultivated by farmers in West Sumatra namely Pucuk Hitam Panyalaian, Madu Pucuk Hitam Panyalaian, Madu Papua dan 100 Hari, while one sample was wild sweet potato. The morphology of these varieties were observed and described.

Primer design

Primers were designed using Geneious 7.0.6 (<https://www.geneious.com>) based on *Ipomoea batatas* cultivar Xushu 18 chloroplast genome sequence (NCBI Reference Sequence: NC_026703.1, nucleotides 1.977-3.485 for matK) that were multiple aligned with other species of *Ipomoea* genus such as *I. tabascanana* (NC_041207.1), *I. ramosissima* (NC_041205.1), *I. x leucantha* (NC_041208.1), *I. splendor-sylvae* (NC_041206.1). Primers specificity were checked by PrimerBLAST and then synthesized by Macrogen, Inc. in South Korea.

DNA extraction

Young leaves at the 3 uppermost nodes were collected and stored in -20°C until the day of DNA extraction. DNA was extracted using simplified SDS protocol demonstrated by (Kiptantiyawati et al., 2014) with slightly modifications. The quality and quantity of DNA was analysed using NanoDrop spectrophotometer (Thermo Scientific). The isolated DNA was stored at -20°C .

matK PCR

PCR conditions of matK amplification was identification. Each PCR tubes were

contained of 12.5 µL of 2 x PCR buffer for KOD FX Neo, 5 µL of 2 mM dNTPs, 0.5 µL of KOD FX Neo (1.0U/µl) TOYOBO (Catalog number KFX-201), 0.75 µL of 10 mM forward primer and reverse primer, 1 µL of DNA template (100 ng/µL) and a certain volume of PCR grade water to get 25 µL final reaction volume. PCR was performed using C1000 thermal cyclor (Bio-Rad) and the cycling conditions were initial denaturation at 94°C for 3 min followed by 35 cycles of denaturation at 98°C for 10 s, annealing at 63°C for 30 s and extension at 68°C for 1 min, and followed by final extension at 68°C for 5 min. The PCR products were separated and visualized in 1.5% agarose gel staining with GelRed (Biotium).

Sequencing and bioinformatic analysis

matK PCR product of each samples were sequenced using Sanger Sequencing methods

by 1st BASE DNA Sequencing Services. Raw sequencing data from forward and reverse primer of each gene were contig using Geneious 7.0.6 (<https://www.geneious.com>) and then BLAST in NCBI.

Results and Discussion

Biology molecular analysis started with bioinformatic study of gene of interest based on *Ipomoea batatas* cultivar Xushu 18 chloroplast genome sequence (NCBI Reference Sequence: NC_026703.1). Designed primer used in this study were matK-F (5'-CAC TTG CTC ATG ATC GTG GTT-3') and matK-R (5'-CGT TGA GGG GGT TAG TGG AA-3'). Primers detailed information generated by Geneious 7.0.6 was described in Table 1.

Table 1. Designed primers information generated by Geneious 7.0.6.

Gene target	Primer details	DNA fold	Amplicon size*
matK	<p>matK-F Sequence (5' to 3'): CACTTGCTCATGATGTTGGTT Type: Primer Length: 21 %GC: 47.6 Hairpin Tm: None Self Dimer Tm: 14.9 Tm: 58.9 created by: primer3</p>		932 bp
	<p>matK-R Sequence (5' to 3'): CGTTGAGGGGTTAGTGGAA Type: Primer Length: 20 %GC: 55.0 Hairpin Tm: None Self Dimer Tm: None Tm: 59.3 created by: primer3</p>		

*The amplicon size was determined according to *Ipomoea batatas* cultivar Xushu 18 chloroplast genome sequence (NCBI Reference Sequence: NC_026703.1).

Some of cultivated sweet potatoes in West Sumatra studied here were Pucuk Hitam Panyalaian, Madu Pucuk Hitam Panyalaian, Madu Papua dan 100 Hari (Fig. 1). Young leaves at the 3 uppermost nodes were collected and processed for total genomic DNA extraction.

DNA extraction process from plants sometimes are quite challenging because of the presence of plant metabolites such as polysaccharide, polyphenols and protein contaminants that will be the inhibitors in

downstream processes (Sharma et al., 2008). Simplified SDS protocol described by (Kiptantiyawati et al., 2014) offered non-hazardous, rapid and inexpensive DNA extraction process. Quality of DNA extracted from five varieties of sweet potatoes using simplified SDS protocol were sufficient (260/280 ratio average 1.6) for amplification matK (Table 2), except for Pucuk Hitam Panyalaian variety that had lowest value of 260/280 ratio indicating poor purity.



Fig. 1. Morphology of four varieties of cultivated sweet potatoes in West Sumatra. **A:** Leaves and tuber of *Pucuk Hitam Panyalaian*. **B:** Leaves and tuber of *Madu Pucuk Hitam Panyalaian*. **C:** Leaves and tuber of *Madu Papua*. **D:** Leaves and tuber of *100 Hari*.

Table 2. Quantity and quality of sweet potatoes DNA extracts

Sweet potatoes variety	DNA yield (µg)	260/280 Ratio
Pucuk Hitam Panyalaian	30.2	1.51
Madu Pucuk Hitam Panyalaian	6.82	1.66
Madu Papua	6.29	1.55
100 Hari	90.45	1.65
Wild sweet potato	35.07	1.63

Plant DNA barcoding PCR using primers matK-F and matK-R were successfully resulting single DNA band in expected amplicon size approximately 900s bp for samples Madu Pucuk Hitam Panyalaian and Madu Papua, except for sample 100 Hari which gives the results of a lower-sized DNA band, approximately 700s bp. Unfortunately, there were no DNA band observed in PCR product of Pucuk Hitam Panyalaian and wild sweet potato. Low purity of DNA samples is thought to be the cause of PCR unsuccessful.

In plant DNA barcoding PCR, quality of DNA plays an important role. Sometimes, the presence of metabolites in plants effect DNA quality during extraction and even closely related species may require different DNA

extraction procedure (Khanuja et al., 1999). This explanation answers the question why some samples did not show the expected results.




Fig. 2. Electropherogram of matK PCR products. Label 1: Madu Pucuk Hitam; 2: Hitam Pucuk; 3: Madu Papua; 4: 100 Hari; 5: Wild sweet potato; L100bp: DNA ladder 100 bp.

Confirmation of matK amplicons of each samples were sequenced using Sanger methods by 1st Base Sequencing Services (Fig. 2). Different amplicon size resulted in some samples indicating that the designed primers used were able to distinguish variation in one species. Plant DNA barcoding

gives contribution in two aspects in biological science those are to provide insight into species level taxonomies and to assist in the process of identifying unknown specimens to known species (Hollingsworth et al., 2011).

Conclusions

Plant DNA barcoding PCR using designed primers matK was successfully resulting single DNA band in different amplicon size in some samples. This indicating that the designed primers used were able to distinguish variation in one species.

Acknowledgments

Profound Thanks towards Research Institution and Community Service (LPPM) of Universitas Negeri Padang, and this study was supported by PNPB Universitas Negeri Padang, in the scheme of Hibah Penelitian Unggulan Perguruan Tinggi Dasar (PUPTD), Contract Number: 927/UN35.13/LT/2019 Fiscal Year 2019.

References

- Badriyya, E., & Achyar, A. (2020). Primer Construction to detect SNP rs11196205 Transcription Factor 7 Like 2 (TCF7L2) Using Amplification Refractory Mutation System (ARMS) PCR to detect Type-2 Diabetes Mellitus. *Bioscience*, 4(2), 151. <https://doi.org/10.24036/0202042108497-0-00>.
- Bartke, N., Fischbeck, A., & Humpf, H.-U. (2006). Analysis of sphingolipids in potatoes (*Solanum tuberosum* L.) and sweet potatoes (*Ipomoea batatas* (L.) Lam.) by reversed phase high-performance liquid chromatography electrospray ionization tandem mass spectrometry (HPLC-ESI-MS/MS). *Molecular Nutrition & Food Research*, 50(12), 1201–1211. <https://doi.org/10.1002/mnfr.200600140>.
- Cai, Z., Qu, Z., Lan, Y., Zhao, S., Ma, X., Wan, Q., Jing, P., & Li, P. (2016). Conventional, ultrasound-assisted, and accelerated-solvent extractions of anthocyanins from purple sweet potatoes. *Food Chemistry*, 197, 266–272. <https://doi.org/10.1016/j.foodchem.2015.10.110>.
- Fazekas, A., Kuzmina, M., Newmaster, S., & Hollingsworth, P. (2012). DNA Barcoding Methods for Land Plants. *Methods in Molecular Biology (Clifton, N.J.)*, 858, 223–252. https://doi.org/10.1007/978-1-61779-591-6_11.
- Hollingsworth, P. M., Graham, S. W., & Little, D. P. (2011). Choosing and using a plant DNA barcode. *PLoS ONE*, 6(5). <https://doi.org/10.1371/journal.pone.0019254>
- Jones, K. M., & de Brauw, A. (2015). Using Agriculture to Improve Child Health: Promoting Orange Sweet Potatoes Reduces Diarrhea. *World Development*, 74, 15–24. <https://doi.org/10.1016/j.worlddev.2015.04.007>.
- Khanuja, S., Shasany, A., Darokar, M., & Kumar, S. (1999). Rapid Isolation of DNA from Dry and Fresh Samples of Plants Producing Large Amounts of Secondary Metabolites and Essential Oils. *Plant Molecular Biology Reporter - PL MOL BIOL REP*, 17, 74. <https://doi.org/10.1023/A:1007528101452>.
- Kiptantiyawati, N., Syarifah, N. L., Maulia, P., Sulistiyowati, E., Putra, M. R., Yudiansyah, Khumaida, N., & Ardie, S. (2014). Sample preservation procedures and simple dna isolation protocols for the tuberous crop, cassava (*Manihot esculenta* crantz.). *Asia-Pacific Journal of Molecular Biology and Biotechnology*, 22, 164–170.
- Mohanraj, R., & Sivasankar, S. (2014). Sweet Potato (*Ipomoea batatas* [L.] Lam)-A valuable medicinal food: A review. *Journal of Medicinal Food*, 17(7), 733–741.
- Pochapski, M. T., Fosquiera, E. C., Esmerino, L. A., dos Santos, E. B., Farago, P. V., Santos, F. A., & Groppo, F. C. (2011). Phytochemical screening, antioxidant, and antimicrobial activities of the crude leaves' extract from *Ipomoea batatas* (L.) Lam. *Pharmacognosy Magazine*, 7(26), 165.
- Sharma, K., Mishra, A. K., & Misra, R. S. (2008). A simple and efficient method for extraction of genomic DNA from tropical tuber crops. *African Journal of Biotechnology*, 7(8), 1018–1022. <https://doi.org/10.4314/ajb.v7i8.58596>.
- Wattoo, J. I., Saleem, M. Z., Shahzad, M. S., Arif, A., & Hameed, A. (2016). rbcl matK. *Journal of Biological Sciences*, 4(1), 3–7.