



# Utilization of Garlic (*Allium sativum*) as a Functional Supplement to Enhance the Health and Performance of Cultured Fish: A Systematic Literature Review

Sheny Permatasari<sup>1\*</sup>

<sup>1</sup> Department of Technology and Management of Applied Aquaculture, School of Vocational Studies, IPB University, Indonesia

\*Correspondence E-mail: [shenypemata@apps.ipb.ac.id](mailto:shenypemata@apps.ipb.ac.id)

## ARTICLE INFO

### Article History:

Received 7 November 2025

Revised 20 December 2025

Accepted 21 December 2025

Published 23 December 2025

### Keywords:

*Allium sativum*,

*Aquaculture*,

*Fish growth*,

*Garlic*,

*Immunity*.

## ABSTRACT

**Background:** The increasing demand for aquaculture production requires the use of foods that not only meet nutritional needs but also promote fish health. The use of natural food additives has become an important alternative to reduce dependence on synthetic antibiotics. One of the most extensively studied natural food additives is garlic (*Allium sativum*). Garlic has been found to eliminate major pathogenic bacteria, strengthen immunity, improve health status, enhance growth, and increase flesh quality in freshwater fish. With its multifunctional properties, garlic represents a promising functional feed additive to enhance growth performance, food utilization, and stress resistance in cultured fish.

**Aims:** This article was prepared using a systematic literature-review approach on the effects of garlic supplementation in aquaculture feeds.

**Methods:** A systematic literature search was conducted for articles published within the last ten years in Scopus- or SINTA-indexed journals. The review included: (1) experimental studies investigating the effects of garlic supplementation in fish feed or rearing media; (2) studies involving growth, physiological, immune, or water quality parameters; and (3) articles focusing on aquaculture species such as tilapia (*Oreochromis niloticus*), common carp (*Cyprinus carpio*), rohu (*Labeo rohita*), rainbow trout (*Oncorhynchus mykiss*), and groupers (*Epinephelus* spp.).

**Results:** The evidence has been shown that garlic can enhance growth performance, food efficiency, and immune response in fish, particularly in species such as tilapia, carp, rainbow trout, rohu and grouper. The effective dosage generally ranges between 0.5–3% garlic powder in food, with variations depending on species and application methods.

**Conclusion/ Recommendation:** Future studies should focus on developing probiotic–herbal feed products based on fermented garlic and assessing the economic and social feasibility of natural phytobiotic applications in intensive aquaculture systems.

**To cite this article:** Permatasari, S. (2025). Utilization of Garlic (*Allium sativum*) as a Functional Supplement to Enhance the Health and Performance of Cultured Fish: A Systematic Literature Review. *Open Global Scientific Journal*, 4(2), 85–94.

This article is under a Creative Commons Attribution-ShareAlike 4.0 International (CC BY-SA 4.0) License. [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/) Copyright ©2025 by author/s

## 1. Introduction

The increasing demand for aquaculture production requires the use of foods that not only meet nutritional needs but also promote fish health. The use of natural food additives has become an important alternative to reduce dependence on synthetic antibiotics, which pose risks of antimicrobial resistance and environmental pollution. One of the most extensively studied natural additives is garlic (*Allium sativum*), a herbal plant rich in bioactive compounds that has long been used in human health applications and is now being developed for aquaculture purposes (Adineh *et al.*, 2020; Valenzuela-Gutiérrez *et al.*, 2021).

Garlic can enhance the immune system to combat diseases and maintain overall health by increasing monocyte activity and phagocytic function (Papu *et al.*, 2014; Erguig *et al.*, 2015). Garlic contains at least 33 sulfur compounds, 17 amino acids, several enzymes, minerals, and vitamins (Kaur & Ansal, 2020). Extracts and isolated compounds of *A. sativum* have been evaluated for various biological activities, including antibacterial, antiviral, antifungal, antiprotozoal, antioxidant, anti-inflammatory, and anticancer properties (Batiha *et al.*, 2020).

The primary active compound in garlic, allicin (allyl 2-propenethiosulfinate or diallyl thiosulfinate), exhibits strong antibacterial, antiviral, and antifungal activity (Adineh *et al.*, 2020). Allicin can activate the non-specific immune system by upregulating cytokine gene expression, thereby enhancing host defense mechanisms when garlic extract is incorporated into fish feed (Fall & Tanekhy, 2015; Erguig *et al.*, 2015). Allicin is recognized as the most potent organosulfur component in garlic, capable of eliminating pathogenic bacteria and parasites, regulating oxidative stress, and improving immune competence. It also enhances gastrointestinal motility and modulates enzyme secretion, which collectively improves digestion and nutrient absorption (Kaur & Ansal, 2020). In addition, other components such as 1-propenyl allyl thiosulfonate, allyl methyl thiosulfonate, (E,Z)-4,5,9-trithiadodeca-1,6,11-triene 9-oxide (ajoene), and  $\gamma$ -L-glutamyl-S-alkyl-L-cysteine function as antioxidants and immunostimulants that enhance non-specific immune responses in fish (Bayan *et al.*, 2014; Valenzuela-Gutiérrez *et al.*, 2021; Bhatwalkar *et al.*, 2021).

Garlic has been found to eliminate major pathogenic bacteria, strengthen immunity, improve health status, enhance growth, and increase flesh quality in freshwater fish such as common carp (*Cyprinus carpio*), tilapia (*Oreochromis niloticus*), rainbow trout (*Oncorhynchus mykiss*), and african catfish (*Clarias gariepinus*) (Hai, 2015; Erguig *et al.*, 2015). Moreover, several studies have demonstrated the immunostimulatory role of garlic in larval and juvenile stages of various fish species, including striped catfish (*Pangasius hypophthalmus*), silver barb (*Barbomyrus gonionotus*), and african catfish (*Clarias gariepinus*), resulting in improved survival and growth performance (Andriani *et al.*, 2017; Hismah *et al.*, 2022; Pangaribuan *et al.*, 2022). With its multifunctional properties, garlic represents a promising functional food additive to enhance growth performance, food utilization, and stress resistance in cultured fish. This review aims to summarize recent scientific findings on the utilization of garlic (*Allium sativum*) as a functional supplement to enhance the health and performance of cultured fish. *This review provides a comprehensive and up-to-date synthesis of recent evidence highlighting garlic (*Allium sativum*) as a multifunctional, antibiotic-free functional feed additive that simultaneously enhances immunity, growth performance, nutrient utilization, and stress resistance across multiple cultured fish species and life stages.*

## 2. Methods


This article was prepared using a systematic literature review approach. The review included: (1) experimental studies investigating the effects of garlic supplementation in fish feed or rearing media; (2) studies involving growth, physiological, immune, or water quality parameters; and (3) articles focusing on aquaculture species such as tilapia (*Oreochromis niloticus*), common carp (*Cyprinus carpio*), african catfish (*Clarias gariepinus*), rohu (*Labeo rohita*), rainbow trout (*Oncorhynchus mykiss*), and groupers (*Epinephelus* spp.).

### Literature selection criteria

Articles were included if they met the following criteria: (i) published within the last ten years (2015–2025); (ii) original research or literature review articles; and (iii) published in peer-reviewed scientific journals indexed in national databases (e.g., Sinta) or international databases (e.g., Scopus, Web of Science, DOAJ).

### Search strategy

Relevant literature was retrieved from Google Scholar, ScienceDirect, SpringerLink, and Researchgate using combination of keywords such as “garlic”, “*Allium sativum*”, “fish growth”, “aquaculture”, and “immunity”. The identification, screening, and eligibility process of systematic literature review approach can be seen in figure 1.



**Figure 1.** The identification, screening, and eligibility process of systematic literature review approach

## Data analysis

Eligible publications were analyzed qualitatively. Findings and conclusions were critically examined and descriptively synthesized to provide an integrated overview of how garlic act as antibacterial agents, immunostimulants, and antioxidants that can improve fish growth and resistance to environmental stress and pathogenic infections. This qualitative-descriptive synthesis forms the scientific basis for developing natural feed additives strategies in intensive aquaculture systems.

## 3. Results and Discussion

### 3.1 Effects of Garlic on Fish Growth

Most studies reported that garlic supplementation in fish diets can enhance growth performance and food conversion efficiency. Based on references found that adding garlic oil (0,5-2%) or garlic extract (1 g/kg) to the diet of *Oreochromis niloticus* improved overall body weight, general health condition, and increased survival rate (Foyal & Javed, 2019; Samson, 2019; Said *et al.*, 2022; Abu-Alya *et al.*, 2022). Other species *Cyprinus carpio*, indicates that adding garlic powder (15 gr/kg), garlic extract (1-5 g/kg) enhanced growth performance, specific growth rate, increased body weight and food efficiency (Manoppo *et al.*, 2016; Karimi Pashaki *et al.*, 2018; Al-Noor *et al.*, 2025). Nyadjeu *et al.*, (2021) and Ukenye *et al.*, (2023) reported that dietary garlic powder (1-2%) of *Clarias gariepinus* improved growth, weight gain, and food utilization.

The growth-promoting mechanism of garlic is closely associated with the stimulation of digestive enzyme secretion and improvement of intestinal morphology. [Huang et al. \(2020\)](#) reported that dietary allicin at levels of 0.005–0.01% enhanced intestinal villi development in *Larimichthys crocea* larvae, thereby improving nutrient absorption. In addition, the antimicrobial properties of garlic help stabilize the intestinal microbiota, which plays a vital role in improving food utilization efficiency ([Bhatwalkar et al., 2021](#); [Delgado et al., 2023](#)). Guroy states that adding garlic powder (1%) enhanced specific growth rate (SGR) and weight gain of rainbow trout (*Oncorhynchus mykiss*). The effects of garlic on fish growth can be seen in table 1.

**Table 1.** The effects of garlic on fish growth based on references

No	Fish Species	Form and Dosage of Garlic	Main Effects on Growth Performance	References
1	<i>Cyprinus carpio</i>	Garlic powder or oil (unspecified dose)	Generally enhanced growth performance	Al-Noor et al., 2025
2	<i>Oreochromis niloticus</i>	Garlic oil (5–15 ml/kg)	Positive effects on growth, hematology, blood biochemistry, hepatosomatic index, and histopathology	Oz et al., 2024
3	<i>Liza ramada</i>	Fermented garlic (1–2%)	Improved growth and antioxidant activity	Basuini et al., 2024
4	<i>Oncorhynchus mykiss</i>	Garlic powder (1%)	Enhanced specific growth rate (SGR) and weight gain	Guroy et al., 2024
5	<i>Oreochromis niloticus</i>	Mixture of lemon, onion, garlic (LOG)	Acted as a growth promoter	Abozaid et al., 2024
6	<i>Clarias gariepinus</i>	Garlic powder (unspecified dose)	Highest weight gain and best performance	Ukenye et al., 2023
7	Multi-spesies	Various garlic forms	Generally improvement in biology performance	Delgado et al., 2023
8	<i>Oreochromis niloticus</i>	Garlic oil (0,5%)	Improved the overall body weight	Abu-Alya et al., 2022
9	<i>Oreochromis niloticus</i>	Garlic oil (0,5-1%)	Improved general health condition	Said et al., 2022
10	<i>Clarias gariepinus</i>	Garlic powder (1-2%)	Better growth, food utilization and body composition	Nyadjeu et al., 2021
11	<i>Labeo rohita</i>	Garlic extract (6.25–25 mg/kg)	Increased SGR and survival rate	Paul et al., 2021
12	<i>Oncorhynchus mykiss</i>	Microencapsulated extract (0–2%)	Improved growth, food conversion ratio (FCR), and body protein content	Adineh et al., 2020
13	<i>Cyprinus carpio</i>	Garlic powder (1–1.75%)	Optimal dose enhanced specific growth rate (SGR) and food conversion	Mohammad, 2020
14	<i>Lates calcalifer</i>	Garlic powder + Ascorbic acid	Improved growth performance, feed utilization, and body composition	Abdelwahab et al., 2020
15	<i>Larimichthys crocea</i>	Allicin (0.005–0.01%)	Enhanced intestinal development and larval growth	Huang et al., 2020
16	Multi-spesies	Various phytobiotics (meta-analysis)	Significant improvement in overall growth performance	Reverter et al., 2020
17	<i>Oreochromis niloticus</i>	Garlic extract (1 g/kg)	Increased survival rate	Foysal & Javed, 2019
18	<i>Oreochromis sp</i>	Garlic powder (1–2%)	Improved survival and growth performance	Samson, 2019
19	<i>Cyprinus carpio</i>	Garlic extract (1–5 g/kg feed)	Increased body weight and food efficiency	Karimi Pashaki et al., 2018
20	<i>Cyprinus carpio</i>	Garlic powder (15 g/kg)	Increased specific body weight gain	Manoppo et al., 2016
21	<i>Oreochromis niloticus</i>	Garlic extract (1%)	Highest weight gain (WG) and specific growth rate (SGR)	Hussein et al., 2016

### 3.2 Effects of Garlic on the Immune System and Disease Resistance

The immunostimulatory effects of garlic have been the primary focus of most related studies. Based on references that adding garlic extract (0,4 g/L or 1-5 g/kg) preventing *Streptococcus iniae* infection, exhibited antiparasitic activity, and adding garlic oil (0,5-1% or 10 ml/kg) improved hematological parameters and antioxidant activity, acted as an immunostimulant (Foysal & Javed, 2019; Reda et al., 2024; Said et al., 2022; Oz et al., 2024). Rezaei et al., (2022) reported increased lysozyme and superoxide dismutase (SOD) activities in crayfish fed diets containing 1% garlic powder. Similarly, Hamed et al., (2020) demonstrated that allicin alleviated liver damage caused by exposure to the pesticide carbofuran in tilapia (*Oreochromis niloticus*) by enhancing the activities of antioxidant enzymes such as superoxide dismutase (SOD) and catalase (CAT).

In pathogen challenge tests, Paul et al., (2021) found that rohu (*Labeo rohita*) fed garlic extract at 6,25-25 mg/kg diet exhibited significantly higher survival rates following infection with *Aeromonas septicemia*. These findings further support the evidence that garlic enhances non-specific immune responses, including increased leukocyte count, phagocytic activity, and serum antibacterial activity in cultured fish (Mahmoud et al., 2019; Abdelwahab et al., 2020; Karimi Pashaki et al., 2018; Rezaei et al., 2022; Delgado et al., 2023; Basuini et al., 2024). The effects of garlic on the immune system and disease resistance can be seen in table 2.

**Table 2.** The effects of garlic on the immune system and disease resistance based on references

No	Fish Species	Form and Dosage of Garlic	Main Effects on the Immune System and Disease Resistance	References
1	<i>Cyprinus carpio</i>	Garlic powder or oil (unspecified dose)	Improved general health and immune condition	Al-Noor et al., 2025
2	<i>Oreochromis niloticus</i>	Garlic oil (10 ml/kg feed)	Improved hematological parameters and antioxidant activity	Oz et al., 2024
3	<i>Oreochromis niloticus</i>	Garlic extract (0.4 g/L water)	Exhibited antiparasitic activity against <i>Dactylogyrus</i> spp.	Reda et al., 2024
4	<i>Oreochromis niloticus</i>	Fermented garlic powder	Exhibited strong antioxidant and antimicrobial effects	Basuini et al., 2024
5	Multi-species	Various garlic forms	Enhanced non-specific immune responses	Delgado et al., 2023
6	<i>Clarias gariepinus</i>	Garlic powder (unspecified dose)	Immunity and disease resistance	Ukenye et al., 2023
7	<i>Postantacus leptodactylus</i>	Garlic powder (1%)	Enhanced immune enzyme activity (lysozyme, phagocytosis)	Rezaei et al., 2022
8	<i>Oreochromis niloticus</i>	Garlic oil (0,5-1%)	Promoted growth and acted as an immunostimulant	Said et al., 2022
9	<i>Oncorhynchus mykiss</i>	Garlic extract mixture (0-2%)	Increased protein ratio and immune response	Oz & Dikel, 2022
10	<i>Epinephelus coioides</i>	Garlic powder (unspecified dose)	High concentrations showed potential toxicity; appropriate dosage is essential	Erazo-Pagador et al., 2022
11	<i>Seriola dorsalis</i>	Garlic extract (1, 4, 16 ml/L water)	Reduced parasitic infection caused by <i>Z. seriolae</i>	Armuelles-Bernal et al., 2022
12	<i>Labeo rohita</i>	Garlic extract (6.25-25 mg/kg feed)	Increased resistance to <i>Aeromonas septicemia</i> infection	Paul et al., 2021
13	<i>Oreochromis niloticus</i>	Garlic extract bionanocomposite	Enhanced oxidative stability and fillet resistance to oxidation	Youssef et al., 2021
14	<i>Oreochromis niloticus</i>	Allicin (0.5-1 g/kg feed)	Increased SOD and CAT enzyme activities; provided hepatic protection against pesticide toxicity	Hamed et al., 2020
15	<i>Oreochromis niloticus</i>	Garlic powder + Ascorbic acid	Improved immune status and antioxidant capacity	Abdelwahab et al., 2020
16	<i>Cyprinus carpio</i>	Garlic extract (unspecified dose)	Decreased <i>Gyrodactylus elegans</i> parasitic infection	Yavuzcan & Bekcan, 2020
17	<i>Oreochromis niloticus</i>	Garlic extract (1 g/kg)	Effective in preventing <i>Streptococcus iniae</i> infection	Foysal & Javed, 2019

18	<i>Oreochromis niloticus</i>	Garlic powder (1,5%)	Significantly reduced lipid peroxidation and exhibited antioxidant effects	Mahmoud <i>et al.</i> , 2019
19	<i>Cyprinus carpio</i>	Garlic extract (1–5 g/kg feed)	Increased WBC count, lysozyme activity, and IgM concentration	Karimi Pashaki <i>et al.</i> , 2018
20	<i>Poecilia reticulata</i>	Garlic powder (1,25 mg/g feed)	Reduced infection intensity on the caudal fin	Kim <i>et al.</i> , 2018
21	<i>Oreochromis niloticus</i>	Garlic extract (1%)	Increased erythrocyte (RBC) and leukocyte (WBC) counts	Hussein <i>et al.</i> , 2016

### 3.3 Effects on Food Efficiency and Fish Physiology

Several studies have reported that garlic supplementation improves feed utilization efficiency by enhancing blood biochemical profiles and liver function. Hussein *et al.*, (2016) state that adding garlic oil (5-15 ml/kg) or garlic extract (1%) increased blood protein levels, digestive enzyme activity, and nutrient utilization efficiency. Abdelwahab *et al.*, (2020) found that the combination of garlic and vitamin C produced synergistic effects on food efficiency and oxidative balance in Nile tilapia (*Oreochromis niloticus*). In addition, Öz *et al.*, (2024) demonstrated that dietary garlic oil (5-15 ml/kg) supplementation improved hematological parameters and total serum protein levels in tilapia.

Positive physiological effects have also been observed in fish reared under extreme environmental conditions. For instance, Wijayanto *et al.*, (2023) reported that garlic supplementation enhanced the growth performance and survival rate of grouper (*Epinephelus* spp.) reared in low-salinity environments. These findings indicate the potential application of garlic in promoting fish adaptation to environmental stress conditions. Also, Basuini *et al.*, (2024) state that adding fermented garlic (1-2%) can increased antioxidant activity and more efficient metabolism of grey mullet (*Liza ramada*). The effects on food efficiency and fish physiology can be seen in table 3.

**Table 3.** The effects on food efficiency and fish physiology based on references

No	Fish Species	Form and Dosage of Garlic	Main Effects on Feed Efficiency and Fish Physiology	References
1	<i>Cyprinus carpio</i>	Garlic powder or oil (unspecified dose)	Increased food efficiency	Al-Noor <i>et al.</i> , 2025
2	<i>Oreochromis niloticus</i>	Garlic oil (5–15 ml/kg)	Increased blood protein levels, digestive enzyme activity, and nutrient utilization efficiency	Öz <i>et al.</i> , 2024
3	<i>Liza ramada</i>	Fermented garlic 1–2%	Increased antioxidant activity and more efficient metabolism	Basuini <i>et al.</i> , 2024
4	<i>Oreochromis niloticus</i>	Various garlic forms	Enhanced metabolic efficiency and digestive enzyme activity	Delgado <i>et al.</i> , 2023
5	<i>Cyprinus carpio</i>	Garlic powder (unspecified dose)	Improved blood and biochemical characteristics	Mohammad MA., 2023
6	<i>Postantacus leptodactylus</i>	Garlic powder (1%)	Improved digestive enzyme activity and nutrient absorption	Rezaei <i>et al.</i> , 2022
7	<i>Labeo rohita</i>	Garlic extract (6.25–25 mg/kg)	Improved food utilization efficiency and reduced oxidative stress	Paul <i>et al.</i> , 2021
8	<i>Oreochromis niloticus</i>	Garlic powder+ Ascorbic acid (unspecified dose)	Enhanced food efficiency, oxidative balance, and liver health	Abdelwahab <i>et al.</i> , 2020
9	<i>Cyprinus carpio</i>	Garlic powder (1–1.75%)	Lower feed conversion ratio (FCR) and improved food efficiency	Mohammad, 2020
10	<i>Oncorhynchus mykiss</i>	Microencapsulated garlic extract (2%)	Reduced FCR, increased body protein, and enhanced digestive enzyme activity	Adineh <i>et al.</i> , 2020
11	<i>Larimichthys crocea</i>	Allicin (0.01%)	Enhanced nutrient absorption through improved intestinal morphology	Huang <i>et al.</i> , 2020
12	Multi-species	Garlic-based phytobiotic (unspecified dose)	Improved average food efficiency by 10–20%	Reverter <i>et al.</i> , 2020

13	<i>Cyprinus carpio</i>	Garlic extract (unspecified dose)	Stress indicators in carp returned to normal after one hour of recovery	Yavuzcan & Bekcan, 2020
14	<i>Cyprinus carpio</i>	Garlic extract (1–5 g/kg feed)	Improved blood biochemical profile and liver function	Karimi Pashaki <i>et al.</i> , 2018
15	<i>Cyprinus carpio</i>	Garlic powder (15 g/kg)	Improved FCR and growth ratio	Manoppo <i>et al.</i> , 2016
16	<i>Oreochromis niloticus</i>	Garlic extract (1%)	Improved immune factors	Hussein <i>et al.</i> , 2016

Overall, dietary garlic supplementation has been shown to consistently improve growth performance, feed utilization, immune function, and stress resistance in various cultured fish species. The effectiveness of garlic is highly dependent on its dosage and form. Across studies, optimal responses were generally observed at 0.5–2% garlic powder, 1–5 g/kg garlic extract, 5–15 mL/kg garlic oil, or 0.005–0.01% allicin, while pathogen resistance was enhanced at lower inclusion levels (6.25–25 mg/kg diet).

These benefits are primarily associated with enhanced digestive efficiency, improved intestinal morphology, modulation of gut microbiota, and stimulation of antioxidant and innate immune responses. When applied within optimal dose ranges, garlic supplementation improved physiological performance without inducing negative effects. Therefore, garlic can be considered an effective natural feed additive for sustainable aquaculture, although further dose–response studies are required to refine species-specific recommendations.

### 3.4. Environmental and Sustainability Aspects

The use of garlic as a natural food additive aligns with the principles of sustainable aquaculture, as it can reduce dependence on antibiotics and chemicals synthetic. [Delgado \*et al.\* \(2023\)](#) and [Reverter \*et al.\* \(2020\)](#), through meta-analysis studies, concluded that the use of herbal plants—including garlic—significantly improves the biological performance of fish without causing toxic effects on the environment. However, [Erazo-Pagador \*et al.\* \(2022\)](#) cautioned that excessively high doses of garlic may be toxic, as indicated by LC<sub>50</sub> values demonstrating potential adverse effects at extreme concentrations. Therefore, determining the optimal dosage is a crucial factor for its practical application in aquaculture systems.

### Research Gaps and Future Directions

Although numerous studies have demonstrated the beneficial effects of garlic supplementation, several scientific gaps remain to be addressed:

1. Lack of standardized dosage and formulation (powder, oil, extract, fermented form) across studies.
2. Limited research on tropical marine fish species compared to freshwater species such as *Oreochromis niloticus* and *Cyprinus carpio*.
3. Few long-term studies evaluating residue effects, chronic toxicity, and interactions with other feed ingredients.
4. Further molecular investigations are required to elucidate the mechanisms of allicin in modulating immune and antioxidant gene expression.

Future studies should focus on developing probiotic–herbal feed products based on fermented garlic and assessing the economic and social feasibility of natural phytobiotic applications in intensive aquaculture systems.

## 4. Conclusion & Recommendations

Garlic (*Allium sativum*) has demonstrated considerable potential as a natural food additive in sustainable aquaculture. Based on the review of 30 scientific articles, dietary garlic supplementation at 0.5–2% garlic powder, 1–5 g/kg garlic extract, 5–15 mL/kg garlic oil, or 0.005–0.01% allicin generally enhances growth performance, food efficiency, and fish resilience to stress and diseases. These positive outcomes are attributed to the antibacterial, antioxidant, and immunostimulant activities of allicin and its bioactive derivatives. Also, the use of garlic as a natural food additive can reduce dependence on antibiotics and chemicals synthetic. Therefore, garlic can be considered an effective natural feed additive for sustainable aquaculture, although further dose–response studies are required to refine species-specific recommendations. Future studies should focus on developing probiotic–herbal feed products based on fermented garlic and assessing the economic and social feasibility of natural phytobiotic applications in intensive aquaculture systems.

## 5. References

Abdelwahab, A. M., El-Bahr, S. M., & Al-Khamees, S. (2020). Influence of Dietary Garlic (*Allium sativum*) and/or Ascorbic Acid on Performance, Feed Utilization, Body Composition and Hemato-Biochemical Parameters of Juvenile Asian Sea Bass (*Lates calcarifer*). *Animals (Basel)*, 10(12), 2396. doi: 10.3390/ani10122396.

Abozaid, H., Elnadi, A. S. M., Mansour, H., Aboelhassan, D. M., Awad, E., El-Nameary, Y. A. A., Omar, H. A. A., Abbas, W. T., & Farag, I. M. (2024). Nutritional Effect of Using a Bioactive Mixture (Lemon, Onion and Garlic) on Growth Performances, Feed Utilization, Immune Status and Gene Expression of Nile tilapia (*Oreochromis niloticus*). *Egyptian Journal of Veterinary Sciences*, 55, 1669-1684. 10.21608/ejvs.2024.266900.1814.

Abu-Alya, I. S., Sheraiba, N. I., Aziz, E. K., Osman, N. A., & Kishta, A. A. (2022). Impact of garlic (*Allium sativum*) and/or ginger (*Zingiber officinale*) oils supplementation on the growth performance, physiological responses, and gene expression of nile tilapia. *Adv. Anim. Vet. Sci.*, 10(11), 2356-2366.

Adineh, H., Harsij, M., Jafaryan, H., & Asadi, M. (2020). The effects of microencapsulated garlic (*Allium sativum*) extract on growth performance, body composition, immune response and antioxidant status of rainbow trout (*Oncorhynchus mykiss*) juveniles. *Journal of Applied Animal Research*, 48(1), 372-378. <https://doi.org/10.1080/09712119.2020.1808473>

Al-Noor, J. M., Najim, S. M., & Al-Tameemi, R. A. (2025). Evaluation of the effects of garlic powder and oil on growth performance and physiological parameters of common carp (*Cyprinus carpio* L.). *Iraqi J. Aquacult.*, 22(1), 187-202.

Andriani, C., Hastuti, S., & Sarjito. (2017). Peran Bawang Putih Dalam Pakan Sebagai Imunostimulan Terhadap Kondisi Kesehatan, Kelulushidupan, dan Pertumbuhan Ikan Tawes (*Puntius javanicus*). *Journal of Aquaculture Management and Technology*, 6 (3), 59-67.

Armuelles-Bernal, C., Cucala-García, L., Barón-Sevilla, B., Díaz-Jiménez, L., Rubio-Godoy, M., & Hernández-Rodríguez, M. (2022). Effect of garlic extract and diallyl sulfide immersion baths in the life stages of *Zeuxapta seriola*, a parasite of California yellowtail (*Seriola dorsalis*). *Aquaculture International*, 30, 1261-1277. <https://doi.org/10.1007/s10499-022-00844-4>

Basuini, M. F. E., Shaban, M. M. E. A., El-Hais, A. M., Soliman, A. A., Abu-Elala, N. M., Teiba, I. I., Alhoshy, M., Sallam, G. R., Shadrack, R. S., Mzengereza, K., & Shehata, A. I. (2024). Exploring the Dual Benefits of Fermented and Non-Fermented Garlic Powder on Growth, Antioxidative Capacity, Immune Responses, and Histology in Gray Mullet (*Liza ramada*). *Fishes*, 9(10), 401. <https://doi.org/10.3390/fishes9100401>

Batiha, G. E., Beshbishi, A. M., Wasef, L. G., Elewa, Y. H. A., Al-Sagan, A. A., El-Hack, M. E. A., Taha, A. E., Elhakim, Y. M. A., & Devkota, H. P. (2020). Chemical Constituents and Pharmacological Activities of Garlic (*Allium sativum* L.): A Review. *Nutrients*, 12(3), 872. doi: 10.3390/nu12030872.

Bayan, L., Koulivand, P. H., & Gorji, A. (2014). Garlic: a review of potential therapeutic effects. *Avicenna J Phytomed*, 4(1), 1.

Bhatwalkar, S. B., Mondal, R., Krishna, S. B. N., Adam, J. K., Govender, P., & Anupam, R. (2021). Antibacterial Properties of Organosulfur Compounds of Garlic (*Allium sativum*). *Front Microbiol*, 12, 613077. doi: 10.3389/fmicb.2021.613077.

Delgado, D. L. C., Caceres, L. L. C., Gomez, S. A. C., & Odio, A. D. (2023). Effect of dietary garlic (*Allium sativum*) on the zootechnical performance and health indicators of aquatic animals: A mini-review. *Veterinary World*, 16(5), 965-976. [www.doi.org/10.14202/vetworld.2023.965-976](http://www.doi.org/10.14202/vetworld.2023.965-976)

Erazo-Pagador, G., Dumaran-Paciente, H. R., & Caloyloy, B. J. (2022). Acute Lethal Toxicity of Dried Garlic (*Allium Sativum*) Powder on Orange-Spotted Grouper (*Epinephelus coioides*) Juveniles under Static Exposure. *Bulletin of the European Association of Fish Pathologists*, 42(1), 28-38. <https://doi.org/10.48045/001c.38092>

Erguig, M., Yahyaoui, A., Fekhaoui, M., & Dakki, M. (2015). The Use of Garlic in Aquaculture. *European Journal of Biotechnology and Bioscience*, 3(8), 28-33.

Fall, J., & Tanekhy. (2015). The Effect of Allicin on Innate Immune Genes of Common Carp (*Cyprinus carpio*

L). *Journal of Applied Biotechnology*, 4(1), 1-12.

Foysal, & Javed, M. D. (2019). Dietary supplementation of garlic (*Allium sativum*) modulates gut microbiota and health status of tilapia (*Oreochromis niloticus*) against *Streptococcus iniae* infection. *Aquaculture Research*, 50(8), 2107–2116. <https://doi.org/10.1111/are.14088>

Güroy, D., Emre, N., Yalim, F. B., Karadal, O., Kaya, D., & Arifoğlu, N. (2024). Interaction of dietary garlic (*Allium sativum*), onion (*Allium cepa*), and probiotic on the growth performance and health status of juvenile rainbow trout (*Oncorhynchus mykiss*). *Aquaculture International*, 32, 4515–4528. <https://doi.org/10.1007/s10499-024-01388-5>

Hai, N. V. (2015). The use of medicinal plants as immuno-stimulants in aquaculture: A review. *Aquaculture*, 446, 88-96.

Hamed, H. S., Ismal, S. M., & Faggio, C. (2020). Journal Pre-proof Effect of allicin on antioxidant defense system, and immune response after carbofuran exposure in Nile tilapia, *Oreochromis niloticus* Effect of allicin on antioxidant defense system, and immune response after carbofuran exposure in Nile tilapia, *Oreochromis niloticus*. *Comparative Biochemistry and Physiology Part C Toxicology & Pharmacology*. 240. 10.1016/j.cbpc.2020.108919

Hismah, N., Amrullah, & Dahlia. (2022). Use of garlic extract (*Allium sativum*) to enhance immunity and growth performance of tilapia seeds (*Oreochromis niloticus*). *Agrokompleks*, 22(2), 18-24.

Huang, W., Yao, C., Liu, Y., Xu, N., Yin, Z., Xu, W., Miao, Y., Mai, K., & Ai, Q. (2020). Dietary Allicin Improved the Survival and Growth of Large Yellow Croaker (*Larimichthys crocea*) Larvae via Promoting Intestinal Development, Alleviating Inflammation and Enhancing Appetite. *Frontiers in Physiology*, 11. <https://doi.org/10.3389/fphys.2020.587674>

Hussein, M. S., El-Zaiat, A. M., & El-Saiad, S. M. (2016). Effects of garlic and onion oil extracts as a natural growth promoters on growth performance, nutrient utilization, whole body composition and hematological parameters of Nile Tilapia (*Oreochromis niloticus*) fingerlings. *Journal of Egyptian Academic Society for Environmental Development. D*, 17(1), 141-15.

Karimi Pashaki, A., Zorriehzahra, S. J., Ghasemi, M., Sharif Rohani, & M., Hosseini, S. (2018). Effects of dietary Garlic extract on some blood, immunity and growth parameters of Common Carp fingerlings (*Cyprinus carpio*). *Sustainable Aquaculture Health Management Journal*, 4 (2), 28-39.

Kaur, H & Ansal, M. (2020). Efficacy of garlic powder as a growth pro moting feed additive for higher growth and improved flesh quality in an Indian Major Carp, *Labeo rohita* (Ham.) fingerlings. *Journal of Entomology and Zoology Studies*, 8, 25-29.

Kim, J. H., Fridman, S., Borochov-Neori, H., Sinai, T., & Zilberg, D. (2018). Evaluating the use of garlic (*Allium sativum*) for the remedy of *Cryptocaryon irritans* in guppies (*Poecilia reticulata*). *Aquaculture Research*. <https://doi.org/10.1111/are.13904>

Mahmoud, R., Aziza, A., Marghani, B., & Eltaysh, R. (2019). Influence of ginger and garlic supplementation on growth performance, whole body composition and oxidative stress in the muscles of nile tilapia (*O. niloticus*). *Adv. Anim. Vet. Sci.*, 7(5), 397-404.

Manoppo, H., Kolopita, M. E. F., & Malatunduh, R. (2016). Growth promoter effect of garlic (*Allium sativum*) on carp (*Cyprinus carpio* L.). *IJPRIF*, 9(4), 283-288.

Mohammad, M. A. (2020). Influence of Different Levels of Garlic *Allium Sativum* Powder Additions on Growth Promoter and Biochemical Composition of Common Carp *Cyprinus Carpio* L. *Plant Archives*, 20(2), 7321–7326.

Mohammad, M. A. (2023). Effect Of Adding Garlic *Allium Sativum* Powder in Diet on Hematological, Biochemical And Histopathological Criteria Of Common Carp *Cyprinus Carpio* L. *Iraqi Journal Of Agricultural Sciences*, 54(4), 1040-1049. <https://doi.org/10.36103/ijas.v54i4.1793>

Nyadjeu, P., Yemdjie, D., Ndjuissi, N., Nguenang, G., Dedou, N., & Tabi-Tomedi M. (2021). Effect of *Zingiber officinale* and *Allium sativum* Powders as Natural Feed Additives Promoting Growth, Feed Utilization and Whole-Body Composition in *Clarias gariepinus* Fry. *Food and Nutrition Sciences*, 12, 526-543. doi: 10.4236/fns.2021.126040

Öz, M., & Dikel, S. (2022). Effect of garlic (*Allium sativum*)-supplemented diet on growth performance, body composition and fatty acid profile of rainbow trout (*Oncorhynchus mykiss*). *Cellular and Molecular Biology*, 68(2), 217–225. DOI: 10.14715/cmb/2022.68.1.26

Oz, M., Inanan, B. E., Ustuner, E., Karagoz, B., & Dikel S. (2024). Effects of dietary garlic (*Allium sativum*) oil

on growth performance, haemato-biochemical and histopathology of cypermethrin-intoxicated Nile tilapia (*Oreochromis niloticus*). *Vet Med Sci*, 10(3), 1449. doi: 10.1002/vms3.1449.

Pangaribuan, J., Yonarta, D., & Rarassari, M. A. (2022). Impact of Additional Onion and Black Garlic on Fish Feed to The Life Sustainability of Lele. *Jurnal Kelautan Perikanan Terapan*, 5(1): 83-88.

Papu, S., Jalvrl, S., Sweta, S., & Singh, B. R. (2014). Medicinal values of garlic (*Allium sativum* L.) in human life: An overview. *Greener Journal of Agricultural Sciences*, 4(6), 265-280.

Paul, S., Rahman, M., Salam, M., Surovy, M., & Islam, T. (2021). Dietary Inclusion of Garlic (*Allium Sativum*) Extract Enhances Growth and Resistance of Rohu (*Labeo Rohita*) Against Motile *Aeromonas septicaemia*. *Annals of Bangladesh Agriculture*, 25(1), 11–22. <https://doi.org/10.3329/aba.v25i1.58151>.

Reda, R., Khalil, A. A., Elhady, M., Tayel, S. I., & Ramadan, E. A. (2024). Anti-parasitic activity of garlic (*Allium sativum*) and onion (*Allium cepa*) extracts against *Dactylogyrus* spp. (Monogenean) in Nile tilapia (*Oreochromis niloticus*): Hematology, immune response, histopathological investigation, and inflammatory cytokine genes of gills. *BMC Vet Res.*, 20(1), 334. doi: 10.1186/s12917-024-04187-5.

Reverter, M., Bontemps, N., Lecchini, D., Banaigs, B., & Sasal, P. (2020). Moving towards more sustainable aquaculture practices: a meta-analysis on the potential of plant-enriched diets to improve fish growth, immunity and disease resistance. *Reviews in Aquaculture*, 1-19. DOI:10.1111/raq.12485

Rezaei, O., Mehrgan, M. S., & Paknejad, H. (2022). Dietary garlic (*Allium sativum*) powder improved zootechnical performance, digestive enzymes activity, and innate immunity in narrow-clawed crayfish (*Postantacus leptodactylus*). *Aquaculture Reports*. 24. 101129. 10.1016/j.aqrep.2022.101129.

Said, I., Sheraiba, N. I., Aziz, E., Osman, N., & Kishta, A. (2022). Impact of Garlic (*Allium sativum*) and/or Ginger (*Zingiber officinale*) oils supplementation on the growth performance, physiological responses, and gene expression of Nile tilapia. *Advances in Animal and Veterinary Sciences*, 10. 10.17582/journal.aavs/2022/10.11.2356.2366.

Samson, J. S. (2019). Effect of garlic (*Allium sativum*) supplemented diets on growth, feed utilization and survival of red tilapia (*Oreochromis* sp.). *International Journal of Agricultural Technology*, 15(4), 637-644.

Ukenye, E. A., Megbowon, I., Oguntade, O. R., Ayo-Olalusi, C. I., Mojekwu, T., Edah, B., Sokenu, B.A., Joseph, J. B., Adeyemi, Y. B., Jubrin, H., Chidume, I., Eze, N. C., Adeleke, R. O., & Igwe, C. (2023). Growth Performance and Immunity Analysis of *Clarias gariepinus* Juveniles Fed with Moringa leaf, Ginger and Garlic Powder. *Nigeria Agricultural Journal*, 54(1), 275–280. <https://doi.org/10.4314/naj.v54i1.252342>.

Valenzuela-Gutierrez, R., Lago-Leston, A., Vargas-Albores, F., Cicala, F., Martinez-Porchas, M. (2021). Exploring the garlic (*Allium sativum*) properties for fish aquaculture. *Fish Physiol Biochem*, 47(4), 1179-1198. doi: 10.1007/s10695-021-00952-7.

Wijayanto, D., Kurohman, F., Nugroho, R. A., & Nursanto, D. B. (2023). The effect of garlic supplementation on the growth, survival and RCR in cultured TGGG hybrid grouper ( $\text{♀}$  tiger grouper  $\times$   $\text{♂}$  giant grouper). *AACL Bioflux*, 16(5), 2554-2561.

Yavuzcan, H., & Bekcan, S. (2020). Control of ectoparasitosis in carp (*Cyprinus carpio*) induced by *Gyrodactylus elegans* (Monogenea) with garlic (*Allium sativum*) and onion (*Allium cepa*) extracts. *Ecocycles*, 6(1), 10–17. <https://doi.org/10.19040/ecocycles.v6i1.157>.

Youssef, R., Abd El-Hakim, N., & Abouelwafa, R. (2021). Garlic-based bionanocomposite film for preserving the quality of tilapia fillets during storage. *RSC Advances*, 11(3), 1545–1554.