



## Efficacy and Safety of Cannabidiol for Anxiety: A Bibliometric Analysis and Systematic Review

Putu Ayu Putri Fajaryani<sup>1</sup>, I Made Agus Gelgel Wirasuta<sup>1,3</sup>, Pande Made Nova Armita Sari<sup>1</sup>, Dyah Kanya Wati<sup>2</sup>, Putu Indah Budi Apsari<sup>4</sup>, Made Ary Sarasmita<sup>1\*</sup>

<sup>1</sup>Department of Pharmacy, Faculty of Mathematics and Natural Sciences, Udayana University, Badung, Bali, Indonesia

<sup>2</sup>Pediatric Consultant, Critical Care Medicine Udayana University, Sanglah Hospital, Denpasar, Bali, Indonesia

<sup>3</sup>Forensic Sciences Laboratory, Institute of Forensic Sciences and Criminology, Udayana University, Badung, Bali, Indonesia

<sup>4</sup>Department of Parasitology, Faculty of Medicine and Health Sciences, Warmadewa University, Denpasar Bali, Indonesia

\*Corresponding Author: E-mail: [arysarasmita@unud.ac.id](mailto:arysarasmita@unud.ac.id)

### ARTICLE INFO

**Manuscript Received:** 06 Nov, 2024

**Revised:** 13 Feb, 2025

**Accepted:** 23 Feb, 2025

**Date of publication:** 02 Oct, 2025

**Volume:** 5

**Issue:** 3

**DOI:** [10.56338/jph.v5i3.6807](https://doi.org/10.56338/jph.v5i3.6807)

### KEYWORDS

Anxiety;  
Anxiolytic;  
Cannabidiol;  
Pharmacology

### ABSTRACT

**Introduction:** A non-psychoactive substance present in the cannabis plant, cannabidiol (CBD), seems to have potential as an anxiolytic agent. This study systematically reviews the efficacy, safety, dosage, mechanisms, and adverse effects of CBD in treating anxiety.

**Methods:** The PRISMA guideline was used to conduct the review.

**Results:** A total of 64 articles were included in the study. It showed that CBD works at the endocannabinoid receptors CB1 and 5-HT1A, which regulate mood and reduce anxiety. In animal models, a range dosage of 5 – 10 mg/kg showed a significant anxiolytic effect in behavioral tests, while in humans, 300 mg/day was proven to be effective in treating various types of anxiety disorders. However, individual responses to CBD showed varying results, and demographic characteristics affect the efficacy of animal and human studies. CBD was also considered safer to be used in the short term compared to diazepam, although the long-term effect remains lacking evidence.

**Conclusion:** This study concludes that CBD has potential as an alternative choice for anxiety. Further research is needed in larger populations with rigorous study design and longer study durations to evaluate its effectiveness and ensure its safety.

**Recommendations for Future Research:** It is hoped that future research can examine the effects of fly resistance and killing on non-target organisms and control environmental conditions.

**Publisher:** Pusat Pengembangan Teknologi Informasi dan Jurnal Universitas Muhammadiyah Palu



OR “Anxiolytic”) and there were no initial year limits. The reliance on specific databases may limit the inclusion of relevant studies, affecting the completeness of this review.

### Inclusion Criteria

The chosen study articles concentrated on the mechanisms of CBD in treating anxiety, appropriate dosage form, dosage or strength, and adverse effects. The main source of this research was in vivo studies written in English. We included animal and human studies written in English. The chosen studies assessed a minimum of two significant points, including cannabidiol and anxiety.

### Exclusion Criteria

Conference papers, thesis dissertations, review articles, papers published in conference proceedings, manuscripts without abstracts, gray literature or unpublished studies and those that did not fit the inclusion criteria were all disqualified to proceed with our review. Articles discussing the association of CBD with the diseases not covered in this review were also excluded from the analysis. These exclusion criteria may introduce selection bias, potentially limiting the generalizability of the findings

### Data extraction and management

#### Bibliometric Analysis

We used VOSviewer (v1.6.19, CWTS, Leiden University) to visualize term relationships in scientific literature, aiding conceptual understanding and study selection. By analyzing co-occurring keywords from titles and abstracts, with a minimum threshold of two occurrences, we generated impactful network, overlay, and density visualizations to highlight key interconnections.

### Systematic Review

We utilized Zotero (AGPL, USA) to collect and manage relevant articles, ensuring alignment with inclusion criteria for analysis. Studies were categorized into two groups: (1) animal studies and (2) human studies. Key data extracted from animal studies included animal type, intervention, dosage, study design, and outcomes (Table 2 and Table 3).

## RESULTS




Figure 2. PRISMA diagram [22]

Relevant articles for treating anxiety using cannabidiol were identified through a systematic search according to the PRISMA guidelines.




Figure 3. Network Visualization

The network visualization of the studies is shown in Figure 3. Based on the color of the nodes, the VOS viewer groups the keywords into seven distinct clusters.




Figure 4. Overlay Visualization

Figure 4 depicts research trends on CBD and anxiety (2012–2022) using an overlay visualization. Keywords are color-coded by publication year in the VOS viewer. Notably, "anxiety" showed peak activity in 2018.




Figure 5. Density Visualization

Figure 5 highlights keyword frequency and intensity related to CBD and anxiety using a heat map. Bright yellow indicates high density (frequent co-occurrence), while dark blue/purple shows lower density. Terms like "cannabidiol" and "anxiety" appeared in 100% of articles, while others, such as "open field test" (12.4%) and "placebo," had lower density, reflecting limited research or emerging topics.

Table 2 shows that CBD has significant anxiolytic effects, particularly in rats, as evidenced by behavioral tests like the elevated plus maze and open field test. Effective doses range from 0.3 to 30 mg/kg, with optimal effects at 5 – 10 mg/kg. Some studies report a biphasic response, where very low or high doses are less effective, suggesting dose-dependent effects influenced by species and experimental conditions.

**Table 2.** Overview of the search for Cannabidiol (CBD) in Animal Models.

Study	Animal	Test	Intervention	Doses and Administration	Results
Uribe - Mariño et al., 2012 (23)	Mice (Male Swiss)	MP	CBD	0.3, 3 and 30 mg/kg; iv	CBD treatment reduced defensive immobility, behavioral index, explosive fleeing, and total fleeing duration.
Guimarães et al., 1990 (24)	Rat (Male Wistar)	EPM	CBD vs Diazepam vs Vehicle	CBD: 2.5, 5.0, 10 and 20 mg/kg Diazepam: 2 mg/kg; i.p	CBD (5 mg/kg) increased open-arm entries but was weaker than diazepam.
Elbatsh et al., 2012 (25)	Rat (Male Lister-hooded)	CER	CBD vs Vehicle	For 14 days, 10 mg/kg. (acute)	Repeated CBD administration significantly increased freezing behavior duration.
Guimarães et al., 1994 (7)	Rat (Male Wistar)	EPM	CBD vs HU-219 vs HU-252 vs HU-261 vs Diazepam	CBD (5 mg/kg), HU-219 (0.03-3.0 mg/kg), HU252 (0.2-5 mg/kg), HU-261 (0.3-10 mg/kg) and Diazepam (2.5 mg/kg); i.p	CBD and HU-219 (0.03–1 mg/kg) increased open-arm entries but were less potent than diazepam.
Moreira, 2006 (26)	Rat (Male Wistar)	VC	CBD vs Diazepam vs CBD + Flumazenil vs Diazepam + Flumazenil	CBD (2.5, 5 or 10 mg/kg), Flumazenil (10 mg/kg), Diazepam (3.0 mg/kg); i.p	Flumazenil did not block CBD's anxiolytic effects; CBD (10 mg/kg) increased both penalized and total licks.
Zieba et al., 2019 (9)	Mice (Male Fmr1 knockout and WT)	OF	CBD vs vehicle	5 and 20 mg/kg; i.p	CBD (5 mg/kg or 10 mg/kg) showed no significant effect on anxiety parameters compared to controls.
		EPM	CBD vs vehicle	5 and 20 mg/kg; i.p	CBD induced anxiolytic effects in both mice, with no differences between genotypes.
Silva-Cardoso et al., 2021 (27)	Rat (Male Wistar)	OF	CBD vs Vehicle	0.3, 3, 10, 30 mg/kg; i.p	CBD (3 mg/kg) effectively reduced anxiety caused by chronic pain.
Almeida et al., 2013 (28)	Rat (Male Wistar)	SI	CBD	1, 5, 15, 30, 60 mg/kg; i.p	CBD (1 mg/kg) increased passive social interactions.
Shu et al., 2024 (29)	Mice (Male ICR)	SIH	CBD vs Diazepam vs CBD with WAY100635	WAY100635 (0,1, 0,3, 1 mg/kg), diazepam (3 mg/kg), CBD (1, 3, 10 mg/kg); i.p	CBD reduces SIH by 55% and 51% at 3 and 10 mg/kg but diazepam at 3 mg/kg shows a stronger 86% reduction.

Study	Animal	Test	Intervention	Doses and Administration	Results
Kaplan et al., 2021 (8)	Mice (Male and Female C57BL/6 J)	EPM	CBD	5, 10 mg/kg; i.p	CBD at 5 mg/kg increased time spent in open vs. closed arms.
Wanner et al., 2021 (30)	Mice (Male and Female Agouti viable yellow (Avy)).	MB	CBD	20 mg/kg/day; oral	F1 female mice treated with CBD during development buried twice as many marbles as controls.
Chaves et al., 2021 (31)	Rat (Male Wistar)	EPM	Experimental design I: vehicle vs CBD vs CBD with WAY100635	WAY 100635: 0.1 mg/kg CBD: 30 mg/kg; i.p	WAY100635 partially reduced CBD's anxiolytic effect.
			Experimental design II: Vehicle vs CBD; vs CBD with AM251; vs CBD with AM630	CBD: 30 mg/kg AM251: 1 mg/kg AM630: 1 mg/kg; i.p	CBD reduces anxiety by increasing time and frequency in open arms, but AM251 diminishes this effect.
Chaves et al., 2023 (32)	Rat (Male Wistar)	EPM	CBD vs Vehicle	30, 60 mg/kg; i.p	CBD increasing open-arm entries, duration, and overall exploratory activity ( $p < 0.05$ )
De Grogerio et al., 2019 (5)	Rat (Male Wistar)	NSF and EPM	CBD vs CBD with WAY100635, CBD with Capsazepine	CBD: 5 mg/kg, WAY100635: 2 mg/kg, Capsazepine: 10 mg/kg; s.c	CBD increased open-arm duration, but its anxiolytic effects were inhibited by WAY100635.
Alegre-Zurano et al., 2021 (33)	Mice (Male CD1)	EPM	Vehicle vs CBD vs CBD with MDPV	CBD: 20 mg/kg, MDPV: 3 – 4 mg/kg; i.p	Mice given MDPV and CBD spent extra time in the open arm
Austriach-Olivares et al., 2022 (34)	Mice (Male WT)	LDB	CBD	10, 20, 30 mg/kg; i.p	CBD at 10 and 20 mg/kg increased time spent in the light box.
		EPM	CBD	10, 20, 30 mg/kg; i.p	CBD (10, 20 mg/kg) increased open-arm percentage, peaking at 20 mg/kg.
		NSFT	CBD	10, 20, 30 mg/kg; i.p	CBD at 10 mg/kg improved motivation and feeding, while 20 mg/kg showed strong anxiolytic effects.
Hsiao et al., 2012 (35)	Rat (Male Wistar)	EPM OF	CBD	0,5 and 1 mg/kg; inject to CeA	CBD (1 mg/kg) increased exploration in the OF and time spent in the open arm.
Chesworth et al., 2022 (36)	Mice (Female WT) and APPxPS 1)	EPM	CBD	20 mg/kg; oral	CBD showed no effect in both mouse groups.
Rock et al., 2017 (37)	Rat (Male)	LDB emergen	CBD vs CBDA	5 mg/kg; i.p	CBD (5 mg/kg) reversed FS stress-induced anxiety and

Study	Animal	Test	Intervention	Doses and Administration	Results
	Sprague-Dawley)	ce test with foot shock stress	CBD, THC	CBD: 1 µg/mL, THC: 1 mg/kg; i.p	increased time spent in the bright room. THC induced anxiety and foot shock stress worsens anxiety.
Melkumnyan et al., 2024 (38)	Mice (Male and Female C57Bl/6)	OF	CBD vs CBD: THC (3:1)	CBD :10 mg/kg, CBD : THC (3:1): 7,5:2,5 mg/kg	At 24-hour alcohol withdrawal, single CBD reduced anxiety.
Marinho et al., 2015 (39)	Rat (Male wistar)	EPM  Restrain stress and EPM	CBD vs CBD with WAY100635	CBD: 15, 30, and 60 nmol, WAY100635: 0,37 nmol; microinjection to intra-infralimbic	CBD (15 and 30) nmol increased exploration of EPM open arms. In stress-restricted rats, CBD treatment did not alter open arm exploration in the EPM compared to untreated groups.
Fogaça et al., 2014 (6)	Rat (Male Wistar)	EPM  Restrain stress and EPM	CBD vs CBD with WAY100635  CBD vs CBD with WAY100635 vs Metyrapone	CBD: 15, 30, and 60 nmol, WAY100635: 0,37 nmol; microinjection to intra-prelimbic  CBD: 15, 30, and 60 nmol, WAY100635: 0,37 nmol; microinjection to intra-prelimbic. Metyrapone: 75 mg/kg;i.p	CBD (30nmol) reducing open-arm exploration in the EPM. CBD's effect on EPM shifted from anxiogenic to anxiolytic-like in rats.
Granjeiro et al., 2011 (40)	Rat (Male Wistar)	EPM	CBD	30 nmol	CBD exhibited increased open-arm entries.
Campos and Guimarães, 2009 (41)	Rat (Male Wistar)	EPM	CBD vs CBD with Capsazepine vs WIN 55,212-2 vs WIN 55,212-2 with Capsazepine	CBD: 30 and 60 nmol, WIN 55,212-2: 3, 10, 30 nmol, Capsazepine: 10 nmol; intra-dIPAG	CBD (30 nmol) and WIN 55,212-2 (3–10 nmol) increased open-arm entries, while higher doses reduced the effect. Capsazepine blocked anxiety from high doses.
Bitencourt et al., 2008 (3)	Rat (Male Wistar)	EPM	CBD vs AM404 vs Diazepam	AM404 (1.0 µg/µl), CBD (2.0 µg/µl), Diazepam (2.85 µg/µl); i.c.v	AM404 and CBD increased open-arm entries in the conditioned group, while diazepam increased entries in both groups.
Salviato et al., 2021 (42)	Rat (Female Wistar)  Rat (Male wistar)	EPM  EPM	CBD  THC: CBD	1 mg/kg and 3 mg/kg; i.p  THC (1 mg/kg): CBD (1 – 3 mg/kg), THC (0,075 mg/kg): CBD (1 mg/kg); i.p	CBD (3 mg/kg) showed increase in %OAT and %OAE compared to control. THC's anxiogenic effects were reduced with CBD (1-3 mg/kg). Combining THC (0.075 mg/kg) and CBD (1 mg/kg) enhanced THC's anxiolytic effects.
Breurer et al., 2016 (43)	Mice (Male swiss)	EPM	CBD  HUF-101	30 mg/kg; i.p  1, 3, and 10 mg/kg. i.p	Time spent in the open arm (%OAT) and entries into the open arm (%OAE) were increased. The number of entries into the open arm (%OAE) and the

Study	Animal	Test	Intervention	Doses and Administration	Results
					duration in the open arm (%OAT) were increased by 3 mg/kg.
			HUF-102	1, 10 and 60 mg/kg; i.p	HUF-102 showed no increase in open-arm time or entries.
			HUF-103	1, 3 and 10 mg/kg; i.p	Significant increases in open-arm time (%OAT) at 3 and 10 mg/kg.
Campos and Guimarães, 2008 (4)	Rat (Male Wistar)	EPM	CBD	15, 30, and 60 nmol	Rats (30 nmol) spent more time in the open arm, showing a bell-shaped dose-response curve with 30 nmol as the optimal dose.
			CBD with AM251	CBD: 30 nmol, AM251: 100 pmol	Rats given 30 nmol CBD with AM251 showed similar open-arm exploration as with CBD alone.
			CBD with WAY100635	CBD: 30 nmol, WAY100635: 0,37 nmol.	CBD (30 nmol) with WAY100635 didn't increase open-arm exploration.
		VC	CBD	15, 30, dan 60 nmol	Rats given 30 nmol CBD drank more despite electric shocks.
Masataka, 2024 (44)	Cat (Male Domesti)	SB	CBD vs placebo	4 mg/kg	Cats given CBD spent more time near their owner without contact.
Schleicher et al., 2019 (45)	Mice (Male and Female C57BL/6 J)	LDB	CBD (3 months/former group) vs CBD (6 months/current groups)	20 mg/kg	CBD-treated mice showed no increased time in the lighted area, unlike vehicle-treated mice.
		EPM	CBD (3 months/former group) vs CBD (6 months/current group)	20 mg/kg	Former group showed no change, while the current group reduced time in the open arm.
Nazario et al., 2015 (46)	Zebrafish (Male and Female WT)	Time in the top area of the aquarium	CBD	0.1, 0.5, 5.0, and 10 mg/kg	CBD showed a U-shaped dose-response, with 0.5 mg/kg significantly increasing surface time.
Fabris et al., 2022 (47)	Rat (Male and female Wistar)	EPM	CBD	0.3, 3, 30 mg/kg; i.p	In males, CBD (3 mg/kg) increased open-arm entries; in females, effects appeared at 0.3 mg/kg in late diestrus.
Pérez-Valenzuela et al., 2023 (48)	Rat (Male and female Sprague Dawley)	LDB	CBD vs THC: CBD	30 mg/kg, 0.3 mg/kg: 30 mg/kg; oral	CBD alone had little effect; combined treatment reduced anxiety in stressed males.
		EPM	CBD vs THC: CBD	30 mg/kg, 0.3 mg/kg: 30 mg/kg; oral	In males, the combination significantly increased open-arm duration.

<b>Study</b>	<b>Animal</b>	<b>Test</b>	<b>Intervention</b>	<b>Doses and Administration</b>	<b>Results</b>	
Huffstetler et al., 2023 (49)	Mice (Male and Female WT C57BL/6 J) and (Kv1.3 <sup>-/-</sup> )	MB	CBD	10 mg/kg and 20 mg/kg; i.p	Male WT mice given CBD (10 mg/kg) buried significantly fewer marbles than controls.	
		EPM	CBD	10 mg/kg and 20 mg/kg; i.p	WT mice showed no change, but Kv1.3 <sup>-/-</sup> females (10 mg/kg) spent more time in closed arms.	
		LDB	CBD	10 mg/kg and 20 mg/kg; i.p	CBD eased anxiety in WT mice but increased it in female Kv1.3 <sup>-/-</sup> mice at 20 mg/kg.	
Campos et al., 2013 (50)	Mice (Male WT C57BL/6 J and GFAP-TK transgenic)	EPM	CBD vs CBD with AM251	CBD: 30 mg/kg (14 days), AM251: 1 mg/kg; i.p	CBD increased open-arm duration, showing anxiolytic effects in stressed WT mice.	
		NSF	CBD vs CBD with AM251	CBD: 30 mg/kg (14 days), AM251: 1 mg/kg; i.p	CBD reduced eating latency in stressed WT mice.	
Fogaça et al., 2018 (51)	Mice (Male C57BL/6)	EPM	CBD	30 mg/kg; i.p	Repeated CBD intake boosted open-arm exploration in stressed mice.	
			CBD with AM251	30 mg/kg, 0.3 mg/kg; i.p	AM251 blocked the CBD's anxiolytic effect.	
			CBD with AM630	30 mg/kg, 0.3 mg/kg; i.p	AM630 blocked the CBD's anxiolytic effect.	
			CBD with WAY100635	30 mg/kg, (0.05 mg/kg and 0.1 mg/kg); i.p	WAY100635 (0.05 mg/kg) did not block CBD's anxiolytic effect.	
			NSF	CBD	30 mg/kg; i.p	CBD decreased the latency to feed in stressed mice.
			CBD with AM251	30 mg/kg 0.3 mg/kg; i.p	AM251 blocked CBD's anxiolytic effect..	
			CBD with AM630	30 mg/kg, 0.3 mg/kg; i.p	AM630 blocks CBD's anxiolytic effect.	
Gomes et al., 2011 (52)	Rat (Male Wistar)	EPM	CBD	15, 30, or 60 nmol; injection into BNST	CBD (60 nmol) significantly increased open-arm entries and duration.	
			CBD with WAY100635	CBD: 15, 30, or 60 nmol, WAY100635: 0.37 nmol; injection into BNST	WAY100635 (0.37 nmol) blocked CBD's anxiolytic effects.	
			VC	CBD	15, 30, or 60 nmol; injection into BNST	CBD (30–60 nmol) increased punished licks, resembling anxiolytic effects.
			CBD with WAY100635	CBD: 15, 30, or 60 nmol, WAY100635: 0.37 nmol; injection into BNST	WAY100635 blocked CBD's anxiolytic effects.	
Shallcross et al., 2019 (53)	Rat (Male Sprague-Dawley)	EPM	CBD	5 mg/kg; i.p	Rats treated with CBD were less frequent in the open arms	
			CPPDB	30 mg/kg; s.c	CPPDB did not affect open-arm duration.	

Study	Animal	Test	Intervention	Doses and Administration	Results
		LDB	CBD	5 mg/kg; i.p	CBD reduced dark box time and sped up light entry.
			CPPDB	30 mg/kg;s.c	CPPDB had no effect on dark box time or light box latency.
Morris et al., 2020 (54)	Dog (Male and Female)	Noise-induced fear response model	CBD	1.4 mg/kg body	CBD doesn't significantly decrease anxiety.
			CBD with Trazodone	CBD: 1.4 mg/kg, Trazodone: 100 mg for dogs 10-20 kg, 200 mg for dogs 20.1-40 kg	No added benefit over trazodone alone; may reduce its cortisol-lowering effect.
Stern et al., 2015 (55)	Rat (Male Wistar)	EPM	CBD	1, 3 mg/kg; i.p	CBD showed no significant effect on open-arm duration or entries versus controls.
			THC with CBD	THC (0.1 mg/kg) and CBD (1.0 mg/kg); i.p	THC (0.1 mg/kg) and CBD (1.0 mg/kg) reduced freezing and boosted open-arm exploration.
		OF	CBD	1, 3 mg/kg; i.p	CBD (1.0 or 3.0 mg/kg) did not affect center duration in the OF test versus controls.
			THC with CBD	THC (0.1 mg/kg) and CBD (1.0 mg/kg); i.p	The combination treatment increased center duration in the OF test.

BNST: Bed Nucleus of the Stria Terminalis; CBD: Cannabidiol; CBDA: Cannabidiol Acid; CeA: Central Amygdala; CER: Conditioned Emotional Response; CPPDB: 3-Cyano-N-(1,3-diphenyl-1H-pyrazol-5-yl)benzamide; dIPAG: Dorsolateral Periaqueductal Gray; EPM: Elevated Plus Maze; i.c.v: intracerebroventricular; i.p: Intraperitoneal; LDB: Light Dark Box; MB: Marble Burying; MDPV: Methylendioxy pyrovalerone; MP: Mock Predatory; NSF: Novelty-Suppressed Feeding; OF: Open Field; s.c: Subcutaneous; SB: Secure Base; SI: Social Interaction; SIH: Stress-Induced Hyperthermia; THC: Tetrahydrocannabinol; VC: Vogel Conflict; VH: Ventral Hippocampus; vs: versus; WT: Wild Type.

Table 3 presents that CBD reduces anxiety across diverse populations, including individuals with social anxiety disorder and PTSD at doses ranging from 150 to 800 mg/day. It has been found that combining CBD with THC or antidepressants could balance the anxiogenic effects of THC and improved therapeutic outcomes.

**Table 3.** Overview of the search for Cannabidiol (CBD)

Study	N Participant (condition)	Age (years)	Drug	Doses and administration	Results
Stanley et al., 2023 (56)	32 (Test Anxiety Induction)	18 – 25	CBD vs placebo	150, 300, 600 mg; oral	Patients on 600 mg CBD showed more physical anxiety signs than placebo and lower-dose groups.
Zuardi et al., 1982 (2)	8 (healthy)	20 - 38	CBD	1 mg/kg; oral	CBD had mild anxiolytic effects, weaker than diazepam.
			CBD: THC	1 mg/kg: 0.5 mg/kg; oral	The THC-CBD combination consistently reduced psychological anxiety.

Study	N Participant (condition)	Age (years)	Drug	Doses and administration	Results
Martin et al., 2021 (57)	538 (anxiety)	≥ 18	Medical cannabis product (CBD dominant 82 %)	61 mg; oral	CBD consistently reduced anxiety in both acute and long-term doses without psychoactive effects.
			Medical cannabis product (balanced THC: CBD 7%)	-	CBD can reduce the anxiety-increasing THC's effect.
Masataka, 2019 (58)	37 (SAD)	18-19	CBD vs placebo	300 mg/day for 4 weeks; oral	Significant decrease in social anxiety scores (FNE & LSAS) compared to placebo
Hurd et al., 2019 (13)	42 (heroin use disorder)	21-65	CBD vs placebo	400 mg; oral once daily for 3 consecutive days	Moderate anxiety and craving reductions lasted up to 7 days post-dose.
				800 mg; oral once daily for 3 consecutive days	CBD sharply reduced anxiety and craving, lasting 7 days, beating 400 mg and placebo.
Berger et al., 2022 (59)	31 (anxiety)	12 - 25	CBD	Starting at 200 mg/day, titrated to max 800 mg/day over 12 weeks; oral	Anxiety dropped 42.6% in 12 weeks, with 40% seeing a 50% reduction.
			Antidepressants with CBD		CBD with antidepressants reduced anxiety more than CBD alone.
Bergamaschi et al., 2011 (10)	24 (SAD) and 12 (healthy)	23	CBD vs Placebo	600 mg; oral	CBD significantly reduced anxiety during public speaking.
Laczkovics et al., 2021 [59]	1 (SAD)	16.9	CBD	Cannabidiol (CBD), 100-600 mg/day (over 8 weeks); Sertraline (100 mg/day, discontinued after 3 weeks of CBD); oral	CBD reduced anxiety and depression, lowering depressive symptoms from moderate to minimal in 8 weeks.
Hutten et al., 2022 (60)	26 (cannabis users)	23.1	THC-dominant cannabis	13.75 mg THC, vaporized	Significant increase in state anxiety compared to placebo.
			CBD-dominant cannabis	13.75 mg THC, vaporized	No significant increase in anxiety compared to placebo.
			THC/CBD-equivalent cannabis	13.75 mg THC + 13.75 mg CBD, vaporized	CBD reduced THC-induced anxiety.
Dahlgren et al., 2022 (61)	14 (anxiety)	22 - 64	CBD solution (contains 9.97 mg/mL CBD, 0.23 mg/mL THC)	1 mL sublingually 3x/day (total: 30 mg CBD, <1 mg THC/day)	Anxiety dropped significantly by week 4, evident by week 1.
Ramani et al., 2024 (62)	109 (healthy)	18 - 50	CBD with L-theanine	CBD:30, 60 mg/day; L-theanine: 175–185 mg; oral	Anxiety, cognitive function, exhaustion, and BDNF levels did not significantly change.
Crippa et al., 2004 (11)	10 (healthy)	25 - 42	CBD	400 mg; oral	Reduced anxiety (p < 0.001) compared to placebo

Study	N Participant (condition)	Age (years)	Drug	Doses and administration	Results
Kwee et al., 2024 (63)	69 (SAD)	33.8	CBD	300 mg, 2 hours before therapy; oral	CBD reduced subjective shock expectancy under mild/unknown threat.
	69 (antidepressants user)	33.8	CBD with serotonergic antidepressants	300 mg CBD, before therapy sessions; oral	CBD impaired fear re-extinction compared to placebo.
Gundugurti et al., 2024 (64)	178 (anxiety)	37.2	Nano-dispersible CBD and Placebo	150 mg/mL (300-600 mg/day) for 12 weeks; oral	Significant reduction in GAD-7 and HAM-A score.
Zuardi et al., 2017 (65)	60 (healthy)	18 - 35	CBD	100 mg, 300 mg, 900 mg; oral	CBD (300mg) reduced anxiety in the post-speech phase (inverted U-shaped curve).
Souza et al., 2022 (14)	300 (frontline healthcare workers during COVID-19)	34.5	CBD	150 mg twice daily (300 mg/day) for 28 days; oral	CBD group demonstrates significant reduction in anxiety (GAD-7 scores) compared to control.
Crippa et al., 2011 (66)	10 (SAD)	22	CBD	600 mg; oral	CBD significantly reduced anxiety during public speaking tests versus placebo.
Bloomfield et al., 2022 (67)	24 (stress-inducing arithmetic task)	18 – 27	CBD	600 mg; oral	No difference behaviour between CBD and placebo's group
Bolsoni et al., 2022 (68)	14 (PTSD; sexual trauma)	18 – 60	CBD and Placebo	300 mg; oral	CBD did not significantly reduce anxiety compared to placebo.
	19 (PTSD; nonsexual trauma)				CBD significantly reduced anxiety and cognitive impairment.
Stack et al., 2023 (69)	198 (anxiety disorders)	18 – 70	CBD	0 – 100 mg/day; oral	50% of participants improved anxiety. Significant reductions in anxiety (p < 0.001)
			CBD (dominant): THC	0-100mg/day; oral	Significant reduction in fatigue and anxiety (p < 0.001)
			THC (dominant): CBD	0-38mg/day; oral	61.1% of participants improved anxiety (p = 0.011)
			CBD: THC(Balance)	0-100mg/day; oral	Significant anxiety reduction (p < 0.001)
Hundal et al., 2018 (12)	32 (high paranoia)	18 – 50	CBD	600mg; oral	CBD increased anxiety slightly (non-significant)
Gournay et al., 2023 (70)	63 (elevated trait worry)	18 – 25	CBD	50, 600mg (twice daily for a week); oral	CBD reduced anxiety symptoms in 2 weeks but did not affect worry severity.

CBD: Cannabidiol; PTSD: Post Traumatic Stress Disorder; SAD: Social Anxiety Disorder; VS: Versus

## **DISCUSSION**

### **Mechanism Action of Cannabidiol (CBD)**

According to the results, cannabidiol (CBD) was effective to treat anxiety in both animal and human studies. CBD acts on the 5HT1A receptor, a serotonin receptor, as well as on the endocannabinoid system receptors, namely CB1 and CB2 (4,31,50,51). Research has shown that serotonin has an important role in reducing anxiety. Studies have also found that 5HT1A antagonists (WAY100635), block the CBD's anxiolytic effect, causing anxious behavior in tests such as the Elevated Plus Maze (42,45,47,49,53), Open Field Test (27,35,38,55) and other anxiety assessments (5,30,34,37). Serotonin is a neurotransmitter that regulates mood and is influenced by the Endocannabinoid System (ECS). 5HT1A receptors are distributed in the amygdala, hippocampus, and prefrontal cortex (PFC), all of which regulate anxiety and fear (72). Activation of 5HT1A receptors improves mood, resulting in happiness and calmness (73).

In addition, research has shown the role of CB1 and CB2 receptors in anxiety (4,31,51,55,74). Co-administration of CBD with CB1 antagonist (AM251) and CB2 antagonist (AM630) eliminated the anxiolytic effect in test animals, as AM251 is a CB1 antagonist and AM630 is a CB2 antagonist (4,31,50,51). The ECS impacts the serotonin and GABA systems, with the activation of CB1 receptors has been shown to increase serotonin release in certain parts of the brain, potentially enhancing mood and leading to anxiolytic effects (75).

### **Optimal Dose of Cannabidiol (CBD)**

The dosage (CBD) varied significantly across both types of studies (see Tables 2 and 3). In animal studies, the dosage between 0.3 mg/kg to 30 mg/kg. Various studies have demonstrated an inverted U-shaped curve between dosage and the anxiolytic effect of CBD, indicating that too low a dose might have no effect, while high doses often led to increased anxiety (anxiogenic). The optimal dosage of CBD appeared to be between 5 mg/kg and 10 mg/kg in animal studies (7,8,26,37,49,53). In human clinical studies, the dosage of CBD ranged from 150 mg to 800 mg per day, with varying durations of use and different patient's condition. The 300 mg dose was the one that provided the most optimal anxiolytic effects (58,64,66,69). The 600 mg dose showed anxiolytic effects in studies involving volunteers with anxiety (10,65,71) Stanley (2023) reported increased physical anxiety symptoms at this dose (56). Gradually increasing the CBD dose to 800 mg showed positive results in the treatment of anxiety (59). Variations in optimal dosage can be explained by differences in CBD metabolism among individuals, particularly the role of the CYP3A enzyme, which contributes the most to the metabolic clearance of CBD. High CYP3A activity clears CBD quickly, reducing its levels and necessitating higher doses, while low activity sustains CBD levels longer, allowing for lower doses. Polymorphisms in the CYP3A enzyme can cause differences in metabolic rate (76). This results in variations in drug and metabolite concentrations in the body, which in turn affects the therapeutic response among individuals. This highlights the critical need for personalized dosing in clinical practice.

### **Sex Differences in Response to Cannabidiol (CBD)**

Sex differences indicate varying responses to cannabidiol (CBD). CBD has anxiolytic effects in both sexes (47). However, female rats only reacted to CBD during the LD phase and at a lower dose than male rats. After a short four-day treatment, the response to CBD persisted in female rats at the late diestrus stage but was absent in the proestrus stage (47). This is attributed to a decrease in allopregnanolone (ALLO). The reduction in ALLO increased the sensitivity of extra synaptic neuronal GABA-A receptors, which are associated with the regulation of anxiety behavior, allowing CBD to work more effectively (48,77). Gao et al (2023) also reported the differences in response between male and female rats (77). Female rats showed minimal response to the CBD and THC combination in the EPM test, with no significant changes in anxiety-like behavior observed in most trials (77). These results point out the importance of considering sex-specific aspects in the therapeutic application of CBD for anxiety treatment.

### **Comparison of Cannabidiol (CBD) with Conventional Drugs**

Conventional drugs used to treat anxiety often present side effects (78). For example, benzodiazepines can lead to dependence, sedation, and psychomotor disorders, while selective serotonin reuptake inhibitors (SSRIs) may initially increase anxiety before exhibiting anxiolytic effects and can also lead to dependence (79). In contrast, CBD does not appear to cause dependence (80). Research conducted by Masataka (2019) found that administering 300 mg of CBD daily for 4 weeks showed no signs of dependence in patients (58). Additionally, a study by Zuardi et al

(1982) directly compared CBD with diazepam. Diazepam demonstrated a strong anxiolytic effect with dizziness, whereas CBD had no such effects (2). Other studies have also proven the safety of CBD in its long-term use (59,60,65). These findings imply that CBD may become safer for treating anxiety in the future.

### **Variability of CBD Effects Based on Participant Conditions**

The effects of CBD on anxiety are not solely dependent on dosage but also on individual characteristics and psychological conditions of the participants. A 600 mg dose of CBD exhibits varying pharmacological responses depending on the type of anxiety experienced. In individuals with social anxiety disorder (SAD), this dosage significantly reduces anxiety symptoms during public speaking tests (10). In contrast, among individuals experiencing anxiety induced by arithmetic stress tests, academic examinations, or those with high paranoia, CBD does not demonstrate a clear anxiolytic effect and, in some cases, even increases anxiety levels (12,56,68). These findings suggest that the effectiveness of CBD depends on the nature and severity of anxiety, indicating that dosage and administration methods may need to be adjusted for optimal therapeutic outcomes.

### **CONCLUSION**

CBD has been shown to reduce anxiety, but its effectiveness varies based on the type of disorder and individual conditions. Further research is needed to ensure consistent findings. Future studies should conduct large-scale RCTs to evaluate long-term safety and efficacy, investigate CBD's effects on specific anxiety disorders, assess sex-based differences in response, and explore genetic factors influencing individual variability.

### **AUTHOR'S CONTRIBUTION STATEMENT**

All authors contributed to concept and design. Putu Ayu Putri Fajaryani and I Made Agus Gelgel Wirasuta contributed to literature search. Putu Ayu Putri Fajaryani contributed to data analysis and writing. I Made Agus Gelgel Wirasuta, Pande Made Nova Armita Sari, Dyah Kanya Wati, Putu Indah Budi Apsari, Made Ary Sarasmita contributed to critical revision of manuscript.

### **CONFLICTS OF INTEREST**

The authors declare that they have no conflict of interest regarding the publication of this paper.

### **DECLARATION OF GENERATIVE AI AND AI-ASSISTED TECHNOLOGIES IN THE WRITING PROCESS**

The authors used AI-assisted technology (e.g., Grammarly) only for language editing purposes.

### **SOURCE OF FUNDING STATEMENTS**

This study was conducted without financial support from any public, commercial, or non-profit funding agencies.

### **ACKNOWLEDGMENTS**

We would like to express my sincere gratitude to Udayana University for providing the facilities and space necessary to conduct this research. The support in terms of access to workspace was invaluable in completing this study.

### **BIBLIOGRAPHY**

1. Ma L, Liu M, Liu C, Zhang H, Yang S, An J, et al. Research Progress on the Mechanism of the Antitumor Effects of Cannabidiol. *Mol Basel Switz*. 2024 Apr 24;29(9). doi: 10.3390/molecules29091943.
2. Zuardi AW, Shirakawa I, Finkelfarb E, Karniol IG. Action of cannabidiol on the anxiety and other effects produced by delta 9-THC in normal subjects. *Psychopharmacology (Berl)*. 1982;76(3):245–50. doi: 10.1007/BF00432554.
3. Bitencourt RM, Pamplona FA, Takahashi RN. Facilitation of contextual fear memory extinction and anti-anxiogenic effects of AM404 and cannabidiol in conditioned rats. *Eur Neuropsychopharmacol J Eur Coll Neuropsychopharmacol*. 2008 Dec;18(12):849–59. doi: 10.1016/j.euroneuro.2008.07.001

4. Campos AC, Guimarães FS. Involvement of 5HT1A receptors in the anxiolytic-like effects of cannabidiol injected into the dorsolateral periaqueductal gray of rats. *Psychopharmacology (Berl)*. 2008 Aug;199(2):223–30. doi: 10.1007/s00213-008-1168-x
5. De Gregorio D, McLaughlin RJ, Posa L, Ochoa-Sanchez R, Enns J, Lopez-Canul M, et al. Cannabidiol modulates serotonergic transmission and reverses both allodynia and anxiety-like behavior in a model of neuropathic pain. *Pain*. 2019 Jan;160(1):136–50. 10.1097/j.pain.0000000000001386.
6. Fogaça MV, Reis FMCV, Campos AC, Guimarães FS. Effects of intra-prelimbic prefrontal cortex injection of cannabidiol on anxiety-like behavior: involvement of 5HT1A receptors and previous stressful experience. *Eur Neuropsychopharmacol J Eur Coll Neuropsychopharmacol*. 2014 Mar;24(3):410–9. doi: 10.1016/j.euroneuro.2013.10.012.
7. Guimarães FS, de Aguiar JC, Mechoulam R, Breuer A. Anxiolytic effect of cannabidiol derivatives in the elevated plus-maze. *Gen Pharmacol*. 1994 Jan;25(1):161–4. doi: 10.1016/0306-3623(94)90027-2.
8. Kaplan JS, Wagner JK, Reid K, McGuinness F, Arvila S, Brooks M, et al. Cannabidiol Exposure During the Mouse Adolescent Period Is Without Harmful Behavioral Effects on Locomotor Activity, Anxiety, and Spatial Memory. *Front Behav Neurosci*. 2021;15:711639. . doi: 10.3389/fnbeh.2021.711639.
9. Zieba J, Sinclair D, Sebree T, Bonn-Miller M, Gutterman D, Siegel S, et al. Cannabidiol (CBD) reduces anxiety-related behavior in mice via an FMRP-independent mechanism. *Pharmacol Biochem Behav*. 2019 Jun;181:93–100. doi: 10.1016/j.pbb.2019.05.002
10. Bergamaschi MM, Queiroz RHC, Chagas MHN, de Oliveira DCG, De Martinis BS, Kapczinski F, et al. Cannabidiol reduces the anxiety induced by simulated public speaking in treatment-naïve social phobia patients. *Neuropsychopharmacol Off Publ Am Coll Neuropsychopharmacol*. 2011 May;36(6):1219–26. doi: 10.1038/npp.2011.6.
11. Crippa JA de S, Zuardi AW, Garrido GEJ, Wichert-Ana L, Guarnieri R, Ferrari L, et al. Effects of cannabidiol (CBD) on regional cerebral blood flow. *Neuropsychopharmacol Off Publ Am Coll Neuropsychopharmacol*. 2004 Feb;29(2):417–26. doi: 10.1038/sj.npp.1300340.
12. Hundal H, Lister R, Evans N, Antley A, Englund A, Murray RM, et al. The effects of cannabidiol on persecutory ideation and anxiety in a high trait paranoid group. *J Psychopharmacol Oxf Engl*. 2018 Mar;32(3):276–82. doi: 10.1177/0269881117737400.
13. Hurd YL, Spriggs S, Alishayev J, Winkel G, Gurgov K, Kudrich C, et al. Cannabidiol for the Reduction of Cue-Induced Craving and Anxiety in Drug-Abstinent Individuals With Heroin Use Disorder: A Double-Blind Randomized Placebo-Controlled Trial. *Am J Psychiatry*. 2019 Nov 1;176(11):911–22. doi: 10.1176/appi.ajp.2019.18101191.
14. Souza JDR, Pacheco JC, Rossi GN, de-Paulo BO, Zuardi AW, Guimarães FS, et al. Adverse Effects of Oral Cannabidiol: An Updated Systematic Review of Randomized Controlled Trials (2020-2022). *Pharmaceutics*. 2022 Nov 25;14(12). doi: 10.1176/appi.ajp.2019.18101191.
15. Abu-Sawwa R, Chase A, Fowowe O, Park Y. Effects of Epidiolex® (Cannabidiol) on seizure-related emergency department visits and hospital admissions: A retrospective cohort study. *Epilepsy Behav*. 2022 Feb 1;127:108538. doi: 10.1016/j.yebeh.2021.108538.
16. Bergeria CL, Spindle TR, Cone EJ, Sholler D, Goffi E, Mitchell JM, et al. Pharmacokinetic Profile of  $\Delta^9$ -Tetrahydrocannabinol, Cannabidiol and Metabolites in Blood following Vaporization and Oral Ingestion of Cannabidiol Products. *J Anal Toxicol*. 2022 Jul 14;46(6):583–91. doi: 10.1093/jat/bkab124
17. Moazen-Zadeh E, Chisholm A, Bachi K, Hurd YL. Pharmacokinetics of Cannabidiol: A systematic review and meta-regression analysis. 2023; doi: 10.1101/2023.02.01.23285341
18. Devinsky O, Cilio MR, Cross H, Fernandez-Ruiz J, French J, Hill C, et al. Cannabidiol: pharmacology and potential therapeutic role in epilepsy and other neuropsychiatric disorders. *Epilepsia*. 2014 Jun;55(6):791–802. doi: 10.1111/epi.12631.
19. Reddy D. The Utility of Cannabidiol in the Treatment of Refractory Epilepsy. *Clin Pharmacol Ther*. 2017 Feb;101(2):182–4. doi: 10.1002/cpt.441.
20. Mechoulam R, Parker LA, Gallily R. Cannabidiol: an overview of some pharmacological aspects. *J Clin Pharmacol*. 2002 Nov;42(S1):11S-19S. doi: 10.1002/j.1552-4604.2002.tb05998.x.

21. Taylor L, Gidal B, Blakey G, Tayo B, Morrison G. A Phase I, Randomized, Double-Blind, Placebo-Controlled, Single Ascending Dose, Multiple Dose, and Food Effect Trial of the Safety, Tolerability and Pharmacokinetics of Highly Purified Cannabidiol in Healthy Subjects. *CNS Drugs*. 2018 Nov;32(11):1053–67. doi: 10.1007/s40263-018-0578-5
22. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. 2021 Mar 29;n71. doi: 10.1136/bmj.n71.
23. Uribe-Mariño A, Francisco A, Castiblanco-Urbina MA, Twardowschy A, Salgado-Rohner CJ, Crippa JAS, et al. Anti-aversive effects of cannabidiol on innate fear-induced behaviors evoked by an ethological model of panic attacks based on a prey vs the wild snake *Epicrates cenchria crassus* confrontation paradigm. *Neuropsychopharmacol Off Publ Am Coll Neuropsychopharmacol*. 2012 Jan;37(2):412–21. doi: 10.1038/npp.2011.188.
24. Guimarães FS. Antianxiety effect of cannabidiol in the elevated plus-maze. *Psychopharmacology (Berl)*. 1990;100(4):558–9. doi: 10.1007/BF02244012.
25. ElBatsh MM, Assareh N, Marsden CA, Kendall DA. Anxiogenic-like effects of chronic cannabidiol administration in rats. *Psychopharmacology (Berl)*. 2012 May;221(2):239–47. doi: 10.1007/s00213-011-2566-z.
26. Moreira FA. Anxiolytic-like effect of cannabidiol in the rat Vogel conflict test. *Prog Neuropsychopharmacol Biol Psychiatry*. 2006;30(8):1466–71. doi: 10.1016/j.pnpbp.2006.06.004.
27. Silva-Cardoso GK, Lazarini-Lopes W, Hallak JE, Crippa JA, Zuardi AW, Garcia-Cairasco N, et al. Cannabidiol effectively reverses mechanical and thermal allodynia, hyperalgesia, and anxious behaviors in a neuropathic pain model: Possible role of CB1 and TRPV1 receptors. *Neuropharmacology*. 2021 Oct 1;197:108712. doi: 10.1016/j.neuropharm.2021.108712
28. Almeida V, Levin R, Peres FF, Niigaki ST, Calzavara MB, Zuardi AW, et al. Cannabidiol exhibits anxiolytic but not antipsychotic property evaluated in the social interaction test. *Prog Neuropsychopharmacol Biol Psychiatry*. 2013 Mar 5;41:30–5. doi: 10.1016/j.pnpbp.2012.10.024.
29. Shu G, He Y, Suo J, Wu C, Gong X, Xiang Y, et al. Cannabidiol exhibits anxiolytic-like effects and antipsychotic-like effects in mice models. *Neurosci Lett*. 2024 Mar 15;826:137723. doi: 10.1016/j.neulet.2024.137723.
30. Wanner NM, Colwell M, Drown C, Faulk C. Developmental cannabidiol exposure increases anxiety and modifies genome-wide brain DNA methylation in adult female mice. *Clin Epigenetics*. 2021 Jan 6;13(1):4. doi: 10.1186/s13148-020-00993-4.
31. Chaves YC, Genaro K, Crippa JA, da Cunha JM, Zanoveli JM. Cannabidiol induces antidepressant and anxiolytic-like effects in experimental type-1 diabetic animals by multiple sites of action. *Metab Brain Dis*. 2021 Apr;36(4):639–52. doi: 10.1007/s11011-020-00667-3
32. Chaves YC, Raymundi AM, Waltrick APF, de Souza Crippa JA, Stern CAJ, da Cunha JM, et al. Cannabidiol modulates contextual fear memory consolidation in animals with experimentally induced type-1 diabetes mellitus. *Acta Neuropsychiatr*. 2023 Feb 20;1–11. doi: 10.1017/neu.2023.13.
33. Alegre-Zurano L, López-Arnau R, Luján MÁ, Camarasa J, Valverde O. Cannabidiol Modulates the Motivational and Anxiety-Like Effects of 3,4-Methylenedioxypyrovalerone (MDPV) in Mice. *Int J Mol Sci*. 2021 Aug 2;22(15). doi: 10.3390/ijms22158304
34. Austrich-Olivares A, García-Gutiérrez MS, Illescas L, Gasparyan A, Manzanares J. Cannabinoid CB1 Receptor Involvement in the Actions of CBD on Anxiety and Coping Behaviors in Mice. *Pharm Basel Switz*. 2022 Apr 13;15(4). doi: 10.3390/ph15040473
35. Hsiao YT, Yi PL, Li CL, Chang FC. Effect of cannabidiol on sleep disruption induced by the repeated combination tests consisting of open field and elevated plus-maze in rats. *Neuropharmacology*. 2012 Jan;62(1):373–84. doi: 10.1016/j.neuropharm.2011.08.013
36. Chesworth R, Cheng D, Staub C, Karl T. Effect of long-term cannabidiol on learning and anxiety in a female Alzheimer’s disease mouse model. *Front Pharmacol*. 2022;13:931384. doi: 10.3389/fphar.2022.931384.
37. Rock EM, Limebeer CL, Petrie GN, Williams LA, Mechoulam R, Parker LA. Effect of prior foot shock stress and  $\Delta(9)$ -tetrahydrocannabinol, cannabidiolic acid, and cannabidiol on anxiety-like responding in the light-dark emergence test in rats. *Psychopharmacology (Berl)*. 2017 Jul;234(14):2207–17. doi: 10.1007/s00213-017-4626-5.

38. Melkumyan M, Annaswamy VM, Evans AM, Showemimo OF, McCullers ZE, Sun D, et al. Effects of cannabidiol, with and without  $\Delta^9$ -tetrahydrocannabinol, on anxiety-like behavior following alcohol withdrawal in mice. *Front Neurosci.* 2024;18:1375440. doi: 10.3389/fnins.2024.1375440
39. Marinho ALZ, Vila-Verde C, Fogaça MV, Guimarães FS. Effects of intra-infralimbic prefrontal cortex injections of cannabidiol in the modulation of emotional behaviors in rats: contribution of 5HT<sub>1A</sub> receptors and stressful experiences. *Behav Brain Res.* 2015 Jun 1;286:49–56. doi: 10.1016/j.bbr.2015.02.023
40. Granjeiro EM, Gomes FV, Guimarães FS, Corrêa FMA, Resstel LBM. Effects of intracisternal administration of cannabidiol on the cardiovascular and behavioral responses to acute restraint stress. *Pharmacol Biochem Behav.* 2011 Oct;99(4):743–8. doi: 10.1016/j.pbb.2011.06.027.
41. Campos AC, Guimarães FS. Evidence for a potential role for TRPV1 receptors in the dorsolateral periaqueductal gray in the attenuation of the anxiolytic effects of cannabinoids. *Prog Neuropsychopharmacol Biol Psychiatry.* 2009 Nov 13;33(8):1517–21. doi: 10.1016/j.pnpbp.2009.08.017.
42. Salviato BZ, Raymundi AM, Rodrigues da Silva T, Salemme BW, Batista Sohn JM, Araújo FS, et al. Female but not male rats show biphasic effects of low doses of  $\Delta(9)$ -tetrahydrocannabinol on anxiety: can cannabidiol interfere with these effects? *Neuropharmacology.* 2021 Sep 15;196:108684. doi: 10.1016/j.neuropharm.2021.108684.
43. Breuer A, Haj CG, Fogaça MV, Gomes FV, Silva NR, Pedrazzi JF, et al. Fluorinated Cannabidiol Derivatives: Enhancement of Activity in Mice Models Predictive of Anxiolytic, Antidepressant and Antipsychotic Effects. *PloS One.* 2016;11(7):e0158779. doi: 10.1016/j.heliyon.2024.e25851.
44. Masataka N. Is cannabidiol (CBD) effective to ease separation anxiety? *Heliyon.* 2024 Feb 15;10(3):e25851. doi: 10.1016/j.heliyon.2024.e25851
45. Schleicher EM, Ott FW, Müller M, Silcher B, Sichler ME, Löw MJ, et al. Prolonged Cannabidiol Treatment Lacks on Detrimental Effects on Memory, Motor Performance and Anxiety in C57BL/6J Mice. *Front Behav Neurosci.* 2019;13:94. doi: 10.3389/fnbeh.2019.00094
46. Nazario LR, Antonioli RJ, Capiotti KM, Hallak JEC, Zuardi AW, Crippa JAS, et al. Caffeine protects against memory loss induced by high and non-anxiolytic dose of cannabidiol in adult zebrafish (*Danio rerio*). *Pharmacol Biochem Behav.* 2015 Aug;135:210–6. doi: 10.1016/j.pbb.2015.06.008.
47. Fabris D, Carvalho MC, Brandão ML, Prado WA, Zuardi AW, Crippa JA, et al. Sex-dependent differences in the anxiolytic-like effect of cannabidiol in the elevated plus-maze. *J Psychopharmacol Oxf Engl.* 2022 Dec;36(12):1371–83. doi: 10.1177/02698811221125440.
48. Pérez-Valenzuela E, Hudson R, Uzuneser T, De Felice M, Szkudlarek H, Rushlow W, et al. Sex-Dependent Synergism of an Edible THC: CBD Formulation in Reducing Anxiety and Depressive-like Symptoms Following Chronic Stress. *Curr Neuropharmacol.* 2023 Sep 13; doi: 10.2174/1570159X21666230912101441.
49. Huffstetler CM, Cochran B, May CA, Maykut N, Silver CR, Cedeno C, et al. Single cannabidiol administration affects anxiety-, obsessive compulsive-, object memory-, and attention-like behaviors in mice in a sex and concentration dependent manner. *Pharmacol Biochem Behav.* 2023 Jan;222:173498. doi: 10.1016/j.pbb.2022.173498.
50. Campos AC, Ortega Z, Palazuelos J, Fogaça MV, Aguiar DC, Díaz-Alonso J, et al. The anxiolytic effect of cannabidiol on chronically stressed mice depends on hippocampal neurogenesis: involvement of the endocannabinoid system. *Int J Neuropsychopharmacol.* 2013 Jul;16(6):1407–19. doi: 10.1017/S1461145712001502.
51. Fogaça MV, Campos AC, Coelho LD, Duman RS, Guimarães FS. The anxiolytic effects of cannabidiol in chronically stressed mice are mediated by the endocannabinoid system: Role of neurogenesis and dendritic remodeling. *Neuropharmacology.* 2018 Jun;135:22–33. doi: 10.1016/j.neuropharm.2018.03.001.
52. Gomes FV, Resstel LBM, Guimarães FS. The anxiolytic-like effects of cannabidiol injected into the bed nucleus of the stria terminalis are mediated by 5-HT<sub>1A</sub> receptors. *Psychopharmacology (Berl).* 2011 Feb;213(2–3):465–73. doi: 10.1007/s00213-010-2036-z.
53. Shallcross J, Hámor P, Bechard AR, Romano M, Knackstedt L, Schwendt M. The Divergent Effects of CDPBB and Cannabidiol on Fear Extinction and Anxiety in a Predator Scent Stress Model of PTSD in Rats. *Front Behav Neurosci.* 2019;13:91. doi: 10.3389/fnbeh.2019.00091

54. Morris EM, Kitts-Morgan SE, Spangler DM, McLeod KR, Costa JHC, Harmon DL. The Impact of Feeding Cannabidiol (CBD) Containing Treats on Canine Response to a Noise-Induced Fear Response Test. *Front Vet Sci.* 2020;7:569565. doi: 10.3389/fvets.2020.569565
55. Stern CAJ, Gazarini L, Vanvossen AC, Zuardi AW, Galve-Roperh I, Guimaraes FS, et al.  $\Delta$ -9-Tetrahydrocannabinol alone and combined with cannabidiol mitigate fear memory through reconsolidation disruption. *Eur Neuropsychopharmacol J Eur Coll Neuropsychopharmacol.* 2015 Jun;25(6):958–65. doi: 10.1016/j.euroneuro.2015.02.001.
56. Stanley TB, Ferretti ML, Bonn-Miller MO, Irons JG. A Double-Blind, Randomized, Placebo-Controlled Test of the Effects of Cannabidiol on Experiences of Test Anxiety Among College Students. *Cannabis Cannabinoid Res.* 2023 Dec;8(6):1090–9. doi: 10.1089/can.2022.0062.
57. Martin EL, Strickland JC, Schlien J, Munson J, Jackson H, Bonn-Miller MO, et al. Antidepressant and Anxiolytic Effects of Medicinal Cannabis Use in an Observational Trial. *Front Psychiatry.* 2021;12:729800. doi: 10.3389/fpsy.2021.729800
58. Masataka N. Anxiolytic Effects of Repeated Cannabidiol Treatment in Teenagers With Social Anxiety Disorders. *Front Psychol.* 2019;10:2466. doi: 10.3389/fpsyg.2019.02466
59. Berger M, Li E, Rice S, Davey CG, Ratheesh A, Adams S, et al. Cannabidiol for Treatment-Resistant Anxiety Disorders in Young People: An Open-Label Trial. *J Clin Psychiatry.* 2022 Aug 3;83(5):21m14130. doi: 10.4088/JCP.21m14130.
60. Laczkovics C, Kothgassner OD, Felnhofer A, Klier CM. Cannabidiol treatment in an adolescent with multiple substance abuse, social anxiety and depression. *Neuropsychiatr Klin Diagn Ther Rehabil Organ Ges Osterreichischer Nervenarzte Psychiater.* 2021 Mar;35(1):31–4. doi: 10.1007/s40211-020-00334-0.
61. Hutten NRPW, Arkell TR, Vinckenbosch F, Schepers J, Kevin RC, Theunissen EL, et al. Cannabis containing equivalent concentrations of delta-9-tetrahydrocannabinol (THC) and cannabidiol (CBD) induces less state anxiety than THC-dominant cannabis. *Psychopharmacology (Berl).* 2022 Nov;239(11):3731–41. doi: 10.1007/s00213-022-06248-9.
62. Dahlgren MK, Lambros AM, Smith RT, Sagar KA, El-Abboud C, Gruber SA. Clinical and cognitive improvement following full-spectrum, high-cannabidiol treatment for anxiety: open-label data from a two-stage, phase 2 clinical trial. *Commun Med.* 2022 Nov 2;2(1):139. doi: 10.1038/s43856-022-00202-8.
63. Ramani A, Townsend R, Rodriguez E, Haughian JM, Stewart LK. Daily cannabidiol and L-theanine beverage consumption does not alter anxiety, fatigue, cognitive function, or natural killer cell function: A randomized, controlled trial in healthy, young adults. *Brain Behav Immun Integr.* 2024 Jan 1;5:100045. doi: 10.1016/j.bbii.2024.100045
64. Kwee CMB, van der Flier FE, Duits P, van Balkom AJLM, Cath DC, Baas JMP. Effects of cannabidiol on fear conditioning in anxiety disorders: decreased threat expectation during retention, but no enhanced fear re-extinction. *Psychopharmacology (Berl).* 2024 Apr;241(4):833–47. doi: 10.1007/s00213-023-06512-6.
65. Gundugurti PR, Banda N, Yadlapalli SSR, Narala A, Thatikonda R, Kocherlakota C, et al. Evaluation of the efficacy, safety, and pharmacokinetics of nanodispersible cannabidiol oral solution (150 mg/mL) versus placebo in mild to moderate anxiety subjects: A double blind multicenter randomized clinical trial. *Asian J Psychiatry.* 2024 Jul;97:104073. doi: 10.1016/j.ajp.2024.104073
66. Zuardi AW, Rodrigues NP, Silva AL, Bernardo SA, Hallak JEC, Guimarães FS, et al. Inverted U-Shaped Dose-Response Curve of the Anxiolytic Effect of Cannabidiol during Public Speaking in Real Life. *Front Pharmacol.* 2017;8:259. doi: 10.3389/fphar.2017.00259.
67. Crippa JAS, Derenusson GN, Ferrari TB, Wichert-Ana L, Duran FLS, Martin-Santos R, et al. Neural basis of anxiolytic effects of cannabidiol (CBD) in generalized social anxiety disorder: a preliminary report. *J Psychopharmacol Oxf Engl.* 2011 Jan;25(1):121–30. doi: 10.1177/0269881110379283.
68. Bloomfield MAP, Yamamori Y, Hindocha C, Jones APM, Yim JLL, Walker HR, et al. The acute effects of cannabidiol on emotional processing and anxiety: a neurocognitive imaging study. *Psychopharmacology (Berl).* 2022 May;239(5):1539–49. doi: 10.1007/s00213-022-06070-3.
69. Bolsoni LM, Crippa JAS, Hallak JEC, Guimarães FS, Zuardi AW. The anxiolytic effect of cannabidiol depends on the nature of the trauma when patients with post-traumatic stress disorder recall their trigger event. *Rev Bras Psiquiatr Sao Paulo Braz* 1999. 2022 Jun;44(3):298–307. doi: 10.1590/1516-4446-2021-2317.

70. Stack SK, Wheate NJ, Moloney NC, Abelev SV, Barlow JW, Schubert EA. The Effectiveness and Adverse Events of Cannabidiol and Tetrahydrocannabinol Used in the Treatment of Anxiety Disorders in a PTSD Subpopulation: An Interim Analysis of an Observational Study. *J Pharm Technol JPT Off Publ Assoc Pharm Tech.* 2023 Aug;39(4):172–82. doi: 10.1177/87551225231180796.
71. Gournay LR, Ferretti ML, Bilsky S, Vance E, Nguyen AM, Mann E, et al. The effects of cannabidiol on worry and anxiety among high trait worriers: a double-blind, randomized placebo controlled trial. *Psychopharmacology (Berl).* 2023 Oct;240(10):2147–61. doi: 10.1007/s00213-023-06437-0.
72. Chang J, Liu X, Xue S, Qiu J. An amygdala-centered effective connectivity network in trait anxiety [Internet]. 2023 [cited 2024 Nov 11]. Available from: <https://www.researchsquare.com/article/rs-3122692/v1>. doi: 10.21203/rs.3.rs-3122692/v1
73. Soria-Fregozo C, Perez-Vega MI, Rodríguez-Landa JF, Germán-Ponciano LJ, García-Ríos RI, Mora-Perez A. Association of 5-HT1A Receptors with Affective Disorders. In: Shad KF, editor. *Serotonin - A Chemical Messenger Between All Types of Living Cells* [Internet]. InTech; 2017 [cited 2024 Nov 11]. Available from: <http://www.intechopen.com/books/serotonin-a-chemical-messenger-between-all-types-of-living-cells/association-of-5-ht1a-receptors-with-affective-disorders>. doi: 10.5772/intechopen.68975.
74. Campos AC, de Paula Soares V, Carvalho MC, Ferreira FR, Vicente MA, Brandão ML, et al. Involvement of serotonin-mediated neurotransmission in the dorsal periaqueductal gray matter on cannabidiol chronic effects in panic-like responses in rats. *Psychopharmacology (Berl).* 2013 Mar;226(1):13–24. doi: 10.1007/s00213-012-2878-7. doi: 10.1007/s00213-012-2878-7.
75. Peters KZ, Cheer JF, Tonini R. Modulating the Neuromodulators: Dopamine, Serotonin, and the Endocannabinoid System. *Trends Neurosci.* 2021 Jun;44(6):464–77. doi: 10.1016/j.tins.2021.02.001
76. Beers JL, Fu D, Jackson KD. Cytochrome P450–Catalyzed Metabolism of Cannabidiol to the Active Metabolite 7-Hydroxy-Cannabidiol. *Drug Metab Dispos.* 2021 Oct;49(10):882–91. <https://doi.org/10.1124/dmd.120.000350>
77. Gao B, Qu Y cui, Cai M yu, Zhang Y yin, Lu H tao, Li H xia, et al. Phytochemical interventions for post-traumatic stress disorder: A cluster co-occurrence network analysis using CiteSpace. *J Integr Med.* 2023 Jul 1;21(4):385–96. doi: 10.1016/j.joim.2023.06.006.
78. Schwinghammer TL, DiPiro JT, DiPiro CV, Ellingrod V. *Pharmacotherapy Handbook, Eleventh Edition.*
79. Dunlop BW, Davis PG. Combination Treatment With Benzodiazepines and SSRIs for Comorbid Anxiety and Depression: A Review. *Prim Care Companion J Clin Psychiatry.*
80. Schoedel KA, Levy-Cooperman N. Polydrug users, use of cannabinoids and abuse potential: A focus on cannabidiol (CBD). In: *Cannabis Use, Neurobiology, Psychology, and Treatment* [Internet]. Elsevier; 2023 [cited 2024 Nov 27]. p. 155–69. Available from: <https://linkinghub.elsevier.com/retrieve/pii/B9780323898621000234>. doi: 10.1016/B978-0-323-89862-1.00023-4