

*Narrative Review*

## MYTHBUSTING THE EFFECT OF BCAA ON WOUND HEALING

Ferdian Tanaka\*, Shaza Nathasya Sandakh, Geovan Hananto, & Antonius Setiadi

*Faculty of Medicine, Universitas Brawijaya, Malang, Indonesia*

### ABSTRACT

Nutrition plays a crucial role in the wound-healing process. Nutrition deficiencies can hinder the wound healing process and prolong the recovery period turning the wound into a chronic wound. One nutrient that is rarely discussed is BCAA (Branched Chain Amino Acid). BCAA are essential amino acids that are believed to accelerate the wound healing process. This study aims to investigate the effects of BCAA on the wound healing process. Studies have shown that BCAAs can increase serum albumin levels by preventing the formation of mRNA-PTB (Polypyrimidine Tract-Binding Protein) in liver cells, which activates and accelerates the wound healing process. Administering BCAA as a single treatment is not optimal; therefore, combined therapy with other essential amino acids is necessary. BCAAs can be administered orally, enterally, parenterally, and topically.

**Keywords:** *Nutrition, BCAA, Wound healing, Protein, Amino acid.*

Nutrisi memegang peranan yang sangat penting dalam proses penyembuhan luka. Kekurangan nutrisi dapat memperlambat proses penyembuhan luka dan berakibat luka menjadi luka kronis. Salah satu nutrisi yang jarang dibahas adalah BCAA (*Branched Chain Amino Acid*). BCAA merupakan asam amino esensial berantai dan melalui beberapa penelitian dipercaya dapat mempercepat proses penyembuhan luka. Studi ini ingin mengetahui efek BCAA terhadap proses penyembuhan luka. Dari studi didapatkan BCAA dapat meningkatkan serum albumin dengan cara melalui mekanisme inhibisi dari formasi mRNA-PTB (*Polypyrimidine Tract-binding Protein*) di sel hepar sehingga dapat mengaktivasi dan mempercepat proses penyembuhan luka. Pemberian BCAA sebagai terapi tunggal tidak optimal, sehingga dibutuhkan terapi kombinasi dengan asam amino esensial lainnya. BCAA dapat diberikan secara peroral, nutrisi enteral, parenteral, dan topikal.

**Kata Kunci:** *Nutrisi, BCAA, Penyembuhan luka, Protein, dan Asam amino*

### Conflicts of Interest Statement:

The authors listed in this manuscript declare the absence of any conflict of interest on the subject matter or materials discussed.

## INTRODUCTION

Any damage to the integrity of living tissue can be considered an injury. The skin, the most significant human organ, has the primary function of protecting the organs. Maintaining the integrity of the skin and maintaining the body's optimal wound healing capacity are essential conditions for healthy survival (1). Wound healing is a complex and dynamic sequence of events. Health professionals need biochemistry, physiology, cell biology, and molecular biology knowledge to make decisions about wound care and ensure that wounds heal optimally and do not become

chronic. In addition, wound care can present challenges and burdens on healthcare systems, as long-term wound care requires significant financial resources (2).

Many factors can affect wound healing, and one of them is diet. Inadequate intake or lack of essential nutrients can seriously affect wound healing (3). One well-studied role of BCAAs is their ability to increase the synthesis of albumin, a protein involved in wound healing. When serum albumin falls by ten g/L, mortality increases by 137% and morbidity by 89%, ultimately leading to inhibition of wound healing (4). BCAAs (branched chain amino acids), consisting of leucine, isoleucine, and

Received: 04-02-2025, Revised: 12-06-2025, Accepted: 06-08-2025

Copyright by Tanaka, et al., 2025. | P-ISSN 2089-6492; E-ISSN 2089-9734 | DOI: 10.14228/jprjournal.v12i2.24

Published by Lingkar Studi Bedah Plastik Foundation. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal. This Article can be viewed at [www.jprjournal.com](http://www.jprjournal.com)

valine, are essential nutrients that cannot be produced endogenously by the body and must be obtained from the diet (5). Other studies show that BCAAs positively affect wound healing in patients with diabetic ulcers (6). Due to the limited research on BCAAs, the authors are interested in the effects of BCAAs on the wound healing process. The results of this review can serve as a theoretical basis or reference for other researchers to investigate the effects of BCAAs on wound healing.

## METHOD

Relevant studies published between 1984 and 2023 were identified through searches in PubMed, Cochrane, and ScienceDirect using various combinations of the following keywords: "BCAA," "wound," "healing," "nutrition," "protein," and "amino acid." Additional research papers were obtained by reviewing the reference lists of relevant publications. Studies with limited credibility and those not written in English were excluded. Data extraction was conducted based on the relevance of the information to the topic rather than following a systematic paper selection approach. For a more detailed explanation of the methodology, refer to *Table 1*.

## WOUND HEALING PROCESS

Wound healing consists of three phases: inflammation and hemostasis, proliferation and maturation. Generally, the wound can take 14 to 21 days to heal. The phase of hemostasis in wound healing begins immediately after skin damage. After injury, open blood vessels activate platelets and interact with collagen to release vasoconstrictors such as thromboxane A2 and prostaglandin 2a, stimulating thrombus formation and local vasoconstriction to stop bleeding (1,7). Blood clot formation is due to the production of fibrin matrix. The fibrin matrix consists of fibrinogen, which is converted to fibrin by thrombin after tissue damage. Fibrin

plays a role in forming a blood clot that helps stop bleeding and also provides a framework or platform for platelets and other reactive cells to deposit in the wound. The fibrin matrix also facilitates healing by providing the structure necessary for developing healing cells. The formation and distribution of fibrin matrix is critical for smooth and effective wound healing (7,8). Immediately after the hemostasis phase, local inflammation occurs around the wound, marking the inflammatory phase. In this phase, an inflammatory reaction occurs, involving various cells and inflammatory mediators, which clear the damaged area of debris, reduce the risk of infection, and begin regeneration of the damaged tissue. In the inflammatory phase, vasodilatation occurs, which increases blood flow to the damaged area, and leukocytes, such as neutrophils and monocytes, are directed to the site of damage to clean the damaged tissue (7).

Inflammatory mediators involved in the inflammatory phase of wound healing are prostaglandins, interleukin-1 (IL-1), tumor necrosis factor-alpha (TNF-alpha), and interferon-gamma (IFN-gamma). These various inflammatory mediators regulate the inflammatory response, facilitate inflammatory cell migration and repair, and clear damaged tissue during the inflammatory phase (8,9). IL-6 also plays a role in inducing fever and increasing the release of acute-phase reactants such as heparin. In addition, IL-6 can influence immune responses, including stimulating the proliferation of T and B cells. This indicates that IL-6 is an essential mediator of the inflammatory response in the wound-healing process (10,11). This inflammatory reaction can be characterized by classic symptoms such as redness, swelling, pain, and warmth around the wound. This phase usually lasts several days to weeks before moving into the next phase of the wound-healing process (7).

The second phase of the wound healing process is the proliferation phase. The spreading phase begins about 2-3 days after the injury and lasts about three weeks.

*Table 1.* Summarizing the search technique

Items	Specification
Date of search	2023.07.13
Databases	PubMed, Cochrane, and ScienceDirect
Search terms used	"BCAA", "wound", and "healing" "Branched-Chain Amino Acids", "wound", and "healing"

Items	Specification
Timeframe	"Nutrition", "wound", and "healing" "Protein", "wound", and "healing" "Amino acid", "wound", and "healing" 1984-2023
Inclusion and exclusion criteria	The process was applied to original research papers and reviews written in English that explore the effects of BCAA on wound healing. Articles that did not specifically address the impact of BCAA on wound healing were excluded from the analysis.
Selection proces	The research was independently conducted by FT, SNS, and GH. The data selection process involved identifying the common findings across the searches performed by all three authors.

Fibroblasts play a crucial role at this stage because they produce large amounts of type III collagen to form a temporary matrix. In addition, fibroblasts produce growth factors that stimulate epithelial cell proliferation, angiogenesis, and granulation tissue formation. Fibroblast cells can transform into myofibroblasts under the cytokine TGF- $\beta$ , which promotes wound contraction (11). Angiotensin II also promotes the proliferative phase by maintaining an inflammatory response that stimulates the differentiation and proliferation of fibroblast cells. This indicates that angiotensin II has a significant anti-inflammatory effect during the proliferative phase (10).

In the regeneration phase, granulation tissue is replaced by permanent scar tissue, and collagen production continues, increasing wound adhesion's strength (2). The production of collagen continues for 4-5 weeks. Newly formed collagen initially has thin properties and lies parallel to the skin. Still, over time, this collagen is absorbed and replaced by thicker collagen accumulating along the skin's stretch lines (11). At this point, the tensile strength of the wound continues to increase along with the production of collagen, increasing from 3% in the first week to 20% in the third week. Three months after the injury, the tensile strength of the wound reaches 80% of healthy skin, but it never reaches 100% (2,3). The remodeling phase is an essential stage of the wound healing process that can lead to minimal scar tissue formation and rapid healing. Disturbances in this phase can lead to hypertrophic scars, keloids, or chronic wounds that are difficult to heal. The remodeling phase reorganizes newly deposited collagen into a structured and muscular tissue based on glycosaminoglycans and proteoglycans and is the final step in the wound healing process (11).

## THE ROLE OF THE NUTRITION IN THE WOUND HEALING PROCESS

Nutrition plays an indispensable role in the wound healing process (12). Nutrients are necessary for cell proliferation, protein synthesis, and chemotactic factors, which are also necessary for wound healing (13). Protein, carbohydrates, fats, vitamins, and minerals play an important role in ensuring that the wound healing process runs optimally, while deficiencies can hinder the process and increase the risk of chronic wounds (14, 15).

Based on their requirements in the body, nutrients can be grouped into two categories: macronutrients and micronutrients (12). Macronutrients are required in more significant amounts by the body than micronutrients, but both groups play equally important roles in wound healing (8). Protein, carbohydrates, and fats are macronutrients, while vitamins and minerals are micronutrients. (16).

Wounds put the body into a catabolic state, increasing energy demands (17). Carbohydrates are the primary source of energy in the body (16). The generated power can sustain high metabolic activity for regeneration (14). Fibroblast proliferation, one of the wound healing stages, is highly sensitive to glucose levels (12). If energy is insufficient, the healing process will be prolonged (8).

Fatty acids also play an essential role in the wound healing (18). Phospholipids are crucial components in the formation of the basement membrane of the cell, and fatty acids are one of the components of phospholipids (19). Fats are transporters of vitamins A, D, E, and K. (20) Fats also help provide subcutaneous insulation and bone protection (15). Fats are crucial for the regeneration of epidermis and skin tissue. (18). Therefore, it can be concluded

that fats play an essential role in maintaining the skin's protective capacity.

Currently, 20 vitamins and 16 minerals are known to affect the wound healing process (12). Vitamin A is a fat-soluble vitamin (20). Vitamin A deficiency impairs collagen synthesis (21). Vitamins A and B can promote wound healing by aiding epidermal growth and re-epithelialization (22). Vitamins C and E are potent antioxidants and are known to aid collagen synthesis (23,24). Vitamin D, another fat-soluble vitamin, is essential for calcium absorption, which is vital for bone formation, supports neuromuscular function, strengthens immunity, and accelerates the wound healing (25,26). Fat-soluble vitamin K is found in various green vegetables, kiwi, meat, and eggs (27). Vitamin K is primarily required for coagulation and bone metabolism and is essential for wound healing (28).

Proteins are the primary building blocks for tissue growth and repair (29). Protein intake is important as catabolism occurs during wound healing (30). Protein deficiency affects wound healing capacity, T-cell activity, phagocytic activity, and antibody levels, prolonging the wound healing process and increasing the risk of infected wounds (12, 13).

Proteins are absorbed from the intestines in the form of amino acids (29). Different types of amino acids are known to aid in the wound healing process. Arginine is a type of amino acid that increases collagen synthesis through mechanisms such as enhancing fibroblast synthesis, promoting matrix deposition, angiogenesis, regeneration, antibacterial activity, and improving blood flow by nitrate oxide synthesis (31). Glutamine can aid in wound healing by acting as an energy source, promoting cell growth, helping to produce nitric oxide, reducing C-reactive protein, and acting as an antioxidant (32). Hydroxymethylbutyrate can reduce protein degradation, enhance protein synthesis, reduce apoptosis, and promote cell growth (33).

Branched-chain amino acids (BCAAs), which include leucine, isoleucine, and valine, aid in wound healing. BCAAs are known to support protein synthesis and reduce muscle proteolysis, acting as an energy source that can be stimulated independently of liver function (6).

## DISCUSSION

Branched-chain amino acids (BCAAs), especially leucine, are essential for effective wound healing (34). One of the well-studied roles of BCAAs is their ability to increase albumin synthesis. Albumin is a protein that has an established relationship with the wound healing process. A decrease in serum albumin level of 10 g/L was associated with a 137% increase in mortality, an 89% increase in morbidity, and impaired wound healing (4). Decreased serum albumin levels are also associated with a weakened immune system and delayed healing of wounds caused by infections. This is evident from white blood cell phagocytosis indicators and inflammatory reactions, especially neutrophils (35). The mechanism by which BCAAs increase albumin synthesis involves the activation of the rapamycin complex (mTOR) by its two substrates, ribosomal protein S6 kinase-1 (S6K1) and translational repressor initiation factor (eIF) 4E-binding protein 1 (4E-BP1) in the liver to start the process of translation and synthesis of proteins (36). Leucine can increase albumin synthesis by inhibiting mRNA-PTB (Polypyrimidine Tract-Binding Protein) complex formation in hepatocytes (37).

The BCAA/AAA ratio positively correlates with albumin synthesis and secretion and response to BCAA supplementation. If the BCAA/AAA ratio is low, adding BCAA does not significantly affect the increase in albumin content (38). It is worth noting that BCAA supplementation alone is not optimal for collagen synthesis. This claim is supported by Rajendram et al. (2015), who found that combining BCAA with glutamine could optimize wound healing compared to BCAA alone due to insufficient collagen in wounds with a single treatment (39).

This is consistent with Murakami et al. (2013) and Crozier et al. (2005), who showed that supplementation of BCAA with glutamine significantly increased skin tropocollagen synthesis, suggesting a synergistic effect between BCAA and glutamine (40,41). Montoya et al. (1989) also showed that the administration of an infusion containing only 45% BCAA did not significantly accelerate wound healing compared to a conventional parenteral infusion containing 8% BCAA (42). Zhang et al. (2004) presented an amino acid mixture containing high doses of leucine,

isoleucine, valine, lysine, phenylalanine, histidine, threonine, methionine, tryptophan, alanine, arginine, glycine, proline, serine, and tyrosine that had anabolic effects on proteins synthesis in rabbit skin wounds. They hypothesized that increasing the availability of amino acids would improve the process of protein synthesis. (43). Wound protein anabolism can initiate the wound healing process because new proteins must be synthesized to repair damaged tissue and improve cellular functions such as cell migration and cell differentiation (43). Mixing BCAA with glycine and BCAA with glycine and arginine synergistically significantly increased the fractional synthesis rate of skin tropocollagen protein in UVB-treated rats (44). In addition, Nair and Chong (2020) found that BCAAs can increase the production of growth hormones, which can help in the wound healing process (45). BCAAs are also related to strengthening the immune system, but the exact mechanism is still not well understood (34).

Moss et al. (1970, 1975, and 1984) found that elevated BCAA levels in postoperative patients decreased protein catabolism and increased protein anabolism. This is reflected in the increased production of plasma proteins and wound healing when the following patients are administered a high-nitrogen-calorie diet (Vivonex R containing 3600-4800 kcal and 150-200 g protein equivalent) via a nasoduodenal tube following elective open cholecystectomy. A significant increase in serum BCAA levels was observed within the first 2 hours after dosing and remained high throughout the duration of tube feeding (46-48). BCAAs can act as a precursor to glutamine and alanine synthesis during catabolic states, providing additional nutrition and maintaining nitrogen balance after surgery (49). Research conducted by Choudry et al. (2006) found that administering the parenteral nutrition fluid BFLUID (containing 30% BCAA) resulted in higher tensile strength compared to the group given Physio35 (which does not contain amino acids) ( $p < 0.05$ ). There was a significant increase in type 1 collagen mRNA expression ( $p < 0.05$ ) compared to the Physio35 group and an increase in type 3 collagen mRNA expression compared to the Physio35 group ( $p < 0.1$ ) (49). Delany HM et al. (2011) showed no difference in wound healing between enteral and

parenteral administration of a high BCAA diet in patients undergoing major surgery. (50).

BCAAs can also be given in topical dressing formulations together with other amino acids. One study showed that topical dressings containing leucine, glycine, lysine, and sodium hyaluronate improved wound healing in rats. The mechanism is thought to involve increased immunolocalization of TGF-beta in the dressing region. Stimulated TGF-beta can increase fibroblast production and eNOS secretion, and it is designed to increase collagen and reduce inflammation. Leucine is an essential amino acid that promotes protein synthesis (51).

It's worth noting that it takes 6.5g of BCAAs to reach plasma concentrations, which is five times higher than human basal levels. Leucine, in particular, is a primary stimulator for collagen synthesis in wound healing, especially in conditions of protein undernutrition. However, it's important to remember that single therapy with BCAAs alone is not effective for dermal collagen synthesis (39).

The maximum tolerated dose for leucine, isoleucine, and valine is 500 mg/kg/day (52). It is important to note that BCAA supplements should be used with caution in individuals with co-morbidities such as diabetes and psychiatric disorders. Increased BCAA concentrations promote insulin resistance, which can accelerate other diabetic complications. BCAAs can also increase aggression and lower mood by reducing serotonin synthesis (53).




Figure 1. Mechanism of BCAA on Wound Healing Process.

## CONCLUSION

BCAA is an essential amino acid that can help in the wound healing process. However, the availability of other essential amino acids such as glutamine, arginine, glycine, and others is necessary to optimize their effectiveness. BCAAs can be administered orally, parenterally, enterally, or topically as wound dressings.

### Correspondence regarding this article should be addressed to:

Ferdian Tanaka, MD. Brawijaya University Faculty of Medicine, Malang, Indonesia, Malang, 65142, Indonesia.  
E-Mail: ferdiantanaka@gmail.com

## ACKNOWLEDGEMENT

None

## REFERENCES

- Almadani YH, Vorstenbosch J, Davison PG, Murphy AM. Wound Healing: A Comprehensive Review. *Semin Plast Surg.* 2021 Aug;35(3):141-4.
- Chhabra S, Chhabra N, Kaur A, Gupta N. Wound Healing Concepts in Clinical Practice of OMFS. *J Maxillofac Oral Surg.* 2017 Dec;16(4):403-23.
- Williams JZ, Barbul A. Nutrition and Wound Healing. *Crit Care Nurs Clin North Am.* 2012 Jun;24(2):179-200.
- Khoirunnisa AA, Mustafa A, Rahman N. Effect of Giving Snakehead Fish Extract (*Channa striata*) on Albumin Levels, Neutrophil, and Lymphocyte in Hypoalbuminemia Patients. 2022;01(02).
- Institute of Medicine (US) Committee on Nutrition T, Erdman J, Oria M, Pillsbury L. Branched-Chain Amino Acids. In: *Nutrition and Traumatic Brain Injury: Improving Acute and Subacute Health Outcomes in Military Personnel* [Internet]. National Academies Press (US); 2011 [cited 2024 Mar 20]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK209312/>
- Yk M, Yh L, Hc L, Ch J, Cc C, Sh L. Effect of Branched-Chain Amino Acids on Wound Healing in Patients with Diabetic Foot Ulcers. *Plast Surg.* 2023;7.
- Wang PH, Huang BS, Horng HC, Yeh CC, Chen YJ. Wound healing. *J Chin Med Assoc.* 2018 Feb;81(2):94-101.
- Anderson K, Hamm RL. Factors That Impair Wound Healing. *J Am Coll Clin Wound Spec.* 2012 Dec;4(4):84-91.
- Gonzalez AC de O, Costa TF, Andrade Z de A, Medrado ARAP. Wound healing - A literature review. *An Bras Dermatol.* 2016;91(5):614-20.
- Buchanan PJ, Kung TA, Cederna PS. Evidence-Based Medicine: Wound Closure. *Plast Reconstr Surg.* 2016 Sep;138(3S):257S-270S.
- Janis JE, Harrison B. Wound Healing: Part I. Basic Science. *Plast Reconstr Surg.* 2016 Sep;138(3S):9S-17S.
- Grada A, Phillips TJ. Nutrition and cutaneous wound healing. *Clin Dermatol.* 2022 Mar;40(2):103-13.
- Dryden SV, Shoemaker WG, Kim JH. Wound Management and Nutrition for Optimal Wound Healing. *Atlas Oral Maxillofac Surg Clin.* 2013 Mar;21(1):37-47.
- De Albuquerque PBS, Rodrigues NER, Silva PMDS, De Oliveira WF, Correia MTDS, Coelho LCBB. The Use of Proteins, Lipids, and Carbohydrates in the Management of Wounds. *Molecules.* 2023 Feb 7;28(4):1580.
- Stechmiller JK. Understanding the Role of Nutrition and Wound Healing. *Nutr Clin Pract.* 2010 Feb;25(1):61-8.
- Campos V, Tappy L, Bally L, Sievenpiper JL, Lê KA. Importance of Carbohydrate Quality: What Does It Mean and How to Measure It? *J Nutr.* 2022 May;152(5):1200-6.
- Ghaly P, Iliopoulos J, Ahmad M. The role of nutrition in wound healing: an overview. *Br J Nurs.* 2021 Mar 11;30(5):S38-42.
- Hokynková A, Nováková M, Babula P, Sedláčková M, Paulová H, Hlaváčová M, et al. Fatty Acid Supplementation Affects Skin Wound Healing in a Rat Model. *Nutrients.* 2022 May 27;14(11):2245.
- Morita S ya, Ikeda Y. Regulation of membrane phospholipid biosynthesis in mammalian cells. *Biochem Pharmacol.* 2022 Dec;206:115296.
- Stevens SL. Fat-Soluble Vitamins. *Nurs Clin North Am.* 2021 Mar;56(1):33-45.
- Zinder R, Cooley R, Vlad LG, Molnar JA. Vitamin A and Wound Healing. *Nutr Clin Pract.* 2019 Dec;34(6):839-49.

22. Barchitta M, Maugeri A, Favara G, Magnano San Lio R, Evola G, Agodi A, et al. Nutrition and Wound Healing: An Overview Focusing on the Beneficial Effects of Curcumin. *Int J Mol Sci*. 2019 Mar 5;20(5):1119.
23. Moores J. Vitamin C: a wound healing perspective. *Br J Community Nurs*. 2013 Dec 1;18(Sup12):S6–11.
24. Yarahmadi A, Saeed Modaghegh MH, Mostafavi-Pour Z, Azarpira N, Mousavian A, Bonakdaran S, et al. The effect of platelet-rich plasma-fibrin glue dressing in combination with oral vitamin E and C for treatment of non-healing diabetic foot ulcers: a randomized, double-blind, parallel-group, clinical trial. *Expert Opin Biol Ther*. 2021 May 4;21(5):687–96.
25. Lin J, Mo X, Yang Y, Tang C, Chen J. Association between vitamin D deficiency and diabetic foot ulcer wound in diabetic subjects: A meta-analysis. *Int Wound J*. 2023 Jan;20(1):55–62.
26. Siregar FD, Hidayat W. The Role of Vitamin D on the Wound Healing Process: A Case Series. *Int Med Case Rep J*. 2023 Apr;Volume 16:227–32.
27. Halder M, Petsophonsakul P, Akbulut A, Pavlic A, Bohan F, Anderson E, et al. Vitamin K: Double Bonds beyond Coagulation Insights into Differences between Vitamin K1 and K2 in Health and Disease. *Int J Mol Sci*. 2019 Feb 19;20(4):896.
28. Tang S, Ruan Z, Ma A, Wang D, Kou J. Effect of vitamin K on wound healing: A systematic review and meta-analysis based on preclinical studies. *Front Pharmacol*. 2022 Dec 2;13:1063349.
29. Wang X, Yu Z, Zhou S, Shen S, Chen W. The Effect of a Compound Protein on Wound Healing and Nutritional Status. Tebyaniyan H, editor. *Evid Based Complement Alternat Med*. 2022 Mar 24;2022:1–12.
30. Rodriguez NA, Jeschke MG, Williams FN, Kamolz L, Herndon DN. Nutrition in Burns: Galveston Contributions. *J Parenter Enter Nutr*. 2011 Nov;35(6):704–14.
31. Chu AS, Delmore B. Arginine: What You Need to Know for Pressure Injury Healing. *Adv Skin Wound Care*. 2021 Dec;34(12):630–6.
32. Arribas-López E, Zand N, Ojo O, Snowden MJ, Kochhar T. The Effect of Amino Acids on Wound Healing: A Systematic Review and Meta-Analysis on Arginine and Glutamine. *Nutrients*. 2021 Jul 22;13(8):2498.
33. Hung SY, Tsai JS, Yeh JT, Chen KH, Lin CN, Yang HM, et al. Amino acids and wound healing in people with limb-threatening diabetic foot ulcers. *J Diabetes Complications*. 2019 Oct;33(10):107403.
34. Thomson RL, Buckley JD. Protein hydrolysates and tissue repair. *Nutr Res Rev*. 2011 Dec;24(2):191–7.
35. Rosyidi RM, Januarman, Priyanto B, Islam AA, Hatta M, Bukhari A. The Effect of Snakehead Fish (*Channa striata*) Extract Capsule to the Albumin Serum Level of Post-operative Neurosurgery Patients. *Biomed Pharmacol J*. 2019 Jun 25;12(2):893–9.
36. Wang X, Proud CG. mTORC1 signaling: what we still don't know. *J Mol Cell Biol*. 2011 Aug;3(4):206–20.
37. Kuwahata M, Yoshimura T, Sawai Y, Amano S, Tomoe Y, Segawa H, et al. Localization of polypyrimidine-tract-binding protein is involved in the regulation of albumin synthesis by branched-chain amino acids in HepG2 cells. *J Nutr Biochem*. 2008 Jul;19(7):438–47.
38. Marchesini G, Bianchi G, Merli M, Amodio P, Panella C, Loguercio C, et al. Nutritional supplementation with branched-chain amino acids in advanced cirrhosis: a double-blind, randomized trial. *Gastroenterology*. 2003 Jun;124(7):1792–801.
39. Rajendram R, Preedy VR, Patel VB, editors. *Branched Chain Amino Acids in Clinical Nutrition: Volume 2* [Internet]. New York, NY: Springer New York; 2015 [cited 2024 Mar 25]. Available from: <https://link.springer.com/10.1007/978-1-4939-1914-7>
40. Murakami H, Shimbo K, Takino Y, Kobayashi H. Combination of BCAAs and glutamine enhances dermal collagen protein synthesis in protein-malnourished rats. *Amino Acids*. 2013 Mar;44(3):969–76.
41. Crozier SJ, Kimball SR, Emmert SW, Anthony JC, Jefferson LS. Oral leucine administration stimulates protein synthesis in rat skeletal muscle. *J Nutr*. 2005 Mar;135(3):376–82.
42. Montoya A, Gómez-Lechón MJ, Castell JV.

- Influence of Branched-Chain Amino Acid Composition of Culture Media on the Synthesis of Plasma Proteins by Serum-Free Cultured Rat Hepatocytes. *In Vitro Cell Dev Biol.* 1989;25(4):358-64.
43. Zhang XJ, Chinkes DL, Wolfe RR. Leucine supplementation has an anabolic effect on proteins in rabbit skin wound and muscle. *J Nutr.* 2004 Dec;134(12):3313-8.
44. Murakami H, Shimbo K, Inoue Y, Takino Y, Kobayashi H. Importance of amino acid composition to improve skin collagen protein synthesis rates in UV-irradiated mice. *Amino Acids.* 2012 Jun;42(6):2481-9.
45. Nair HK, Chong SS. Evaluating the efficacy of protein-based nutritional supplements in patients with chronic wounds: a pilot study. *Case Rep.* 2020;3(3).
46. Moss G, Koblenz G. Postoperative positive nitrogen balance: effects on wound and plasma protein synthesis. *Surg Forum.* 1970;21:71-3.
47. Moss G, Bierenbaum A, Bova F, Slavin JA. Postoperative metabolic patterns following immediate total nutritional support: hormone levels, DNA synthesis, nitrogen balance, and accelerated wound healing. *J Surg Res.* 1976 Dec;21(6):383-93.
48. CLINICAL AND EXPERIMENTAL EFFECTS OF IMMEDIATE POSTCOLECTOMY... : Critical Care Medicine [Internet]. [cited 2024 Mar 31]. Available from: [https://journals.lww.com/ccmjourn/citation/1984/03000/clinical\\_and\\_experimental\\_effects\\_of\\_immediate.23.aspx](https://journals.lww.com/ccmjourn/citation/1984/03000/clinical_and_experimental_effects_of_immediate.23.aspx)
49. Choudry HA, Pan M, Karinch AM, Souba WW. Branched-chain amino acid-enriched nutritional support in surgical and cancer patients. *J Nutr.* 2006 Jan;136(1 Suppl):314S-8S.
50. Delany HM, Teh E, Dwarka B, Levenson SM. Infusion of Enteral vs Parenteral Nutrients Using High-Concentration Branch-Chain Amino Acids: Effect on Wound Healing in the Postoperative Rat. *J Parenter Enter Nutr.* 1991 Jul;15(4):464-8.
51. Topical application of dressing with amino acids improves cutaneous wound healing in aged rats - PubMed [Internet]. [cited 2024 Mar 27]. Available from: <https://pubmed.ncbi.nlm.nih.gov/19560799/>
52. Elango R. Tolerable Upper Intake Level for Individual Amino Acids in Humans: A Narrative Review of Recent Clinical Studies. *Adv Nutr.* 2023 Jul;14(4):885-94.
53. Holeček M. Side Effects of Amino Acid Supplements. *Physiol Res.* 2022 Mar 22;29-45.