

## **Formulation And Evaluation Of Lip Balm Preparation From Watermelon Seed Extract (*Citrullus lanatus*) As Lip Moisturizer**

**Alter Y. Runtu<sup>1\*</sup>, Soniya Tumuju<sup>1</sup>, Douglas N. Pareta<sup>1</sup>, Andry Natanel<sup>1</sup>,  
Anggelina A. Pangalila<sup>1</sup>, Aldolfina Sumangando<sup>2</sup>, Marshella M. Mokoagouw<sup>1</sup>**

<sup>1</sup>Department of Pharmacy, Faculty of Mathematics and Natural Sciences, Christian University of Indonesia in Tomohon

<sup>2</sup>Department of Biology, Faculty of Mathematics and Natural Sciences, Christian University of Indonesia in Tomohon

\*Corresponding author; [alter18runtu@gmail.com](mailto:alter18runtu@gmail.com)

Accepted: 26 September 2025; Approved : 16 Oktober 2025

### **ABSTRAK**

Penelitian ini bertujuan untuk memformulasikan dan mengevaluasi sediaan lip balm yang mengandung ekstrak biji semangka (*Citrullus lanatus*) sebagai agen pelembap bibir alami yang aman dan efektif, seiring meningkatnya kebutuhan akan kosmetik berbahan dasar tanaman. Ekstrak diperoleh melalui metode maserasi, kemudian dibuat empat formula lip balm dengan konsentrasi ekstrak sebesar 0%, 3%, 6%, dan 9%. Evaluasi sediaan meliputi uji karakteristik fisik, uji iritasi, uji hedonik, serta uji stabilitas selama penyimpanan 10 minggu. Hasil penelitian menunjukkan bahwa seluruh formula bersifat stabil, memiliki nilai pH dan titik lebur yang sesuai, serta tidak menimbulkan iritasi. Peningkatan konsentrasi ekstrak secara signifikan meningkatkan daya sebar dan efektivitas pelembap, dengan formula 9% menjadi yang paling disukai oleh panelis. Dengan demikian, ekstrak biji semangka berpotensi sebagai bahan aktif alami yang aman dan efektif untuk sediaan perawatan bibir.

**Kata kunci:** *Citrullus lanatus*, lip balm, ekstrak biji semangka, pelembap alami, evaluasi formulasi.

### **ABSTRACT**

This study aimed to formulate and evaluate a lip balm preparation containing watermelon seed extract (*Citrullus lanatus*) as a safe and effective natural lip moisturizer, in response to the increasing demand for plant-based cosmetic products. The extract was obtained using the maceration method, and four lip balm formulations were prepared with extract concentrations of 0%, 3%, 6%, and 9%. The preparations were evaluated through physical characterization tests, irritation assessment, hedonic evaluation, and stability testing over a 10-week storage period. The results indicated that all formulations were stable, exhibited acceptable pH values and melting points, and did not cause irritation. Increasing extract concentration significantly improved spreadability and moisturizing performance, with the 9% formulation receiving the highest preference among panelists. These findings suggest that watermelon seed extract has strong potential as a natural and safe active ingredient for lip care formulations.

**Keywords:** *Citrullus lanatus*, lip balm, watermelon seed extract, natural moisturizer, formulation evaluation.

## **1. INTRODUCTION**

Cosmetics, particularly lip care products, have become an essential necessity because the

skin on the lips is highly vulnerable to environmental factors such<sup>1</sup> as extreme temperatures and UV exposure, which may cause dryness and chapping<sup>2</sup>. Although many

commercial products still contain synthetic<sup>3</sup> ingredients, such as petroleum jelly, which may potentially lead to adverse effects, there is an increasing trend toward the use of natural-based cosmetic ingredients<sup>4</sup>.

Previous research has shown the potential of watermelon fruit and rind extracts as moisturizers and UV protectors<sup>3</sup>. However, exploration of watermelon seeds for similar applications is still limited. Empirically, watermelon seeds have long been used in traditional medicine<sup>5</sup>, and phytochemical studies show the presence of beneficial compounds such as flavonoids and essential fatty acids (especially linoleic acid)<sup>6</sup>, which has natural emollient properties and potentially helps maintain lip moisture<sup>5</sup>. Therefore, this study aims to fill this gap by formulating a lip balm preparation using watermelon seed extract as the main active ingredient and testing its effectiveness as a lip moisturizer.

## 2. RESEARCH METHOD

### Time and Place of Research

This research was conducted from March to April 2025 at the Laboratory of the Faculty of Mathematics and Natural Sciences, Indonesian Christian University Tomohon.

### Materials and Instruments

The instruments used in this research included an analytical balance, blender, spatula, porcelain dish, measuring cylinder, water bath, digital pH meter, microscope slide, beakers, filter paper, glass jars, and a rotary evaporator.

The materials consisted of watermelon seed extract, n-hexane, cera alba, lanolin, methylparaben, glycerin, olive oil, and rabbits used as experimental subjects for the irritation test.

This research employed an experimental laboratory approach aimed at formulating, preparing, and evaluating lip balm preparations containing watermelon seed extract<sup>7</sup>.

### Research procedure

#### 1. Preparation of Watermelon Seed Extract

Fresh watermelon seeds were obtained from a plantation in Matani Village, South Minahasa. A total of 705 g of seeds were washed and oven-dried at 60°C for 60 minutes<sup>7</sup> until the weight decreased to 455 g. The dried seeds were then ground into powder<sup>8</sup>. Extraction was carried out using the maceration method by soaking 455 g of seed powder in 1 L of n-hexane for 3 × 24 hours. The resulting macerate was filtered and concentrated using a rotary evaporator at 40°C until a thick extract was obtained<sup>2</sup>.

#### 2. Formulation of Lip Balm Preparations

Four lip balm formulations were prepared: a control formulation without extract (F0) and three formulations containing watermelon seed extract at concentrations of 3% (F1), 6% (F2), and 9% (F3). All formulations used the same base composition, consisting of 10% cera alba, 10% lanolin, 0.03% methylparaben, 5% glycerin, and olive oil added up to 10 g.

The preparations were produced using the melting method. Cera alba and lanolin were melted at 62–65°C, followed by the addition of methylparaben, glycerin, and olive oil with continuous stirring until a homogeneous mixture was formed<sup>8</sup>. Watermelon seed extract was then incorporated into each formula, poured into molds, and allowed to solidify at room temperature<sup>4</sup>.

### Type of research

**Table 1.** Formula LipBalm

| Material                | Formula ( amount %) |       |       |       | Description      |
|-------------------------|---------------------|-------|-------|-------|------------------|
|                         | F0                  | F1    | F2    | F3    |                  |
| (1)                     | (2)                 | (3)   | (4)   | (5)   | (6)              |
| Watermelon seed extract | -                   | 3     | 6     | 9     | Active substance |
| Cera alba               | 10                  | 10    | 10    | 10    | Candle           |
| Lanolin                 | 10                  | 10    | 10    | 10    | Fat              |
| Metil paraben           | 0,03                | 0,03  | 0,03  | 0,03  | Preservative     |
| Gliserin                | 5                   | 5     | 5     | 5     | Humectant        |
| Olive oil               | Ad 10               | Ad 10 | Ad 10 | Ad 10 | Oil phase        |

### 3. Evaluation of Lip Balm Preparations

The lip balm formulations prepared in this research were evaluated through the following tests:

**Organoleptic Test:** Visual observation of changes in color, odor, and texture during one week of storage<sup>9</sup>.

**Homogeneity Test:** Examination of the preparation between two glass plates to detect coarse particles<sup>2</sup>.

**pH Test:** Measurement using a digital pH meter; preparations were considered acceptable if the pH ranged from 4.5 to 7<sup>10</sup>.

**Spreadability Test:** Determination of spread diameter using the glass plate method<sup>11</sup>.

**Melting Point Test:** Assessment of melting point at 50°C<sup>12</sup>.

**Irritation Test:** Conducted on rabbits using the patch test method, with erythema and edema observed for 72 hours<sup>13</sup>.

**Hedonic Test:** Preference evaluation by 30 panelists regarding aroma, texture, ease of application, and moisturizing effect<sup>14</sup>.

### 4. Stability Test

Stability testing in this research was carried out over a period of 10 weeks by monitoring organoleptic characteristics, pH, and homogeneity of the formulations<sup>7</sup>.

### Data Analysis

Quantitative data obtained in this research from melting point, spreadability, and hedonic tests were analyzed using Analysis of Variance (ANOVA) with SPSS software to determine significant differences among formulations. Qualitative data from organoleptic evaluation, homogeneity, pH, and irritation tests were presented descriptively in tabular form.

## 3. RESULTS AND DISCUSSION

### Production of Watermelon Seed Extract

The extraction of watermelon seeds obtained from the Empat Bersaudara plantation in Matani Village, South Minahasa, produced a thick extract with an orange to reddish-yellow appearance. The extraction yielded 20 g of concentrated extract from 455 g of dried seed powder. The procedure began with oven-drying at 60°C to reduce the moisture content of the seeds, followed by maceration using n-hexane as the solvent. This method was selected because n-hexane effectively extracts lipid components, including linoleic acid and other essential fatty acids, which function as natural emollients. The resulting extract was subsequently incorporated as the active ingredient into lip balm formulations at concentrations of 3% (F1), 6% (F2), and 9% (F3), while the formulation without extract served as the negative control (F0).

### Evaluation Results of LipBalm Preparation

#### 1. Organoleptic test

Organoleptic observations conducted over one week of storage demonstrated that all lip balm formulations exhibited good physical stability. The control formulation (F0) appeared cloudy white with a mild beeswax aroma. In contrast, formulations containing watermelon seed extract (F1, F2, and F3) showed colors ranging from pale yellow to bright yellow and possessed a characteristic nutty odor derived from the extract. All formulations maintained a semi-solid consistency, which is considered suitable for lip application. The gradual increase in color intensity with higher extract concentrations indicates that watermelon seed extract contributes to the pigmentation of the preparation<sup>15</sup>.

**Table 2.** Organoleptic Test Results

| Formula | Color         | Smell            | Texture    |
|---------|---------------|------------------|------------|
| (1)     | (2)           | (3)              | (4)        |
| F0      | Murky white   | A little beeswax | Semi solid |
| F1      | Pale yellow   | A little nutty   | Semi solid |
| F2      | Pale yellow   | A little nutty   | Semi solid |
| F3      | Bright yellow | A little nutty   | Semi solid |

**2. Homogeneity Test**

The homogeneity test confirmed that all formulations (F0–F3) exhibited a uniform distribution of ingredients. No coarse particles or granules were observed when the preparations were spread between two glass plates.

This result indicates that the melting and mixing process successfully produced homogeneous lip balm preparations. The stability of the formulations may be attributed to the wax–oil phase formed by *cera alba* and olive oil, supported by lanolin, which acts as an emulsifying agent<sup>12</sup>.

**Table 3.** Homogeneity Test Results

| Formula | Homogeneous | Non-homogeneous |
|---------|-------------|-----------------|
| (1)     | (2)         | (3)             |
| F0      | √           | -               |
| F1      | √           | -               |
| F2      | √           | -               |
| F3      | √           | -               |

**3. Melting Point Test**

The melting point evaluation demonstrated that all formulations complied with the Indonesian National Standard (SNI 16-4399, 1996), which specifies an acceptable melting point range of 50–70°C. The average melting point increased with higher extract concentrations, recorded as 50°C (F0), 51°C (F1), 52°C (F2), and 55°C (F3). This finding suggests that watermelon seed extract influences the structural properties of the

preparation, enhancing its resistance to heat<sup>10</sup>. Statistical analysis using ANOVA (p-value < 0.001 and F-value = 289.000) demonstrated a significant difference in melting points between formulas. This confirms that the extract concentration substantially affects the thermal stability of the product, which plays an important role in maintaining the shape and consistency of lip balm at room temperature<sup>16</sup>.

**Table 4.** Melting Point Test Result

| Formula | Replication | Temperature |
|---------|-------------|-------------|
| (1)     | (2)         | (3)         |
| F0      | 1           | 50°C        |
|         | 2           | 50°C        |
| F1      | 1           | 51°C        |
|         | 2           | 51,1°C      |
| F2      | 1           | 52°C        |
|         | 2           | 52,3°C      |
| F3      | 1           | 55°C        |
|         | 2           | 55,2°C      |

**4. Spreadability Test**

Spreadability is an essential parameter that determines the ease of product application. In this research, all lip balm formulations demonstrated adequate spreadability, with diameters ranging from 3.25 cm to 4.50 cm. These values fall within the optimal spreadability range of 3–5 cm.

ANOVA results indicated a statistically significant difference in spreadability among formulations (p-value = 0.022; F = 10.692), suggesting that the concentration of watermelon seed extract influences the spreading capacity of the preparation<sup>16</sup>.

**Table 5.** Dispersal Test Results

| Formula | Replication | Spread power (cm) |
|---------|-------------|-------------------|
| (1)     | (2)         | (3)               |
| F0      | 1           | 3,2               |
|         | 2           | 3,3               |
| F1      | 1           | 3,4               |
|         | 2           | 4                 |
| F2      | 1           | 3,6               |
|         | 2           | 4,4               |
| F3      | 1           | 3,5               |
|         | 2           | 4,5               |

5. pH Test

The pH test results indicate that all formulas have pH values that are safe and comfortable for lip skin, ranging from 6.50 to 6.86. These values are within the physiological

pH range of lip skin (4.5–7.0), which aims to ensure that the lip balm preparations in this study will not cause irritation. The decrease in pH occurs with the addition of the extract, but it remains safe and stable<sup>10</sup>.

**Table 6.** pH Test Results


| Formula | Homogeneous |
|---------|-------------|
| (1)     | (2)         |
| F0      | 6,86        |
| F1      | 6,68        |
| F2      | 6,50        |
| F3      | 6,59        |

6. Hedonic Test

A hedonic evaluation involving 30 panelists revealed variations in preference among the formulations, particularly regarding moisturizing performance and ease of application. The formulation containing 9% watermelon seed extract (F3) was the most favored,

receiving the highest preference scores for moisture (70%) and ease of application (81.43%).

ANOVA results confirmed significant differences among formulations, indicating that higher extract concentrations enhance product performance due to the presence of emollient fatty acids<sup>17</sup>.



**Figure 1.** Hedonic Test Results

**Table 7.** Results of the Repeated Measures ANOVA Test

| Parameter           | <i>Mauchly's Test (p)</i> | <i>E(df1, df2)</i> | <i>p-value</i> | <i>Partial Eta Squared</i> | description                                                                                                    |
|---------------------|---------------------------|--------------------|----------------|----------------------------|----------------------------------------------------------------------------------------------------------------|
| humidityari         | 0,694                     | F(2,58)<br>= 9,256 | <<br>0,001     | 0,242                      | There is a significant difference between the formulas. Formula 1 differs significantly from formulas 2 and 3. |
| Aroma               | 0,192                     | F(2,58)<br>= 0,335 | <<br>0,717     | 0,011                      | There is no significant difference between formulas in terms of aroma                                          |
| Dosage form         | 0,060                     | F(2,58)<br>= 0,921 | <<br>0,404     | 0,031                      | There is no significant difference between the formulas. The effect of the formula is very small.              |
| Ease of application | 0,253                     | F(2,58)<br>= 9,667 | <<br>0,001     | 0,250                      | There are significant differences between the formulas. The effect is medium to large.                         |

7. Irritation test

Irritation test results showed that all formulas, including those containing watermelon seed extract, did not cause erythema (redness) or edema (swelling).

This proves that the formulation is safe and does not cause skin irritation. This safety is supported by the natural emollient content in the extract, which effectively maintains moisture without side effects<sup>12</sup>.

**Table 8.** Irritation Test Results

| Formula | Reaction       |                |
|---------|----------------|----------------|
|         | Errythema      | Edema          |
| (1)     | (2)            | (3)            |
| F0      | non-irritating | non-irritating |
| F1      | non-irritating | non-irritating |
| F2      | non-irritating | non-irritating |
| F3      | non-irritating | non-irritating |

**Lip Balm Stability Evaluation**

The stability evaluation of the lip balm preparation was carried out over 10 weeks, showing very good results, as all formulas maintained their organoleptic characteristics and homogeneity without any significant changes in color, rancid odor, or phase separation.

pH re-testing after 10 weeks showed good pH stability, with all formulas remaining within a safe range for lip skin. Although F1 and F2 had a slight increase in pH, their values are still acceptable for lip balm products. This stability indicates strong formulation integrity.

**Table 9.** LipBalm Stability Evaluation Results

| Test parameter | Formula    | 1 week                                                                                          | 10 week                                                                                                      |
|----------------|------------|-------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------|
| Organoleptic   | F0         | Semi-solid with a cloudy white color, this preparation has the characteristic aroma of beeswax. | Its semi-solid consistency has a pale yellow to bright yellow color with a stable aroma, resembling beeswax. |
|                | F1, F2, F3 | Having a semi-solid texture, this substance is pale yellow to bright yellow in color with a     | ranging from pale yellow to bright yellow, with a semi-solid texture and a slightly                          |

|             |                |                       |                               |
|-------------|----------------|-----------------------|-------------------------------|
|             |                | slightly nutty aroma. | nutty aroma.                  |
| homogeneity | F0, F1, F2, F3 | homogeneous           | Homogeneous                   |
| pH          | F0, F1, F2, F3 | 6.50 - 6.86           | F1(7.28), F2(7.42), F3(6.86). |

#### 4. CONCLUSION

Based on the research conducted, watermelon seed extract (*Citrullus lanatus*) was successfully formulated into a lip balm preparation that met the required physical evaluation standards, including organoleptic stability, homogeneity, appropriate pH, melting point, and satisfactory spreadability. All formulations were proven to be safe, as no signs of erythema or edema were observed during the irritation test, indicating that the preparations were non-irritating. Statistical analysis demonstrated that increasing the concentration of watermelon seed extract significantly enhanced the moisturizing effect and ease of application of the lip balm. Among all formulations, formula F3 containing 9% extract was the most preferred by panelists, particularly in terms of moisture improvement and user comfort. Meanwhile, the extract concentration did not significantly influence the aroma or the physical form of the preparation. Overall, this research concludes that watermelon seed extract has strong potential as an active natural ingredient for producing lip balm products that are safe, stable, effective, and well-accepted by consumers, with the optimal performance observed at a concentration of 9%.

#### 5. REFERENCES

- Hidayah F, Erwiyani AR. Knowledge level, attitudes, and use of lip balm for lip care among pharmacy students at Ngudi Waluyo University. *Pro Heal J Ilm Kesehat*. 2022;4(1):179-183. doi:10.35473/proheallth.v4i1.1553
- Nazliniwaty, Laila L, Wahyuni M. Utilization of pomegranate peel extract (*Punica granatum* L.) in lip balm formulation. *J Jamu Indones*. 2019;4(3):87-92. doi:10.29244/jji.v4i3.153
- Adjeng ANT, Koedoes YA, Ali NFM, Palogan ANA, Damayanti E. Education on safe cosmetic ingredients and usage in Suka Banjar Village, Gedong Tataan, Pesawaran Regency. *J Kreat Pengabd Kpd Masy*. 2023;6(1):89-102. doi:10.33024/jkpm.v6i1.8041
- Handayani R. Lip balm formulation from Arabica coffee seed extract (*Coffea arabica* L.) Java Preanger as an emollient. *J Ilm Farm Farmasyifa*. 2021;4(1):105-111. doi:10.29313/jiff.v4i1.6497
- Amaliah Dahlia A, Pratama M. Antioxidant activity test of ethanol extract of watermelon rind and seeds (*Citrullus lanatus*) using the DPPH (1,1-diphenyl-2-picrylhydrazyl) method. *Makassar Nat Prod J*. 2023;1(2):2023-2102.
- Irabor GE, Ebhoaye JE, Odia A. Qualitative and quantitative screening of phytochemical compounds in watermelon (*Citrullus lanatus*) seeds cultivated in Esan West Local Government Area, Edo State. *Int J Eng Appl Sci Technol*. 2020;5(1):268-273. doi:10.33564/ijeast.2020.v05i01.040
- Okzelia SD. Formulation and evaluation of gel from white watermelon rind extract (*Citrullus lanatus* [Thunb.] Matsum & Nakai) as a skin moisturizer. *J Sabdariffarma*. 2022;9(2):33-44. doi:10.53675/jsfar.v3i2.394
- Rahmadini WS. *Formulation and evaluation of lip balm stick containing white watermelon rind extract (Citrullus lanatus) with variations in cera alba concentration*. Thesis. Politeknik Kesehatan Palembang; 2022.
- Simamora F, Janice J, Amir W. Effectiveness test of white rind extract cream from red watermelon (*Citrullus lanatus*) on increasing elasticity, sebum, and hydration in white mice (*Mus musculus*) skin. *Int J Heal Pharm*. 2022;3(3):322-330. doi:10.51601/ijhp.v3i3.173
- Harahap NH. *Lip balm formulation containing pumpkin seed oil as a lip moisturizer*. Undergraduate thesis. Universitas Sumatera Utara; 2020.

- 
11. Ambari Y, Hapsari FND, Ningsih AW, Nurrosyidah IH, Sinaga B. Formulation study of lip balm containing sappan wood extract (*Caesalpinia sappan* L.) with beeswax variation. *J Islam Pharm.* 2020;5(2):36-45. doi:10.18860/jip.v5i2.10434
  12. Islamiah N, Sukrasono, Simanullang G. Formulation and physical stability evaluation of lip balm containing rice bran oil. *Media Farm Indones.* 2023;18(2). doi:10.53359/mfi.v18i2.230
  13. Amalia I, Prabandari S, Susiyarti. Formulation and physical property evaluation of lip balm containing strawberry ethanol extract (*Fragaria* sp). *Politek Harapan Bersama.* 2021;1(9):3.
  14. Kamal S, Surya EMK. Lip balm formulation using gambir catechin (*Uncaria gambir* Roxb.) and hedonic evaluation. *Glob Conf Ser Sci Technol.* 2020;5:33-38.
  15. Ardini D, Sumardilah SD. Effects of aloe vera extract lip balm as a lip moisturizer. *J Kesehat Metro Sai Wawai.* 2021;14(1):10-18. doi:10.26630/jkm.v13i1.2677
  16. Henson RN. Analysis of variance (ANOVA). In: *Encyclopedia of Psychopharmacology*. Vol 1. Elsevier Inc; 2015. doi:10.1016/B978-0-12-397025-1.00319-5
  17. Lailatul S, Yanti EF. Formulation of lip balm preparation containing red dragon fruit extract. 2024;7(1):1-11.