



## Relationship of Soil Physicochemical Properties and Existence of *Phytophthora* sp. in Pineapple Plantations

Deva Aziz Nanda Martin\*<sup>1</sup> and Ali Rahmat<sup>2</sup>

<sup>1</sup>Department of Agrotechnology, Faculty of Agriculture Science, University of Lampung, Jln. Sumantri Brojonegoro No. 01, Bandar Lampung, Lampung 35141, Indonesia

<sup>2</sup>Unite Graduate School of Agriculture Science, Gifu University, 1-1 Yanagido, Gifu 501-1193, Japan

\*Correspondence: e-mail: devan369martin@gmail.com

### ABSTRACT

One of the tropical fruits with high economic value is pineapple. Pineapple plantation has several limitations, especially in cultivating period and diseases caused by *Phytophthora* sp. (called heart rot disease). The spread of *Phytophthora* sp. in pineapple plantation is influenced by physical of soil (called as soil borne pathogen). Here, the purpose of this study was to examine physical properties of soil (compaction), specifically on heart rot disease in pineapple plantation. We used pineapple farm locating in Central Lampung, Indonesia. In this paper, we analyzed the disease impact from the soil compaction point of view. We investigated the difference between soil samples with and without sick plant. The results showed that the disease area were affected by *Phytophthora* sp. area have high soil compaction. Indeed, this soil condition is good for growing *Phytophthora* sp.

© 2017 Tim Pengembang Journal UPI

### ARTICLE INFO

Article History:

Submitted/Received 14 Dec 2016

First revised 01 Jan 2017

Accepted 30 Mar 2017

First available online 01 Apr 2017

Publication date 01 Apr 2017

**Keyword:**

*Phytophthora* sp.,

*Pathogen*,

*Soil compaction*,

*Pineapple plantation*,

*Heart rot*.

## 1. INTRODUCTION

Indonesia is the best country for production fruits and vegetable. (Prasetyo, 2016). One of the fruits with a large area and high production in Indonesia is Pineapple (*Ananas comosus* Merr.). This makes pineapple to be the most important economic plants in the tropic area. Pineapple is grown mainly for fresh, canned fruits and juice. This fruit is also the source of bromelain and enzyme used in pharmaceuticals (Pornsuriya *et al.*, 2008).

In the cultivation process, heart rot symptoms are commonly found in the fields and become major losses in pineapple production. Heart rot is caused by *Phytophthora* sp. (Bortholomew, 2003). Soil-borne disease caused by *Phytophthora* sp. species is serious in the tropics because of frequent heavy rainfall favorable to the spread of sporangia and zoospores as well as a lack of cold temperature in the winter to kill the pathogens (Ann, 1994). Some chemical treatments were typically used to control heart rot disease, but they make soil to be getting worse and unsustainable for the future. (Muhammed *et al.*, 2014)

As soil-borne pathogens, soil properties influence the existence of *Phytophthora* sp. (Azcón-Aguilar & Barea, 1997) In this experiment, we investigated effect of soil physicochemical properties on *Phytophthora* sp. in the pineapple plantation. The research of this research can give information how to manage and make policy in the cultivation system of pineapple.

## 2. MATERIALS AND METHODS

The experiment was conducted Pineapple plantation located in Central Lampung, Lampung Province, Indonesia. The

experiment was done with the following three steps; First, field survey and take a soil sample (location was indicated an attack by *Phytophthora* sp. and without *Phytophthora* sp.). Second, analysis of the existence of *Phytophthora* spp. from the soil sample following Postulate Koch in biotechnology laboratory of the University of Lampung. Third, analysis of soil properties (soil compaction) in soil science laboratory of the University of Lampung. The experiment was conducted from December 2015 to February 2016. The data was collected and tested using a T-test.

## 3. RESULTS AND DISCUSSION

The survey results showed that the soil are classified as the attacked area by *Phytophthora* sp. based on the symptom of heart rot disease. We found pineapple infected by heart rot disease has the average percentage of 26.85%. Furthermore, in the same location, we took soil for sampling, namely a sick plant (soil Phy) and healthy plant (soil No-Phy).

### 3.1. Disease Identification

The symptom of heart rot disease in this study is identified by the colour of the heart leaves. Leaves changes to yellow or light coppery brown. Later, the that leaves (causing the leaf edges to roll under) turn into brown and eventually die. Once symptom is detected, young leaves are easily pulled from the plant, and the basal white leaf tissue at the base of the leaves become water-soaked and rotten with a foul smell. This is due to the invasion of secondary organisms. The growing point of the stem becomes yellowish-brown with a draks line between healthy and diseased areas (Figure 1).

The results of the isolation and identification of the disease from infected leaves of pineapple plants by *Phytophthora* sp. are shown in **Figure 2**. As shown in this figure, images of spores of *Phytophthora* sp. was isolated from leaves pineapple. This result is in a good agreement with other report. (Omane *et al.*, 2012)


### 3.2. Soil Properties

Compaction is rearrangement of aggregates and primary particles into a state of higher bulk density-and lower porosity

when a load (or stress, pressure) is applied to a soil (Allmaras *et al.*, 1988; Fox & Kamprath, 1970). The first impact of compaction is the loss of pore space between aggregates (interaggregate pore space) as the soil volume decreased. The loss of interaggregate pore space has a major effect on water infiltration and drainage, gas exchange and aeration (oxygen diffusion), mechanical resistance to root penetration and proliferation, heat movement, and biological activity of both soilborne pathogens and the host organism (Allmaras *et al.*, 1988)




**Figure 1.** Symptom of heart rot disease caused by *Phytophthora* sp. in pineapple plantation in Indonesia. Figure in the left is the low magnification of picture, whereas figure in the right is the high magnification.



**Figure 2.** Microscope images of spores (left) and hyphae (right) of *Phytophthora* sp. was isolated from leaves pineapple.

Problems were induced by excessive compaction are runoff, soil erosion, slow infiltration and soil crusting, impaired or delayed internal drainage, decreased soil water storage, shallow and sparse plant rooting, reduced nutrient and water uptake, accelerated denitrification, production of toxic materials due to soil reducing conditions, fewer field days, and more root disease from pathogens such as *Pythium*, *Phytophthora*, *Aphanomyces*, *Fusarium* and *Rhizoctonia* spp. Most of these problems are linked to adverse water relations caused by retarded infiltration, less airfilled pore space, and impeded water movement (Allmaras *et al.*, 1988). Soil compaction alters soil microbial structure and function as a result of the limitation of air permeability and oxygen availability, which has implications for soil nutrition and soil-borne disease (Ishak *et. al.*, 2016).

Results showed that there is an impact from the soil compaction (See **Figure 3**). Soil compaction in a land with an attack by *Phytophthora* sp. is higher than that without *Phytophthora* sp. on T-test. Based on **Figure 3** on T-test, there is no significant difference for the value of the soil compaction at depth 10-50 cm. The tillage is done only to a depth of 50 cm, and the pineapple roots grow to a depth of 50 cm (Gunawan, 2007). However, soil compaction at a depth of 60-70 cm is significantly different. Phy-soil sample has higher soil compaction than that in No-Phy soil sample. Thus, when there is a rain, soil in No-Phy condition can infiltrate more water. However, because of soil compaction on soil in Phy-soil, puddle can be promoted. Puddles allow sporangium formation of *Phytophthora* sp. if mycelium soaked by water.



**Figure 3.** Soil compaction.

Soil No-Phy sample has possibility higher soil porosity than that in Phy-soil sample. Thus, when there is a rain, or irrigation, soil for no-Phy sample can absorb and hold more water than that for Phy-soil sample. Low soil porosity and high soil compaction promote puddle. The rain water will destroy the soil aggregates, while the water will splash the soil particles and are transported by soil surface runoff. This phenomenon is known as erosion (Soepardi, 1983). Erosion carries soil particles or organic materials, which contain structure of *Phytophthora* sp. (Weste & Taylor, 1971). This condition makes *Phytophthora* sp. to spread rapidly in the plantation area. (Weste, 1975)

Excessive soil compaction affects the rate and distribution of root growth, and thus affects the chances of successful host-to-pathogen contact and the dynamics of root-pathogen interactions. For example, poor aeration or axial constraints on root development (such as mechanical resistance) may reduce the rate of root tip

advance by as much as 75% and may also induce formation of laterals much closer to the root tip. The result is a more compacted and stressed root system. Exudates produced by elongating and/or stressed roots stimulate dormant fungal pathogen structures (e.g. chlamydospores, microsclerotia) to germinate and grow. The overall result is that soilborne pathogens in compacted soils are more likely to intercept young lateral root and obtain sufficient nutrients to infect that root. (Allmaras *et al.*, 1988)

#### 4. CONCLUSIONS

Based on research, we concluded that there is a correlation between soil physicochemical properties and existence of *Phytophthora* sp. *Phytophthora* sp. can grow with maximum growth rate in high soil compaction. This soil condition support its growth because of its low porosity, low infiltration, and easy to be flooding. This study gives information on how to treat the *Phytophthora* sp. in the plantation.

## 5. ACKNOWLEDGEMENTS

The authors thank to University of Lampung, Indonesia and Gifu University, Japan.

## 6. AUTHORS' NOTE

The author(s) declare(s) that there is no conflict of interest regarding the publication of this article. Authors confirmed that the data and the paper are free of plagiarism.

## 7. REFERENCES

Allmaras, R. R., Kraft, J. M., & Miller, D. E. (1988). Effects of soil compaction and incorporated crop residue on root health. *Annual review of phytopathology*, 26(1), 219-243.

Ann, P. J. (1994). Survey of soils suppressive to three species of *phytophthora* in taiwan. *Soil biology and biochemistry*, 26(9), 1239-1248.

Azcón-Aguilar, C., & Barea, J. M. (1997). Arbuscular mycorrhizas and biological control of soil-borne plant pathogens—an overview of the mechanisms involved. *Mycorrhiza*, 6(6), 457-464.

Bartholomew, D. P., Paull, R. E., & Rohrbach, K. G. (2002). *The pineapple: botany, production, and uses*. CABI. Wallingford, UK

Fox, R. L., & Kamprath, E. J. (1970). Phosphate sorption isotherms for evaluating the phosphate requirements of soils. *Soil science society of america journal*, 34(6), 902-907.

Gunawan, E. (2007). *Kajian pertumbuhan dan produksi nenas pada lahan gambut dan lahan aluvial di kalimantan barat*. TesisIPB: Bogor. (Master thesis, Indonesian language)

Ishak, L., McHenry, M.T. & Brown, P.H. (2016). Soil compaction and its effects on soil microbial communities in Capsicum growing soil. *Acta horticultura*, 1123, 123-130

Omame, E., Oduro, K. A., Cornelius, E. W., Opoku, I. Y., Akrofi, A. Y., Sharma, K., & Bandyopadhyay, R. (2012). First report of leaf blight of taro (*Colocasia esculenta*) caused by *Phytophthora colocasiae* in Ghana. *Plant disease*, 96(2), 292-292.

Prasetyo, E. (2016). Detection of mangotree varieties based on image processing. *Indonesian journal of science and technology*, 1(2), 203-215.

Pornsuriya, C., Wang, H.K., Lin, F.C., & Soytong, K. (2008). First report of pineapple root rot caused by *phythium graminicola*. *Journal of agriculture technology*, 4(1), 139-150.

Soepardi, G. (1983). *Sifat dan Ciri Tanah*. Departemen Ilmu-ilmu Tanah Fakultas Pertanian IPB: Bogor. (Indonesian)

Weste, G. (1975). The distribution of *phytophthora cinnamomy* within the national park, wilson promontory. *Australian journal of botany*, 23(1), 67-76.

Weste, G., & Taylor, P. (1971). The invasion of native forest by *Phytophthora cinnamomy* 1. Brisbane Ranges. *Australian journal of botany*, 19(2), 281-294.